



<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

## TOS PROTOCOL AND PROCEDURE: PLANT DIVERSITY SAMPLING

<b>PREPARED BY</b>	<b>ORGANIZATION</b>	<b>DATE</b>
David T. Barnett	FSU	11/17/2015

<b>APPROVALS</b>	<b>ORGANIZATION</b>	<b>APPROVAL DATE</b>
Mike Stewart	PSE	1/25/2016
Andrea Thorpe	PROJ SCI	1/27/2016

<b>RELEASED BY</b>	<b>ORGANIZATION</b>	<b>RELEASE DATE</b>
Anne Balsley	CM	1/27/2016

See configuration management system for approval history.

© 2016 NEON Inc. All rights reserved.

The National Ecological Observatory Network is a project solely funded by the National Science Foundation and managed under cooperative agreement by NEON, Inc. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

## Change Record

REVISION	DATE	ECO #	DESCRIPTION OF CHANGE
A_DRAFT	10/03/2011	ECO-00280	Draft release
B_DRAFT	01/13/2014	ECO-01140	Draft release. Will be finalized in next rev.
C	03/18/2014	ECO-01668	Production release, template change, and other changes as detailed in Appendix C (rev C only)
D	11/03/2014	ECO-02341	Migration to new protocol template
E	02/24/2015	ECO-02536	Naming convention for subplots was changed (see Figure 1). Enter 0.5 for estimates of cover <1%.
F	1/27/2016	ECO-03451	<ul style="list-style-type: none"> <li>-Removed directive to record species height.</li> <li>-Changed directions for recording cover to include all vegetative cover under 3m regardless of height of individual.</li> <li>-Added the Tower Base Plots to the description of which plot types to sample.</li> <li>-Updated sample timing and bout number information (Appendix E).</li> <li>-Changed definition of 'standing dead' to include woody species</li> <li>-More clearly defined 'wood'</li> </ul>

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

**TABLE OF CONTENTS**

**1 OVERVIEW .....1**

1.1 Background ..... 1

1.2 Scope..... 1

1.2.1 NEON Science Requirements and Data Products ..... 1

1.3 Acknowledgments..... 1

**2 RELATED DOCUMENTS AND ACRONYMS .....2**

2.1 Applicable Documents ..... 2

2.2 Reference Documents..... 2

2.3 Acronyms ..... 2

2.4 Definitions..... 2

**3 METHOD .....3**

**4 SAMPLING SCHEDULE .....4**

4.1 Sampling Frequency and Timing ..... 4

4.2 Criteria for Determining Onset and Cessation of Sampling ..... 4

4.3 Timing for Laboratory Processing and Analysis ..... 4

4.4 Sampling Timing Contingencies ..... 5

**5 SAFETY .....5**

**6 PERSONNEL AND EQUIPMENT.....6**

6.1 Equipment..... 6

6.2 Training Requirements..... 10

6.3 Specialized Skills..... 10

6.4 Estimated Time ..... 10

**7 STANDARD OPERATING PROCEDURES.....11**

**SOP A PREPARING FOR SAMPLING.....11**

**SOP B FIELD SAMPLING .....11**

**SOP C LABORATORY PROCESSING AND ANALYSES.....22**

**SOP D DATA ENTRY AND VERIFICATION .....24**

**SOP E SAMPLE SHIPMENT .....25**

**8 REFERENCES .....26**

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

**APPENDIX A DATASHEETS .....26**

**APPENDIX B QUICK REFERENCES .....27**

**APPENDIX C CHECKLISTS .....33**

**APPENDIX D REMINDERS .....34**

**APPENDIX E ESTIMATED DATES FOR ONSET AND CESSATION OF SAMPLING .....35**

**APPENDIX F SITE-SPECIFIC INFORMATION.....38**

**LIST OF TABLES AND FIGURES**

**Table 1.** Equipment list – Materials and supplies required for one crew for the plant sampling procedure ..... 6

**Table 2.** Equipment list – Laboratory processing ..... 9

**Table 3.** Variables to be observed in the 1-m<sup>2</sup> nested subplot. .... 13

**Table 4.** Identification qualifier codes (idQ) to designate unknown species or those species with uncertain identification in the field or after identification in the lab..... 20

**Table 5.** Datasheets associated with this protocol ..... 26

**Table 6.** Estimated dates of historical temperature thresholds ..... 35

**Figure 1.** The square, multi-scale plot ..... 12

**Figure 2.** Estimates of cover should include all vegetative material < 3m in height. For herbaceous growth (A), and shrubs (B) < 3m, record the total combined cover by species; for tall trees with no woody branches or foliar growth < 3m (C) record basal area and the presence of individuals > 3m should be noted for that species; for trees (D) and shrubs (E) > 3m that also have vegetative growth < 3m, record the cover of vegetative growth < 3m and indicate the presence of individuals > 3m in height..... 16

**Figure 3.** The 1 m<sup>2</sup> subplot is calibrated with black and white marks to make estimates of plant species cover more accurate and repeatable..... 18

**Figure 4.** The plot will have some permanent markers and will also require temporary flags that are placed each time the plot is measured..... 29

**Figure 5.** Assembled vegetation sampling subplot frame ..... **Error! Bookmark not defined.**

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

## **1 OVERVIEW**

### **1.1 Background**

The purpose of plant diversity sampling is to describe inter- and intra-annual variation of the presence and abundance of plant species at NEON sites. In addition to providing habitat for local fauna, the flora at each site integrates a variety of biotic and abiotic factors that respond to environmental change.

Plant species diversity will be measured once or twice annually in the field. The plot-based method yields plant species data at multiple scales that will provide an understanding of changes in composition, distribution, and abundance of native and non-native plant species. The data will be comparable within and across NEON sites and to other continental vegetation efforts to allow for a comprehensive understanding of the impacts of the drivers of change on the diversity of plant species and the functional role they play in ecological systems.

This document provides detailed guidance for assessing plant diversity in the Distributed Base Plots and Tower Base Plots in the field, the collection and handling of unknown plant species, and the collection of voucher specimens for training and archiving purposes.

### **1.2 Scope**

This document provides a change-controlled version of Observatory protocols and procedures. Documentation of content changes (i.e. changes in particular tasks or safety practices) will occur via this change-controlled document, not through field manuals or training materials.

#### **1.2.1 NEON Science Requirements and Data Products**

This protocol fulfills Observatory science requirements that reside in NEON’s Dynamic Object-Oriented Requirements System (DOORS). Copies of approved science requirements have been exported from DOORS and are available in NEON’s document repository, or upon request.

Execution of this protocol procures samples and/or generates raw data satisfying NEON Observatory scientific requirements. These data and samples are used to create NEON data products, and are documented in the NEON Scientific Data Products Catalog (RD[03]).

### **1.3 Acknowledgments**

This protocol is based on a technique for sampling plant species diversity in a multi-scale plot that was created for use in The Carolina Vegetation Survey, the Whittaker, and the Modified-Whittaker plot design. Special thanks belong to Ben Chemel, Tom Stohlgren, Geneva Chong, and Robert Peet.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

## 2 RELATED DOCUMENTS AND ACRONYMS

### 2.1 Applicable Documents

Applicable documents contain higher-level information that is implemented in the current document. Examples include designs, plans, or standards.

AD[01]	NEON.DOC.004300	EHS Safety Policy and Program Manual
AD[02]	NEON.DOC.004316	Operations Field Safety and Security Plan
AD[03]	NEON.DOC.000724	Domain Chemical Hygiene Plan and Biosafety Manual
AD[04]	NEON.DOC.050005	Field Operations Job Instruction Training Plan
AD[05]	NEON.DOC.000912	NEON Science Design for Plant Diversity
AD[06]	NEON.DOC.014051	Field Audit Plan
AD[07]	NEON.DOC.001399	NEON Raw Data Ingest Workbook for TOS Plant Diversity

### 2.2 Reference Documents

Reference documents contain information that supports or complements the current document. Examples include related protocols, datasheets, or general-information references.

RD[01]	NEON.DOC.000008	NEON Acronym List
RD[02]	NEON.DOC.000243	NEON Glossary of Terms
RD[03]	NEON.DOC.002652	NEON Level 1, Level 2 and Level 3 Data Products Catalog
RD[04]	NEON.DOC.001271	NEON Protocol and Procedure: Manual Data Transcription
RD[05]	NEON.DOC.001579	Datasheets for TOS Protocol and Procedure: Plant Diversity Sampling
RD[06]	NEON.DOC.001024	TOS Protocol and Procedure: Canopy Foliage Chemistry and Leaf Mass Per Area Measurements
RD[07]	NEON.DOC.014040	TOS Protocol and Procedure: Plant Phenology
RD[08]	NEON.DOC.000987	TOS Protocol and Procedure: Measurement of Vegetation Structure
RD[09]	NEON.DOC.001025	TOS Protocol and Procedure: Plot Establishment
RD[10]	NEON.DOC.001702	NEON Herbarium Specimen Label and Annotation Generation
RD[11]	NEON.DOC.001237	NEON Theoretical Basis Document: TOS Plant Diversity – QA/QC of Raw Field And Lab Data
RD[12]	NEON.DOC.003564	NEON Standard Operating Procedure: Plant Voucher Specimen Preparation

### 2.3 Acronyms

All acronyms used in this document are defined in RD[01].

### 2.4 Definitions

N/A

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

**3 METHOD**

This document describes the collection of plant diversity information designated to inform the objectives and associated requirements of the National Ecological Observatory Network (NEON). Plant diversity sampling shall occur according to a sample design – a statistically rigorous system that directs the spatial distribution of observations – at plots distributed across NEON sites. Plant species composition or presence and abundance data shall be collected in multi-scale plots, estimates of cover being limited to 1m<sup>2</sup> subplots that shall be nested in larger plots where plant species composition will be recorded.

Even experienced botanists will not know every species encountered in each plot. Typically it is not cost effective, and sometimes impossible, to spend time identifying a plant in the field. Therefore, instructions for the collection and identification of unidentified species are provided.

Voucher specimens provide a permanent record of the NEON naming convention, use of authorities, validation, and a means to track taxonomic naming conventions through time. The samples must be of archival quality. Specimens should be collected when floristic parts are present. Twenty to forty of the common species found in plots should be collected over the first several years of sampling. Vouchers to be housed at the Domain Support Facility should be dried, pressed and mounted. Vouchers to be sent to archive facility must meet herbaria standards, and should be dried, and pressed. Plant tissue will also be collected and stored at an archive facility to enable the ecological community to conduct a variety of genetic investigations (RD[06]).

Standard Operating Procedures (SOPs), in Section 7 of this document, provide detailed step-by-step directions, contingency plans, sampling tips, and best practices for implementing this sampling procedure. The value of NEON data hinges on consistent implementation of this protocol across all NEON domains, for the life of the project. It is therefore essential that field personnel carry out this protocol as outlined in this document. In the event that local conditions create uncertainty about carrying out these steps, it is critical that technicians document the problem and enter it in NEON’s problem tracking system.

The procedures described in this protocol will be audited according to the Field Audit Plan (AD[06]). Additional quality assurance will be performed on data collected via these procedures according to the NEON Algorithm Theoretical Basis Document for TOS Plant Diversity: QA/QC of Raw Field and Lab Data (RD[11]).

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

## 4 SAMPLING SCHEDULE

### 4.1 Sampling Frequency and Timing

The timing of field sampling will be determined annually by Field Operations based on criteria for sampling and approved annually by Science (Appendix E).

Complete sampling in approximately a 1-2 month period around peak flowering (Appendix E). Significant delays may change the detectability of species and influence the comparability of sampling bouts.

### 4.2 Criteria for Determining Onset and Cessation of Sampling

Sample bouts will be timed to maximize the number of plant species detected at a NEON site. Observations will generally be made at peak phenology (when most species are flowering) to facilitate identification of individuals. Many NEON sites will not have a single peak in phenology due to plant adaptations that take advantage of different climatic conditions. Sites with more than one peak may require multiple sampling bouts (Appendix E).

The specific timing of sampling bouts will be tied to observations from the NEON phenology measurements (RD[07]), but the timing of sampling has generally been determined to support planning purposes (Appendix E).

### 4.3 Timing for Laboratory Processing and Analysis

There are four conditions that require the collection of plant specimens that result in lab activities:

1. Identification at the Domain Support Facility. These species could not be efficiently identified in the field, were collected and will be identified by NEON staff. If possible, and to increase efficiency of field sampling, identify the species and soon after the collection, update morphospecies on the datasheet or in the mobile data recorder if the data has not been uploaded or on the appropriate space in the internal NEON documents.
2. Inclusion at Domain Support Facility herbarium. These specimens should be pressed, dried, identified, mounted and labeled. A quality specimen might require the collection of two individuals should identification require destruction of the sample (e.g., flower and/or ovary dissection).
3. Identification by external expert. These specimens will be pressed and dried and prepared for shipment following the field season.
4. Archive at external facility. These specimens should be pressed, dried, and identified if possible.

Numbers three and four above are not mutually exclusive. Some specimens will be sent for identification and then sent to an archive facility (RD [12]).

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

Specimens should not be left in the refrigerator for more than two days. They can be placed in a press, stored in a well-ventilated location, and identified at a later date. Identification often requires a variety of dichotomous keys, a dissecting microscope, a dissecting kit, and a herbarium with voucher specimens for verification. Any specimen destined for an archive should be placed in the -80C freezer for 24 hours after it is completely dried for decontamination (RD [12]).

#### 4.4 Sampling Timing Contingencies

When unexpected field conditions require deviations from this protocol, the following field implementation guidance must be followed to ensure quality standards are met:

- If the data collection cannot be completed due to safety or logistical reasons, sampling must resume at the plot during the same sampling bout (approximately 45 days) if the plot is to be considered complete. Delay of sampling should be recorded in data about the plot-sampling effort.
- Any changes that the plot undergoes should be noted in the data associated with the plot. For example, disturbance caused by a large tree falling in the plot, severe flood damage and erosion, or the disturbance from mammals – bison, cows, small mammals – during (between visits to a single plot) or between bouts should be noted.
- Deviations associated with the collection of data should not be made from this protocol. The number of people collecting data, the tools for defining the plot boundary, and the amount of material collected for the identification of unknown plant species may be altered to meet the needs of Operational constraints.

## 5 SAFETY

This document identifies procedure-specific safety hazards and associated safety requirements. It does not describe general safety practices or site-specific safety practices.

Personnel working at a NEON site must be compliant with safe field work practices as outlined in the Operations Field Safety and Security Plan (AD[02]) and EHS Safety Policy and Program Manual (AD[01]). Additional safety issues associated with this field procedure are outlined below. The Field Operations Manager and the Lead Field Technician have primary authority to stop work activities based on unsafe field conditions; however, all employees have the responsibility and right to stop their work in unsafe conditions.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

## 6 PERSONNEL AND EQUIPMENT

### 6.1 Equipment

The following equipment is needed to implement the procedures in this document. Equipment lists are organized by task. They do not include standard field and laboratory supplies such as charging stations, first aid kits, drying ovens, ultra-low refrigerators, etc.

**Table 1.** Equipment list – Materials and supplies required for one crew for the plant sampling procedure

Item No.	R/S	Description	Purpose	Quantity*	Special Handling
<b>Durable Items</b>					
MX104361	R	Chaining pins or other suitable anchor	Anchor measuring tapes	3	N
	S	Cooler	Chill perishable plant vouchers in field	5	N
MX100696	R	Digital camera and SD card, 12 megapixel	Capture images of plants for species identification	1	N
MX100703	R	GPS receiver, recreational accuracy	Navigate to sampling location	1	N
MX100358	R	Ice pack	Chill perishable plant vouchers in field	Many	N
MX103211	R	Magnifier hand-lens, 20X	Aid in species identification	Many	N
MX104369	R	Measuring tape, minimum 50 m	Determine plot boundary	3	N

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

Item No.	R/S	Description	Purpose	Quantity*	Special Handling
	R	Pruning shear	Collect voucher specimens	1 ea.	N
	R	Sampling frame, 1m <sup>2</sup>	Delineate 1m <sup>2</sup> subplot	1	N
GN10001	S	Small carabiner and ring binder	Organize and carry unknown plant vouchers	1	N
	R	Weeder	Collect voucher specimens	1	N
<b>Consumable items</b>					
	S	AA battery	Spare battery for GPS receiver		
	R	Adhesive label	Label unknown and voucher specimens	1 sheet	N
		All weather copy paper	Print datasheets		
	S	Digital camera battery	Spare battery		
	R	Field notebook	Record field notes	1	N
MX103940	S	Flagging tape	Delineate sampling area	1	N
MX100592	R	Resealable plastic bag, 1 gal	Organize and carry unknown plant vouchers and plant tissue	> 40	N
	R	Survey marking flag, PVC or fiberglass stake	Delineate sampling area	Many	N

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

Item No.	R/S	Description	Purpose	Quantity*	Special Handling
		Tabloid newspaper pages	Press collected individuals for identification		
<b>Resources</b>					
RD[05], RD[06]	R	Field datasheet	Record data	1	N
	S	Field guide, regional flora reference guide and/or key	Identify unknown species	1	N
	S	Field guide, species list	Identify unknown species	1	N

R/S=Required/Suggested

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

**Table 2.** Equipment list – Laboratory processing

Item No.	R/S	Description	Purpose	Quantity*	Special Handling
<b>Durable Items</b>					
MX103213	R	Botany dissection kit	Identify unknown species	1	N
MX103214	R	Cardboard ventilator	Pressing plants		
MX102896	R	Microscope	Aid in species identification	1	N
MX100312	R	Paper blotters	Press collected individuals for identification	Many	N
MX100316	R	Plant press	Press collected individuals for identification	2	N
	R	Scissors or pruning shear	Prepare voucher specimen for mounting	1 ea.	N
<b>Consumable items</b>					
		Tabloid newspaper pages	Press collected individuals for identification		
<b>Resources</b>					
	R	Field guide, regional flora reference guide and/or key	Identify unknown species		N

R/S=Required/Suggested

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

**6.2 Training Requirements**

All technicians must complete required safety training and protocol-specific training for safety and implementation of this protocol as required in Field Operations Job Instruction Training Plan (AD[04]).

**6.3 Specialized Skills**

A minimum of two technicians is required for each plant diversity sampling team. It is mandatory that one technician have experience with the identification of plants – preferably in the habitats found at the site where observations will be made, be able to use a dichotomous key, and have experience identifying plant specimens in the lab with a dissecting microscope and associated tools. At each site this technician must be able to identify most of the species in the field.

It is important that careful attention be paid to the person responsible for identifying plants in the plot. This is important because unknown plant species collected during the effort will be tracked by this identity (measuredBy) and the unknown plant name. If multiple botanists are identifying species within a single plot, a single botanist should be responsible for collecting unknown species. Alternatively, botanists can be responsible for all of the species in a 100m<sup>2</sup> subplot where the botanist (measuredBy) can be recorded on the datasheet or handheld device.

**6.4 Estimated Time**

A plot should take 1-5 hours for a team of two to complete. The time required will vary depending on a number of factors, such as skill level, species richness at the site, and environmental conditions. The timeframe is an estimate based on completion of a task by a skilled two-person team (i.e., not the time it takes at the beginning of the field season). Use this estimate as framework for assessing progress and see the sections about searching the 10m<sup>2</sup> and 100m<sup>2</sup> subplots for more guidance. If a task is taking significantly longer than the estimated time, a problem ticket should be submitted.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

## 7 STANDARD OPERATING PROCEDURES

### SOP A Preparing for Sampling

Assemble nested subplot frames

- The cover and identity of plant species will be recorded in 1m<sup>2</sup> frames.

Prepare data collection tools

- Prepare the mobile data recorder and datasheets (RD[05]) for collecting data prior to leaving for the field. Be sure electronics are charged, software is loaded, and data from prior sampling efforts have been downloaded and deleted. Be prepared to use provided paper datasheets if the electronic device fails (e.g. dunked in a creek, lost, or crashes) or if data is to be collected on paper data sheets.
- Plant species identified in the field will be recorded according to the USDA PLANTS database code. These codes will be downloaded onto the hand-held device, but having a printed version of species found at the site during previous sampling years and the associated code is strongly recommended given the importance of tracking species by the correct codes. The Domain-specific plant lists and codes are also available on the internal NEON internal document locations. The quality of the data depends on the correct species-code linkages.

Organize equipment and consumable items

- Plastic bags will be used to collect unknown plant species. Prior to going to the field be sure to have an ample number of loose bags. Adhesive labels will be needed and working permanent markers and pencils.


### SOP B Field Sampling

#### B.1 Plot Establishment

Plant diversity sampling occurs in a square-shaped plot measuring 20m on a side and containing four 100m<sup>2</sup> subplots (Figure 1). Each subplot contains nested subplots: a 1m<sup>2</sup> subplot nested in a 10m<sup>2</sup> subplot in each of two corners. For comparison of data across space and through time, it is important that the dimensions of these plots and subplots be consistent across plots and sites. This protocol assumes that plots will be marked by a center point and four corners. The permanent markers define the corners of the plot and should maintain comparability through time. If this is not the case, plots must be established during each sampling bout according to the Plot Establishment Protocol (RD[09]). While delineating subplots, please take care to avoid trampling the plot – particularly the 1m<sup>2</sup> subplots.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

1. Delineate the sides of the 100m<sup>2</sup> subplot, the 10m<sup>2</sup> nested subplot (3.16 m from the nearest permanent marker at the plot corners or center), and the 1m<sup>2</sup> nested subplot with flags or appropriate markers.
  - a. Instructions in Appendix 0 assume the plot was established with precise square and exact 20m plot sides and that the tape can be stretched between corners with no obstacles.
  - b. Instructions in Appendix B.2 recognize an inevitable lack of absolute precision of the established markers and obstacles that are likely to obstruct the tape when stretched between markers.
2. The 1m<sup>2</sup> nested subplot is delineated with a rigid frame anchored at the corner by a permanent plot marker, a secondary marker at most sites, or marked during setup (in the case of 40.1.1 and 32.4.1).



**Figure 1.** The square, multi-scale plot is used to record plant species composition and cover. The plot includes nested subplots at specific locations within the plot.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

## B.2 General Collection

The plot-based collection requires photography, observation of primarily abiotic elements – termed ‘variables’ – in 1m<sup>2</sup> nested subplots, and observations of vascular plant species at multiple spatial scales.

1. Metadata. Record plotID, boutNumber, the primary botanist (measuredBy), additional staff (recordedBy), and date which should reflect the day the sampling was completed (if working with paper datasheets; this information is captured automatically by the handheld).
2. 1m<sup>2</sup> Nested Subplot. Photograph nested subplots (**photographs suspended until further notice**), record variable cover estimates, and identify and record cover and presence of individuals greater than 3m in height of vascular plant species in the subplot.
  - 1) **Photograph 1m<sup>2</sup> nested subplot**. Take a plane-view picture of the nested 1m<sup>2</sup> nested subplot such that the subplot frame fills the photograph while standing at the middle of the south edge outside the 1m<sup>2</sup> nested subplot. The amount of the subplot frame and vegetation to be captured in the image will depend on the height and density of vegetation; vegetation should not be manipulated to make subplot frame visible. If it is not possible to stand at the south edge, move to the west, north, east edge in that order. Once uploaded and back in the lab, the photograph name should include: plot\_ID, Module, subplot number, and date in the following format: CPER\_001\_PlantDiversity\_31.1.1\_20130812.
  - 2) **Measure and record variables**. Estimate and record the combined cover of each variable of abiotic (non-living) elements, non-vascular plant species, and overstory cover of species in each 1m<sup>2</sup> nested subplot (Table 3). Cover of any one variable shall not exceed 100 percent, but the total cover of multiple variables may be (but very rarely will be) greater than 100 percent. Observations should reflect those variables that cover the surface of the subplot (e.g. the moss growing on a rock, but not that part of the rock under the moss, or the litter on top of the soil but not the soil under the litter).

**Table 3.** Variables to be observed in the 1m<sup>2</sup> nested subplot.

Variable name	Description
Soil	Particles < 5mm diameter.
Rock	Inorganic particles ≥ 5mm diameter.
Wood	Woody organic material ≥ 5mm including living roots and material severed from the original source of growth and on the ground, including dead trees (either self-supported by roots, severed from roots, or uprooted) that are leaning > 45 degrees from vertical. Include the basal area and any woody organic material < 300cm in the cover estimate.
Litter	Unrooted organic material lying on the ground such as grass, leaves, pine needles, and twigs < 5 mm diameter.
Standing Dead	Standing dead woody material that is not severed from the original source of

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

	growth and not leaning > 45 degrees (please note ‘wood’ in the otherRemarks field). Also include desiccated herbaceous organic material from the previous calendar year or that cannot be identified (please note ‘herbaceous’ in the otherRemarks field). Species that might have been included had the sampling bout been longer or earlier in the year should be included in the plot species list.
Water	Standing or flowing water.
Lichen	Cyanobacteria or green algae.
Moss	Bryophytes. Typically small (1 – 10cm but up to 50cm), mosses, liverworts, hornworts.
Other non-vascular	Algae, please include macrofungi and biological soil crusts here as well and note in comments.
Scat	Animal dung, make note of species it originated from if possible in comments.
Overstory	An estimate of the total vegetation greater than or equal to 3m above the 1m <sup>2</sup> nested subplot; should include species not rooted in the nested subplot.

### 3) Measure and record plant species data.

- 1) Record the presence of living vascular plant species with stems emerging from within the 1m<sup>2</sup> nested subplot by entering Natural Resource Conservation Service (NRCS) US Department of Agriculture (USDA) PLANTS database code in the taxonID field for each species. A list of plants and codes, the NEON master taxon list of plant species codes can be found on the NEON intranet. In the event that a code is not available in the field and a code different from the NEON master code/NRCS code is used on a datasheet, the full scientific name associated with that code must be provided with the datasheet, either via annotation (see RD[04]) or by attaching a key to each datasheet so the correct code can be entered into the web-based user interface. If entering data into a mobile data recorder, any part of the scientific name or NRCS code can be entered to initiate filtering to the species of interest. If no species are found in the nested 1m<sup>2</sup> subplot, either click “No plant cover present” on the electronic device or record the same in the taxonIDRemarks field of the first line of the datasheet. Species-specific comments should be entered in the taxonIDRemarks field.

If an exact species determination can’t be made:

- Describe uncertainty about the genus or species with the appropriate identification qualifier in the idQ field (see B.3, Morphologically Challenging Species for correct use of idQ codes).
- Group or “lump” taxonomically similar species by entering the accepted NRCS genus code followed by sp/spp (depending on number of species in each plot), or enter possible species in the taxonIDRemarks field (see B.3, Morphologically Challenging Species).

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

- If a species cannot be identified in the field and the individual does not have parts needed for identification, enter the lowest taxonomic rank that can be determined (e.g., family or genus) followed by the sp. suffix (see section B.4, Unknown Plant Species). If it is not possible to identify to genus or to family please enter the code '2Plant Unknown Plant' on the datasheet, webUI, and or handheld device (please see the FAQ for plant diversity data entry on the NEON intranet for more information).
- If a species cannot be identified in the field and the individual does have observable identifiable characteristics, enter the lowest rank taxon code (e.g. genus), a unique name to track the species in morphospeciesID, and any descriptive comments in the morphospeciesIDRemarks (see section B.4, Unknown Plant Species). If it is not possible to identify to genus or to family please enter the code '2Plant Unknown Plant' on the datasheet, webUI, and or handheld device (please see the FAQ for plant diversity data entry on the NEON intranet for more information).

If there is not an NRCS code available in the master species list for a species identified:

- Check the spelling of the species and check the USDA PLANTS online database
- See if there is an updated synonym in the USDA PLANTS database that is in the master list and enter that synonym
- Conduct other searches for other synonyms that might be in the USDA PLANTS database
- If none of these solutions yields an appropriate solution, enter OTHE and put the scientific name and appropriate NRCS code in the comments for that field (please see the FAQ for entering plant data on the NEON intranet for more specifics on the use of OTHE).

- 2) Estimate the combined cover of plant material < 300cm in height of all individuals by species in the nested 1m<sup>2</sup> subplot. Measure cover as the percentage of ground surface obscured by the vertical projection of all aboveground parts of each species (Figure 2).
- For all individuals or stems of each species < 300cm in height, include the combined cover of all living vegetation (woody, foliar, herbaceous) AND select 'N' for Plant Height Over 300cm
  - For individuals or stems of each species > 300cm in height, record the combined cover of all plant material (the basal diameter, branches, foliage) < 300cm in height AND select 'Y' for Plant Height >300cm
  - If there are individuals or stems of a single species both < 300cm and > 300cm in a single 1m<sup>2</sup> subplot, enter the combined cover of all vegetation < 300cm (as above), AND select 'Y' for Plant Height >300cm

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

- Only estimate cover of plants, or portions of plant, with stems that originate within or have some part of the stem inside the subplot frame. Epiphytes not actually rooted on the ground of the nested subplot, but that are rooted to trees in the space extending above the nested subplot should be included. Record cover of those individuals <300cm in height from the ground. For those individuals > 300cm, record the identity of the species and check the ‘Plant Height Over 300cm’. It is understood that the identity and precise cover may be difficult to ascertain, in which case it might be necessary to identify to a higher taxonomic level.
- Estimate cover to nearest 1%.
- Enter 0.5 for estimates of cover <1%.



Figure 2. Estimates of cover should include all vegetative material < 300cm in height. For herbaceous growth (A), and shrubs (B) < 300cm, record the total combined cover by species; for tall trees with no woody branches or foliar growth < 300cm (C) record basal area and a height of > 300cm should be noted for that species; for trees (D) and shrubs (E) > 300cm that also have vegetative growth < 300cm, record the cover of vegetative growth < 300cm and indicate the presence of individuals > 300cm in height for that species. There will be instances when herbaceous growth <300cm (A) and trees >300cm (C) of the same species are found in the same 1m<sup>2</sup> subplot, in these cases record the combined cover and indicate the presence of species > 300cm.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

- There will often be spatial overlap of plant species. Cover should be recorded as the total aerial coverage for each species; estimates should not exceed 100 percent for a single species, but combined species total subplot cover may be greater than 100%.

Cover estimates can be made more repeatable across observer, plots, and sites with calibration:


- Familiarize yourself with what particular cover estimates (e.g., 1%, 10%, 15%, etc.) look like and use them as reference sizes. For example, if you know that 1% cover is about the same size as your fist, use your fist as a reference.
  - Each 1m<sup>2</sup> nested subplot frame is calibrated in 10cm sections to make cover estimates easier (Figure 3).
  - Visually group species together into a percent cover.
  - Fine tune estimate by subtracting out any spaces or gaps.
  - Check that your combined abiotic and select biotic variables (minus overstory) and your cover estimates for each taxonID sum to a total of at least 100%.
- 3) 10m<sup>2</sup> Nested Subplot. Record the identity of all species with stems in each 10m<sup>2</sup> nested subplot as described for the 1m<sup>2</sup> nested subplot. It is not necessary to record all (but see below) species already documented in those 1m<sup>2</sup> nested subplots in each respective 10m<sup>2</sup> nested subplot. However, it is acceptable to list species that were observed in the smaller, 1m<sup>2</sup> nested subplot (it may be difficult to remember, especially given the repetitive nature of the plot, and it is preferable to record species if there is any question) as these records will be handled when processed with NEON cyberinfrastructure. If there are no additional vascular plant species beyond those which have already been recorded in the nested subplot (for example 31.1.10 has no additional species other than what was found in 31.1.1) please add one plant species found in the nested 1m<sup>2</sup> subplot. Only click “NO PLANT COVER PRESENT” on the handheld and the web user interface if there are no plant species found when searching the remainder of the 10m<sup>2</sup> subplot.

There is no specific time that should be spent looking for plant species during search efforts. The search is best thought of in terms of a species-accumulation curve. Initial searching is likely to result in more species. A general guideline: if new species are being found, keep searching. If after five to ten minutes of gently moving dominant species to look for small and locally rare individuals – even crawling if necessary – no new species are found, then spend another five minutes and move on.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

- 4) 100m<sup>2</sup> Subplot. Record the identity of all plant species with stems in each 100m<sup>2</sup> subplot as described for the 10m<sup>2</sup> nested subplot. It is not necessary to record all species already documented in nested subplots. However, it is acceptable to list species that were observed in the smaller, 1m<sup>2</sup> and 10m<sup>2</sup> nested subplots (it may be difficult to remember, especially given the repetitive nature of the plot, and it is preferable to record species if there is any question) as these records will be handled when processed with NEON cyberinfrastructure. If there are no additional vascular plant species beyond those which have already been recorded in the nested subplots (for example 31.1.100 has no additional species other than what was found in 31.1.10) please add one plant species found in any of the nested subplots. Only click “NO PLANT COVER PRESENT” on the handheld and the web user interface if there are no plant species found when searching the remainder of the 100m<sup>2</sup> subplot.

As with searching the 10m<sup>2</sup> nested subplot, there is no specific time that should be spent looking for plant species during search efforts. The search is best thought of in terms of a species-accumulation curve. Initial searching is likely to result in more species. A general guideline: if new species are being found, keep searching, covering the entire area in a systematic manner such as walking lines or a grid. If after ten minutes of gently moving dominant species to look for small and locally rare individuals – even crawling if necessary – while searching the entire subplot and no new species are found, then spend another ten to fifteen minutes and move on.



**Figure 3.** The 1 m<sup>2</sup> subplot is calibrated with black and white marks to make estimates of plant species cover more accurate and repeatable

**B.3 Morphologically Challenging Species**

Identification qualifiers. In some cases there may be uncertainty regarding the identity of an individual. The lowest taxonomic rank that can be determined should be entered, and the

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

appropriate identification qualifier code applied (Table 4). Codes should be applied to the finest level of taxonomy recorded. For example, CS (“Roughly equals but “not sure” about the species) should only be applied if both a genus and species are recorded, while CG (“Roughly equals but “not sure” about the genus) should only be entered if a species-level determination is not possible. These codes should also only be applied to the recorded observation if there is uncertainty regarding the determination. If a taxonomic definition is not possible at a particular resolution (e.g. NEVER enter cf. species code when *Carex* spp. – for example – is entered) or if a morphospeciesID is used for an unknown species (see below), identification qualifier codes should not be applied. Another example: If it is thought that an individual might belong to *Achnatherum* sp., enter ‘cf.genus’ in the identification qualifier field to indicate this uncertainty. However, if BROMU (*Bromus* sp.), is selected to indicate that the species is unknown (see section B.4 below), an identification qualifier of cf. species should NOT be entered.

- Lumping (also see ‘Recording unknown species’ below). Some species will be difficult to differentiate. Because comparability of cover estimates must be consistent through time, it is necessary to identify these groups of species (e.g. genus) and consistently lump them through time and across field sampling efforts. It is difficult to know what this lumping might look like prior to the first field sampling year. A list of lumped species – specific to each site - should be developed over the first year of sampling and, if possible, based on conversations with botanists who work at the site or in the region. These lumped species should be identified to genus, and the proper NRCS code followed by the spp. suffix should be recorded.
- Cryptic species (slash species). Cryptic species issues arise when two species that are morphologically indistinguishable in the field co-occur (or might co-occur) at a site. NEON intends to add these species pairs to the master taxon lists to account for this. If a cryptic species pair is not currently available in the master list, the proposed species pair must be entered in the crypticSpeciesGroups spreadsheet on the NEON intranet. In the case that it is and will remain difficult to differentiate between two species of a single genus, enter the NRCS genus code followed by the sp. suffix (e.g., *Triticum* sp.) in the taxonID field, and enter the code for the two species in the taxonIDRemarks field (e.g. TRSA5/TRAE2) until the proposed “slash” pair is incorporated into the master list and is available on the webUI and/or field handheld device.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

**Table 4.** Identification qualifier codes (idQ) to designate unknown species or those species with uncertain identification in the field or after identification in the lab.

idqCode	identificationQualifier	Description
CS	cf. species	Roughly equals but "not sure" about the species
AS	aff. species	"Similar to, but is not" the species
CG	cf. genus	Roughly equals but "not sure" about the genus
AG	aff. genus	"Similar to, but is not" the genus
CB	cf. subspecies	Roughly equals but "not sure" about the subspecies
AB	aff. subspecies	"Similar to, but is not" the subspecies
CF	cf. family	Roughly equals but "not sure" about the family
AF	aff. family	"Similar to, but is not" the family
CV	cf. variety	Roughly equals but "not sure" about the variety
AV	aff. Variety	"Similar to, but is not" the variety

#### B.4 Unknown Plant Species

If a species determination cannot be made in the field, the presence of unknown species should be recorded or an individual should be collected for identification in the lab or with the assistance of expert botanists.

- Recording species that can't be identified to species. Species that can't be identified and do not possess sufficient parts to allow identification in the lab or with external help (remember NEON has a process to solicit assistance with plant identification from expert botanists) will likely be encountered. The NEON master taxon lists include codes for instances when identification below a given taxonomic rank (e.g., family, genus) cannot be made. These are indicated by a 'sp.' or 'spp.' in the scientific name. **Use the 'sp.' designation when only one unknown species is present (e.g., single individual is found, or sufficient reproductive parts are present to assume all individuals are the same species). Spp. is used when the group of individuals in question might belong to more than one species (e.g., many individuals, morphologically distinct features not discernable).** When one of these taxa is selected, an identification qualifier is not needed, unless the lowest taxonomic rank indicated (e.g., family, genus) is uncertain.
  - If there is likely only one species (can be multiple individuals) within any particular plot/nested subplot, record the lowest taxon rank with the sp. suffix (e.g. *Triticum* sp.) even if multiple unknown species or a different unknown species of the same family/genus are found in a different plot/nested subplot.
  - If there are multiple species within any particular plot/nested subplot, record the lowest taxon rank with the spp. suffix (e.g. *Triticum* spp.).
  - If neither the genus nor the family can be determined, enter '2Plant Unknown Plant' in the taxonID field (datasheet or electronic device).
  - For example, if you select BROMU (*Bromus* sp.), an identification qualifier of cf. species is unnecessary, as the 'sp,' indicates that the species is unknown. If, however, you think

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

that the individual might belong to *Achnatherum* sp., you would enter cf. genus into the identification qualifier field to indicate this uncertainty.

- Collecting and recording unknown species (morphospecies). Tracking unknown species that can later be identified is expected during the course of this work. If domain staff or an external facility are likely able to subsequently identify the tracked individual or ‘morphospecies’, the full scientific name associated with that morphospecies must subsequently be provided with the datasheet, either via annotation (seed RD[04]) or by attaching a key to each datasheet. If domain staff is not able to identify a given morphospecies prior to data entry (in the case of the webUI), the morphospecies ID must be recorded in the appropriate spreadsheet on the NEON intranet. When generating morphospecies, known taxonomic information should be recorded, a morphospeciesID should be created to track the species, and a specimen should be collected or photographed.
  - Taxonomy and morphospeciesID
    - a. Record the lowest taxon rank (family or genus) in the taxonID field.
    - b. Enter a descriptive name that will be memorable should the morphospecies be found in other plots in the morphospeciesID field.
    - c. Enter a description of the individual that might be useful when keying the plant in the lab (e.g. pubescent ligules, acidic moist habitat) in the morphospeciesIDRemarks field.
  - Collecting a specimen
    - a. Given NEON’s long-term monitoring efforts, unknown species should be collected from outside the 40 m x 40 m plot. Finding the same unknown species can sometimes take considerable time.
    - b. Collect representative parts of the entire individual, including the roots, flowers (if possible), and vegetative growth of grasses and forbs. A piece of a branch is usually sufficient for trees and shrubs. If a flower cannot be found, technicians can keep an eye open for an individual in flower for the rest of the sampling effort, but are not expected to return to a particular plot for the exclusive purpose of finding the individual in flower at a later date.
    - c. Place unknown specimens in sealable plastic bags (see **Error! Reference source not found.** for optional system for organization and transport of specimens). A cooler with an ice pack may also be used (optional) to prevent wilting of specimens, and may be particularly useful on hot days and/or when there is little shade available. Label plant with the unique (to the technician) morphospeciesID, measuredBy (botanist), date, GPS coordinates, elevation, and plot number (where species was initially found, if appropriate and if possible).
    - d. If collection of an individual is not possible, take photograph(s) of the individual (including flowers and other parts crucial to identification) and record photographic information in the morphospeciesIDRemarks field. Once

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

uploaded, the photograph should be labeled with morphospeciesID, plotID, and date as follows: alternatePappusHerb\_CPER\_001\_20130812.

- e. At the end of the field day, place plastic bags in a refrigerator until they are identified and/or placed in a plant press and dried for identification at a later date. It is imperative that the label information remain associated with the specimen. Specimens should not be left in the refrigerator for more than two days. Identification often requires a variety of dichotomous keys, a dissecting microscope, a dissecting kit, and a herbarium with voucher specimens for verification.

## B.5 Collections and Voucher Specimens

Plant species will be collected at NEON sites to: 1) facilitate the identification of species not identified in the field (see section B.4 above), 2) to create an herbarium in the domain support facility for training and quality assurance purposes (RD[12]), 3) contribute to a long-term record of plant diversity observations as part of the NEON archive program (the exact number to be collected and archived is pending), and 4) to create an archival-quality voucher for plant genetic collections (RD[06]).

The following guidelines should be considered when collecting specimens:

- 1) Select specimens in good condition, free of damage from insects and/or disease.
- 2) If possible, all parts of a plant should be collected, the roots, stems, flowers, fruits, and seeds. Collect at least stems, leaves, and flowers or fruit of herbaceous plants, and twigs, leaves, and flowers or catkins of trees and shrubs.
- 3) Place all specimens of a single species from one locality into one collection bag.
- 4) Add the plant tissue sampleID to the label if the specimen is a voucher for the plant tissue collection (RD[06]).

## B.6 Refreshing Sampling Kit

Be sure to have sufficient plastic bags, adhesive labels, and permanent markers for the next field sampling effort.

## SOP C Laboratory Processing and Analyses

The lab component of the plant biodiversity sampling includes three parts: plant pressing and drying (RD[12]), identification, and processing of a subset of collected species for voucher specimens. The exact order and need for each step with each specimen will depend on scheduling and time, and the objectives for the particular specimen.

Some species will come from the field and be identified fresh (without pressing), within two days of collection. If the specimen was destroyed during identification or was not intended for vouchering, there

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

is no need to save and press the specimen. Not every unknown plant species must be vouchered and submitted to the archive.

Some species will be collected and the botanist will not have time to identify them within two days, or will not be able to identify the specimen. These specimens should be pressed for identification at a later time (RD[12]), either by the botanist (with the help of an herbarium and/or books) or by sending them to an external expert.

Other specimens will be collected specifically for archive at the Domain Support Facility or an external archive facility. Specimens collected for external facilities should be treated with extra care to preserve diagnostic parts and pressed until dry but not mounted or labeled pending further directions from the archive. Those specimens destined for storage at the NEON Domain Support Facility should be mounted and labels should be prepared.

**C.1 Sample Processing Timing**

Specimens should not be left in the refrigerator for more than two days. They can be placed in the press, stored in a well-ventilated location, and identified at a later date. Specimens may remain in this state for months. Special care should be taken to ensure that specimens are not vulnerable to damage from insects.

**C.2 Identification of Unknown Species**

Ideally, a fresh (not pressed and dried) duplicate specimen – if the species is to be included in one of the archives described above - that is not to be included in the herbarium should be used for identification. Identification requires basic knowledge of morphological characteristics of different plant families, plant keys, access to a herbarium (University of Wyoming, University of Florida, etc.), a clean bench space in the lab, and a dissecting microscope, and dissecting kit. If there is any doubt, a duplicate specimen should be submitted to a taxonomic expert for identification, or specimen can be compared to online or local herbarium collections. Attempts should first be made to identify unknowns at the Domain Support Facility and possibly with the assistance of a herbarium. If identification is not possible, specimens can be sent to experts in regional flora.

After unknown specimens are identified, update the information in the data sheets, files or database. If the species is identified prior to submission of the data, it is not necessary to retain the original morphospecies name in the datasheets or entered data. Please visit the NEON intranet in the morphospeciesTracking folder to enter the final keyed identity determination of morphospecies.

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

## SOP D Data Entry and Verification

**The importance of thorough, accurate data transcription cannot be overstated; the value of the efforts in the field is only manifested once the data are properly entered for delivery to NEON’s end users.**

As a best practice, field data collected on paper datasheets should be digitally transcribed within 7 days of collection or the end of a sampling bout (where applicable). However, given logistical constraints, the maximum timeline for entering data is within 14 days of collection or the end of a sampling bout (where applicable).

Before entering data, all personnel must read RD[04] for complete instructions regarding manual data transcription. Prior to entering data via a web user interface (webUI), each technician shall enter a plot (or subplot) of data from one bout into the protocol-specific webUI housed on the Training portal, as described in RD[04].

Protocol-specific instructions and the associated data ingest workbook for entering data can be found on the NEON intranet in the FSU-FOPs folder. Prior to entering data please be sure to check the codes of each species to be sure that that appropriate NRCS code is attributed to the species detected. Due to the volume of plant species in US reflected in the USDA PLANTS database, the codes are often a bit more cryptic than four letters corresponding to genus and species. If the data is collected on a paper datasheet, it is preferable that the person who collected the data also enters the data or is at least familiar with flora at the site to reduce the possibility of errors in the data entry process. If the wrong code was used on the paper datasheet, the correct NRCS code must be annotated on the sheet.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

**SOP E Sample Shipment**

Information included in this SOP conveys science-based packaging, shipping, and handling requirements, not lab-specific or logistical demands. For that information, reference the [Collection shipping document](#) on [Collection's NEON intranet site](#). More information regarding the shipment of species to be identified by external experts, voucher specimens, and the desiccated plant tissue (RD[06]) will become available after institutions and facilities are identified.

Shipping details are TBD and will be included in a future revision of this protocol (as of rev F).

**E.1 Handling Hazardous Material**

N/A

**E.2 Supplies/Containers**

TBD

**E.3 Timelines**

TBD

**E.4 Conditions**

TBD

**E.5 Grouping/Splitting Samples**

TBD

**E.6 Return of Materials or Containers**

TBD

**E.7 Shipping Inventory**

TBD

**E.8 Laboratory Contact Information and Shipping/Receipt Days**

See the [CLA shipping document](#) on [CLA's NEON intranet site](#).

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

## 8 REFERENCES

### APPENDIX A DATASHEETS

The following datasheets are associated with this protocol:

**Table 5.** Datasheets associated with this protocol

NEON Doc. #	Title
NEON.DOC.001579	Datasheets for TOS Protocol and Procedure: Plant Diversity Sampling

These datasheets can be found in Agile or the NEON Document Warehouse.

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

## APPENDIX B QUICK REFERENCES

Tape measures and flags shall delineate and anchor the subplots. In relatively flat terrain with few obstacles such as trees, logs, and rocks, stretching tape around multiple sides of the plot results in precise placement of subplots (see B.1). In most cases obstacles will often result in a tape not reading exactly 40 m after stretching from point 31 past point 33 (south side of plot) to point 51 (east side of plot). In these cases (see B.2), the 1 m<sup>2</sup> and 10 m<sup>2</sup> subplots should be 1 m and 3.16 m in linear distance from the closest corner (typically a primary or secondary marker) – subplots should be anchored at the proximal plot corners and plot center (points 31, 33, 41, 49, 51). For example, 1 m and 3.16 m should be subtracted from whatever reading is displayed on the tape at point 33 to define the south side of subplots 32.2.1 and 32.2.10. Because there is not typically a marker at point 40 or point 42, these markers should be placed as close to 10 m north of points 31 and 33.

If permitted by the site host, and time and material allow, there is no reason from a NEON Science perspective not to increase the number of markers left at a plot. Placing more secondary markers could reduce plot establishment required for plant diversity sampling and increase repeatability of data.

### B.1 Delineating a precise plot with little to obstruct the tape on the perimeter.

The perimeter of the plot and subplots shall be delineated by tape measures and subplot frames as follows (Figure 4):

- 1) Begin in the south-west corner of the plot (point 31), at most sites this permanent marker will be labeled with information about the plot.
- 2) Anchor a 50 m tape and extend it towards the south-east corner (point 33).
  - a. Walk on the south side of the tape to avoid trampling plants inside the 20 x 20 m plot.
  - b. While pulling the tape, insert pin flags into the ground touching the outside edge of the tape at 1 m, 3.16 m, 10 m, 16.84 m, and 19 m.
- 3) Anchor the tape at the 20 m at the south-east corner of the plot (point 33) and pull it towards the marker at the north-east corner (point 51) of the plot.
  - a. Walk on the east side of the tape to avoid trampling plants inside the 20 x 20 m plot.
  - b. While pulling the tape, insert pin flags into the ground touching the outside edge of the tape at 21 m, 23.16 m, 30 m, 36.84 m, and 39 m.
- 4) Return to the south-west corner (point 31) of the plot.
- 5) Anchor the second 50 m tape and extend it towards the north-west corner (point 49).

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

- a. Walk on the west side of the tape to avoid trampling plants inside the 20 x 20 m plot.
  - b. While pulling the tape, insert pin flags into the ground touching the outside edge of the tape at 1 m, 3.16 m, 10 m, 16.84 m, and 19 m.
- 6) Anchor the tape at the 20 m at the north-west corner (point 49) of the plot and pull it towards the marker at the north-east corner (point 51) of the plot.
- a. Walk on the north side of the tape to avoid trampling plants inside the 20 x 20 m plot.
  - b. While pulling the tape, insert pin flags into the ground touching the outside edge of the tape at, 21 m, 23.16 m, 30 m, 36.84 m, and 39 m.
- 7) Anchor a third tape at the center of the plot (point 41) and extend it south toward the flag that at 10m.
- a. Insert pin flags into the ground at 1 m and 3.16 m.
- 8) Return to the center and extend the tape east toward the flag that at 30 m.
- a. Insert pin flags into the ground at 1 m, 3.16 m, 6.84 m, and 9 m.
- 9) Return to the center and extend the tape north toward the flag at 30 m.
- a. Insert pin flags into the ground at 1 m and 3.16 m.
- 10) Return to the center and extend the tape west toward the flag at 10 m. Insert pin flags into the ground at 1 m, 3.16 m, 6.84 m, and 9 m.




Figure 4. The plot will have some permanent markers and will also require temporary flags that are placed each time the plot is measured.

**B.2 Plot delineation with some lack of precision in plot and some obstacles (this will be most cases).**

This method is very similar to the previously described, but it recognizes that deviations in the distance between markers and obstacles in the tape may prevent the measures from working as described in Appendix 0 (e.g. if the tape must go around a tree between the southwest corner and the south-east corner the tape may not intersect the permanent marker at 20 m). The important difference is that subplots will be established from the nearest permanent marker. The idea is to delineate the plot boundary by connecting the permanent markers with the tape measure. The tape should be kept as close as possible to the ground, be forced through shrubs, and around trees to maintain the straightest line possible between markers. With two people, one person can anchor the tape at the south-west corner and pull the tape towards a person standing at the destination marker, or one person can hold the tape at the south-west corner and a second person can pull the tape towards the target marker. A compass might be helpful for establishing the direction the tape should be pulled.

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

After the tape is extended the subplot and 10 m markers can be established by pulling the tape tight from the nearest permanent marker and accounting for trees and other obstacles as needed. A string or equivalent material that measures 3.16 m is likely easier to use for establishing sides of the 10m<sup>2</sup> subplot. The perimeter of the plot and subplots can be delineated by tape measures and subplot frames as follows (Figure 4):

- 1) Record date and plot number.
- 2) Begin in the south-west corner of the plot (point 31), at most sites this permanent marker will be labeled with information about the plot.
- 3) Anchor a 50 m tape and extend it towards the south-east corner (point 33), walking on the south side of the tape and following a path that creates the straightest possible line towards the marker in the south-east corner.
- 4) Wrap the tape at the south-east corner/permanent marker (point 33) and extend it to the north-east corner (point 51) at approximately 40 m on the tape.
- 5) Return to the south-west corner (point 31) and while pulling the tape tight towards the south-east corner (point 33), insert pin flags into the ground touching the outside edge of the tape at 1 m, 3.16 m, 10 m.
- 6) Proceed to the south-east corner (point 33) and pull the tape tight (either wrapped around the marker or/or with a second person holding) from the south-east corner back towards the south-west corner (point 31) and insert flags at a distance of 1 m and 3.16 m from the south-east corner on the south edge of the plot.
- 7) With the tape anchored at the south-east corner (point 33), pull it tight towards the north-east corner (point 51) of the plot and insert pin flags at 1 m, 3.16 m, and 10 m from the south-east corner along the east side of the plot.
- 8) From this 10 m mark on the east edge of the plot, pull the tape tight back towards the south-east corner (point 33) and insert flags at a distance of 1 m and 3.16 m from the 10 m mark towards the south-east corner.
- 9) Proceed to the north-east corner (point 51) of the plot and pull the tape tight from the north-east corner back towards the south-east corner (point 33) and insert flags at a distance of 1 m and 3.16 m from the north-east corner on the east edge of the plot.

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

- 10) Return to the south-west corner (point 31) of the plot. Anchor the second 50 m tape and extend it towards the north-west corner (point 49), walking on the west side of the tape and following a path that creates the straightest possible line towards the marker at the north-west corner (point 49).
- 11) Wrap the tape at the north-west corner (point 49)/permanent marker and extend it to the north-east corner (point 51) at approximately 40 m on the tape.
- 12) Return to the south-west corner (point 31) and while pulling the tape tight towards the north-west corner (point 49), insert pin flags into the ground touching the outside edge of the tape at 1 m, 3.16 m, 10 m on the west side of the plot.
- 13) From this 10 m mark on the west edge of the plot, pull the tape tight towards the north-west corner (point 49) and place flags towards the north-west corner (point 49) at a distance of 1 m and 3.16 m from the 10 m mark on the west edge of the plot.
- 14) Proceed to the north-west corner (point 49) and pull the tape tight (either wrapped around the marker or/or with a second person holding) from the north-west corner (point 49) back towards the south-west corner (point 31) and insert flags at a distance of 1 m and 3.16 m from the north-west corner (point 49) on the west edge of the plot.
- 15) With the tape anchored at the north-west corner (point 49), pull it tight towards the north-east corner (point 51) of the plot and insert pin flags at 1 m, 3.16 m, and 10 m along the north side of the plot.
- 16) Proceed to the north-east corner (point 51) of the plot and pull the tape tight from the north-east corner (point 51) back towards the north-west corner (point 49) and insert flags at a distance of 1 m and 3.16 m from the north-east corner (point 51) on the north edge of the plot.
- 17) Proceed to the center of the plot (point 41).
- 18) Extend the third tape from the middle of the plot towards the 10 m mark on the north edge of the plot and while pulling the tape tight from the center, insert flags at a distance of 1 m and 3.16 m from the center.
- 19) Repeat the previous step in each direction from the plot center.

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

20) The boundary of the 10m<sup>2</sup> nested subplots can be defined by tape measures and pin flags. For 10m<sup>2</sup> nested subplots on the perimeter, a tape can be extended from a previously inserted survey or pin flag that is 3.16m from the corner where subplots are nested. To maintain a square nested subplot, this tape can target a pin flag that is 3.16m from a corner or center on the perimeter of an opposite side of the 100m<sup>2</sup> subplot (10 m away). Locating and aiming this targeted flag may require the help of a second person in dense vegetation. For example, the edge of the 10m<sup>2</sup> nested subplot in corner 1 of subplot 31 can be defined by stretching a tape from the flag at 3.16m on the south edge of the subplot toward the flag 3.16 m towards the center of the plot from the west edge. Delineating the boundary of the 10m<sup>2</sup> nested subplots anchored at the center of the plot requires that the target flag be added 3.16m from the flag at the middle of the a 20m edge of the plot. For example to defining the edge of the 10m<sup>2</sup> nested subplot in corner 1 of subplot 41 would require a flag 3.16m from the flag that is 10m between point 49 and 51 or the between point 33 and 51.

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

**APPENDIX C CHECKLISTS**

N/A

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

**APPENDIX D REMINDERS**

N/A

**APPENDIX E ESTIMATED DATES FOR ONSET AND CESSATION OF SAMPLING**

The dates in the table below are based on historic records and are estimates for the start and stop dates of sampling. It is essential that domain staff monitor real-time conditions to determine when to start (and stop) sampling, as described in Section 4 of this protocol. Dates for earliest onset of greenness and onset of minimum greenness that define the growing season can be found in the Plant Phenology protocol (RD[07]). The floristic peak – that time when the greatest number of species can possess parts required for identification – will occur inside that time period. The dates listed below reflect approximate start and end dates based on these greenness data. Please adjust this timing based on experience, expert opinion, and data from the Phenology sampling. If a site might require a second bout, please contact NEON TOS Science.

**Table 6.** Estimated dates of historical temperature thresholds

Domain	Site	# of Bouts	Approx. Start Date 1	Approx. End Date 1	Approx. Start Date 2	Approx. End Date 2
01	all	1	June	August		
02	all	1	April/May	August		
03	all	1	April	August		
04	all	1	TBD			
05	all	1	May	August		
06	all	1	April/May	August		
07	all	1	May	September		
08	all	1	April	September		
09	all	1	May/June	August		
10	CPER	2	May	June	August	September
	RMNP	1	June	August		
	STER	1	May	August		
11	CLBJ	TBD	April	June	August	September
	OAES	2	April	June		

Title: TOS Protocol and Procedure: Plant Diversity Sampling		Date: 1/27/2016
NEON Doc. #: NEON.DOC.014042	Author: D. Barnett	Revision: F

12	YELL	1	June	August		
13	NIWO	1	June	August		
	MOAB	1	March	July		
14	all	2	Feb	April	July	October
15	ONAQ	1	March	July		
16	all	1	May	August		
17	SJER	1	January	March		
	SOAP	1	May	August		
	TEAK	1	June	August		
18	all	1	June	August		
19	all	1	June	August		
20	OLAA	TBD	TBD			



<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

<i>Title:</i> TOS Protocol and Procedure: Plant Diversity Sampling		<i>Date:</i> 1/27/2016
<i>NEON Doc. #:</i> NEON.DOC.014042	<i>Author:</i> D. Barnett	<i>Revision:</i> F

**APPENDIX F    SITE-SPECIFIC INFORMATION**

N/A