



<i>Title:</i> Aquatic Site Sampling Design - NEON Domain 05		<i>Date:</i> 05/23/2019
<i>NEON Doc. #:</i> NEON.DOC.003604	<i>Author:</i> S. Parker	<i>Revision:</i> A

Aquatic Site Sampling Design – NEON Domain 05

PREPARED BY	ORGANIZATION	DATE
Stephanie Parker	AQU	3/26/2019
Nora Catolico	AQU	1/24/2018
Brandon Jensen	AQU	11/3/2016
Keli Goodman	AQU	5/04/2016
Jenna Stewart	AQU	4/20/2015
Charlotte Roehm	AQU	

APPROVALS	ORGANIZATION	APPROVAL DATE
Kate Thibault	SCI	04/04/2019
Mike Stewart	PSE	04/06/2019
David Barlow	ENG	05/22/2019

RELEASED BY	ORGANIZATION	RELEASE DATE
Anne Balsley	CM	05/23/2019

See configuration management system for approval history.

The National Ecological Observatory Network is a project solely funded by the National Science Foundation and managed under cooperative agreement by Battelle. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.



<i>Title:</i> Aquatic Site Sampling Design - NEON Domain 05		<i>Date:</i> 05/23/2019
<i>NEON Doc. #:</i> NEON.DOC.003604	<i>Author:</i> S. Parker	<i>Revision:</i> A

Change Record

REVISION	DATE	ECO #	DESCRIPTION OF CHANGE
Historic changes			Removed Round Lake; CM updated with new template and changes based on riparian habitat assessment timing; Updated bathymetry timing; Updated Crampton sampling wells and map; Updated LIRO map; Updated bio contingencies and water and sediment chem sampling locations in seepage lakes
A	05/23/2019	ECO-06095	Initial release



TABLE OF CONTENTS

1 DESCRIPTION.....3

1.1 Purpose 3

1.2 Scope..... 3

2 RELATED DOCUMENTS AND ACRONYMS3

2.1 Applicable Documents 3

2.2 Reference Documents..... 3

2.3 Acronyms 4

3 TEMPORAL SAMPLING STRATEGY4

3.1 Rationale 4

3.2 Approach..... 5

4 SAMPLING DATES7

4.1 Sensor Maintenance 8

4.2 Water Chemistry Sampling Dates 8

4.3 Groundwater Chemistry Sampling Dates..... 9

4.4 Biology Bout, Sediment Chemistry Sampling, and Riparian Assessment Dates 10

4.4.1 Suggested Biology and Sediment Chemistry Bout..... 12

4.4.2 Other Biology Sampling..... 12

5 PROTOCOL DISTURBANCE AND PRIORITIZATION14

6 SPATIAL SAMPLING STRATEGY15

6.1 General Site Sampling Locations..... 15

6.2 Site-Access and Instrument Locations 18

6.3 Riparian Sampling Locations 20

7 REFERENCES22

APPENDIX A LAKES.....23

APPENDIX B OBSOLETE LOCATIONS, CRAM24



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

LIST OF TABLES AND FIGURES

Table 1. Duration, frequency and prioritization of field activities and long term monitoring for NEON lake sites as a function of targeted constraints and driving variables. For associated lab hours, see Appendix A. (*May be scheduled more frequently if a stochastic event significantly alters the lake basin.)..... 6

Table 2. Rule sets for sampling modules in lakes. Deviations may be allowed with science approval..... 7

Table 3. Proposed water chemistry sampling dates for D05 Crampton and Little Rock Lakes. Dates are estimated based on available local data and may shift based on actual site conditions. Please note that dates are suggested, but should be adjusted as necessary, following the guidelines above. Although Tuesdays are the target, sampling may be shifted so that water chemistry sampling at LIRO occurs on Mondays while sampling at CRAM occurs on Tuesdays. 9

Table 4. Groundwater Observation Wells at D05 Crampton and Little Rock Lakes. **Wells for groundwater chemistry sampling are denoted in bold text.** 10

Table 5. Proposed groundwater chemistry sampling dates for D05 Crampton Lake..... 10

Table 6. Proposed Biological sampling windows for D05 Crampton Lake and Little Rock Lake. Fish sampling and Sediment Chemistry will take place during Bouts 1 and 3. The riparian habitat assessment peak greenness window may not coincide with the bout windows. 11

Table 7 Disturbance Criteria for lake sampling. Impact level: high (4), medium/high (3), medium/low (2), low (1), none (0). Bathymetry/morphology spans the entire permitted area. Sensors are located at the deepest point in the lake, and near the lake inlet and outlet. 14

Table 8. Module-specific sampling locations..... 15

Table 9. CRAM Sampling Locations. Proposed coordinates are determined prior to sampling at HQ. Field Science coordinates are groundtruthed in the field and reported to science. If available in the table, Field Science coordinates should be used for sampling..... 17

Table 10. LIRO Sampling Locations. Proposed coordinates are determined prior to sampling at HQ. Field Science coordinates are groundtruthed in the field and reported to science. If available in the table, Field Science coordinates should be used for sampling..... 17

Figure 1. Proposed bouts for biological sampling in D05 Crampton Lake. Sediment Chemistry and fish sampling occurs during Bouts 1 and 3. 13

Figure 2. Proposed bouts for biological sampling in D05 Little Rock Lake. Sediment Chemistry and fish sampling occurs during Bouts 1 and 3. 13

Figure 3. General diagram for an AQU site showing sampling locations in a seepage lake system. 16

Figure 4. Site access and instrument locations at D05 Crampton Lake..... 18

Figure 5. Site access and instrument locations at D05 Little Rock Lake. 19

Figure 6. CRAM ideal riparian sampling design 20

Figure 7. LIRO ideal riparian sampling design..... 21



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

1 DESCRIPTION

1.1 Purpose

The goal of the National Ecological Observatory Network (NEON) is to enable understanding and forecasting of the impacts of climate change, land use change, and invasive species on continental-scale ecology.

A disparity exists in the scale of organisms and their effects on the global environment (Hargrove & Pickering, 1992). While environmental impacts often occur at the largest scales, small scale biological and physical processes need to be understood in order to document responses of organisms, communities, populations and other small scale phenomena (Keller et al., 2008). Data will be gathered from the level of gene to ecosystem at a local to continental scale using standardized field procedures and sample processing. In order to address this disparity, NEON will approach the Grand Challenge questions through an analysis of processes, interactions and responses occurring across spatial and temporal scales.

The local data collected at NEON sites within the 20 Domains will be integrated with the targeted regional data from NEON airborne instrumentation. This will provide a direct linkage in spatial and temporal scaling from NEON’s distributed sensor network and in-situ field measurements, coupled with individual plant or canopy measurements to plot or stand level observations, and ultimately to the continental scale.

1.2 Scope

This document outlines the Domain 05 site-specific sampling strategy proposed for NEON Aquatic field sampling activities and other directly associated activities that will be used to address key data products related to the overarching Grand Challenge questions. It provides the sampling rationale for given parameters.

2 RELATED DOCUMENTS AND ACRONYMS

2.1 Applicable Documents

Applicable documents contain information that shall be applied in the current document. Examples are higher level requirements documents, standards, rules and regulations.

AD[01]	NEON.DOC.000001	NEON Observatory Design
AD[02]	NEON.DOC.002652	NEON Level 1, Level 2, Level 3 Data Products Catalog
AD[03]	NEON.DOC.005011	NEON Coordinate Systems Specification

2.2 Reference Documents

Reference documents contain information complementing, explaining, detailing, or otherwise supporting the information included in the current document.



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

RD [01]	NEON.DOC.000008	NEON Acronym List
RD [02]	NEON.DOC.000243	NEON Glossary of Terms
RD [03]	NEON.DOC.001152	NEON Aquatic Sample Strategy Document
RD [04]	NEON.DOC.001085	AOS Protocol and Procedure: Stream Discharge
RD [05]	NEON.DOC.001197	AOS Protocol and Procedure: Bathymetry and Morphology of Lakes and Non-Wadeable Streams
RD [06]	NEON.DOC.002905	AOS Protocol and Procedure: Water Chemistry Sampling in Surface Waters and Groundwater
RD [07]	NEON.DOC.001886	AOS Protocol and Procedure: Stable Isotope Sampling in Surface and Ground Waters
RD [08]	NEON.DOC.001199	AOS Protocol and Procedure: Surface Water Dissolved Gas Sampling
RD [09]	NEON.DOC.001191	AOS Protocol and Procedure: Sediment Chemistry Sampling in Lakes and Non-Wadeable Streams
RD [10]	NEON.DOC.003044	AOS Protocol and Procedure: Aquatic Microbe Sampling
RD [11]	NEON.DOC.003045	AOS Protocol and Procedure: Periphyton, Seston, and Phytoplankton Sampling
RD [12]	NEON.DOC.003039	AOS Protocol and Procedure: Aquatic Plant, Bryophyte, Lichen, and Macroalgae Sampling
RD [13]	NEON.DOC.003046	AOS Protocol and Procedure: Aquatic Macroinvertebrate Sampling
RD [14]	NEON.DOC.001194	AOS Protocol and Procedure: Zooplankton Sampling in Lakes
RD [15]	NEON.DOC.003826	AOS Protocol and Procedure: Riparian Habitat Assessment
RD [16]	NEON.DOC.001296	AOS Protocol and Procedure: Fish Sampling in Lakes
RD [17]	NEON.DOC.004613	NEON Preventative Maintenance Procedure: AIS Buoy

2.3 Acronyms

C0-C3	Buoy sensor set
IN	Inlet sensor set
OT	Outlet sensor set
GDD	Growing degree days
MGC	Multivariate geographic clustering
MODIS	Moderate Resolution Imaging Spectroradiometer
NOAA	National Oceanic and Atmospheric Administration
NCDC (NCEI)	National Centers for Environmental Information

3 TEMPORAL SAMPLING STRATEGY

3.1 Rationale

NEON designed a set of domains based on a statistically rigorous analysis using national data sets for eco-climatic variables, based upon algorithms for multivariate geographic clustering (MGC) (Hargrove & Hoffman, 1999, 2004). The MGC approach identified nine primary climate state variables that could define the domains, allowing for regionalization of primary features within each domain. In order to replicate the strategy used for the large scale spatial design of NEON, Aquatics has adapted this approach and



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

modified the list of the nine variables by identifying variables that were equally pertinent to the large scale temporal design, and by adding critical variables that affect physical, biological and chemical parameters in aquatic environments.

Aquatic ecosystems exhibit physical, chemical and biological variability over a wide range of spatial and temporal scales (Steele, 1978). This has resulted in a movement towards research approaches that utilize concurrent field based, buoy, aircraft, and satellite sampling strategies in order to measure physical, chemical and biological distributions over large areas synoptically and over long time periods. The integration of such sampling strategies across scales is an integral part of NEON’s approach to the addressing the Grand Challenge questions (Keller et al., 2008).

NEON must be able to extrapolate relationships between drivers (climate change, land use change, and biological invasions) and ecological consequences to areas that are not sampled by NEON facilities but where partial, extensively sampled, or gridded information is available. In order to obtain this NEON’s temporal sampling strategy must be equally designed to detect and quantify trends over time, as well as characterizing the spatial pattern of those trends. The sampling approach at the field scale, hence, must address the temporal.

3.2 Approach

Sampling strategies must cover a range of temporal scales and must address issues of duration and frequency of sampling activities. The design of the temporal strategy for NEON Aquatics addresses both the duration and frequency of the field activities as well as the small scale but long-term continuous monitoring data collection. In addition, prioritization of the physical, biological and chemical parameters needs to be identified. The general layout of a NEON lake site are presented in Section 6.

NEON Aquatics has proposed the following approach in order to determine the sampling duration and frequency that will yield the best estimate of composition and/or concentration of the physical, biological and chemical parameters (Table 1)

Physical/Chemical: Air temperature has been identified as the main variable defining the timing and frequency of sampling for physical and chemical parameters. Air temperature controls the dynamics of ice-on and ice-off events as well as stratification and turnover events.

Biological: Degree days, water temperature, and riparian greenness are the primary variables identified for defining the timing and frequency of sampling of most biological parameters.

Sampling modules may also have specific rule sets that dictate the order and timing of collection, as well as time constraints on laboratory work to maintain viable samples. The rule sets below (Table 2) have been identified for specific sampling modules.



Table 1. Duration, frequency and prioritization of field activities and long term monitoring for NEON lake sites as a function of targeted constraints and driving variables. For associated lab hours, see Appendix A. (*May be scheduled more frequently if a stochastic event significantly alters the lake basin.)

Sampling Module	Sampling Duration (hrs)	Sampling Frequency (x per year)	Constraints on Sampling	Driving Metrics for Sampling	Priority
Sensor Maintenance					
Surface water	1-2	26	Water Temperature Discharge	None	High
Meteorological	1-2	26	Weather	None	High
Groundwater (light)	1-2	26	Weather	None	High
Groundwater (full)	2-4	4	Weather	None	High
Well redevelopment	4	1	Weather	None	High
Physical					
Bathymetry	8-40	1 per 5 yrs*	Wind Ice-off	Riparian greenness	Low to Medium
Biological					
Surface Microbes	2-4	6	Ice-off Wind	Precipitation Water Temperature	High
Aquatic plants and Macroalgae	3-8	3	Ice-off Wind	Precipitation Light (PAR)	High
Macroinvertebrates	3	3	Ice-off Wind	Precipitation Water Temperature	High
Zooplankton	3	3	Ice-off Wind	Precipitation Water Temperature	High
Periphyton and phytoplankton	3	3	Ice-off Wind	Precipitation Light (PAR)	High
Fish	8-40	2	Ice-off Wind	Precipitation Water Temperature	Medium
Riparian habitat assessment	2-4	1	Wind	Riparian greenness	Low
Chemical					
Surface water chemistry	1-3	12	Ice-off Wind	Precipitation Water Temperature	High
Dissolved gas	1	12	Ice-off Wind	Precipitation Water Temperature	Medium
Isotopes	2	12	Ice-off Precipitation	Precipitation Water Temperature	High
Sediment chemistry	4-8	2	Ice-off Wind	Flow Regime Water Temperature	Low to medium
Groundwater chemistry	8-20	2	Sufficient Water in Well	Groundwater Elevation, Seasonal (spring, fall)	Medium to High



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

Table 2. Rule sets for sampling modules in lakes. Deviations may be allowed with science approval.

Protocol	Rule set
Water chemistry, dissolved gas, and isotopes	Should be completed first to reduce the risk of contamination. However, if completing multiple protocols that could take more than a few hours, collect chemistry samples last to reduce the time between collection and processing/shipping.
	Collect recurrent samples on Tuesdays, when possible.
	Alkalinity/ANC lab processing must begin within 24 hours of collection, or the sample must be flagged.
Surface water microbes	Sample in conjunction with recurrent (usually Tuesday) water chemistry.
	Filters must be flash-frozen in the field, and kept frozen until storage in -80 °C freezer. If processing in the domain lab, freeze at -80 °C within 4 hours of collection.
	Cell counts must be preserved in the field. Maximum time to preservation if bad weather = 4 hours.
Aquatic plants	Lab processing must begin within 48 hours of collection, or the sample must be flagged. AFDM samples may be dried and placed in desiccators until enough room is available in the muffle furnace.
	Biomass collection (clip harvest) only occurs during Bout 2.
Macroinvertebrates	Must be preserved within 1 hour of collection.
	Preservative change must occur within 12-72 hours of collection.
Zooplankton	Must be preserved with 30 minutes of collection.
Periphyton/Phytoplankton	Lab processing must begin within 24 hours of collection. AFDM samples may be dried and placed in desiccators until enough room is available in the muffle furnace. Minimum lab processing time spans 2 days.
	Sample must be kept cool (~4 °C) and dark until processing at the domain lab.
	Chlorophyll filters must be shipped to the external facility within 7 days of collection.
Sediment chemistry	Start field collection after non-fish biological sampling to minimize disturbance.
Fish	Schedule within 2 weeks of macroinvertebrate collection (biology bouts 1 and 3). Contingency situations may cause this time to be greater than 2 weeks.
	If conditions do not allow for fish sampling during bout 1, then sample when safe conditions allow up to 2 weeks before the start of bout 2. If conditions do not allow for fish sampling to occur during bout 3, then sample when safe conditions allow up to 30 days beyond the end of bout 3.
Bathymetry	Bathymetry occurs every 5 years unless extreme events warrant more frequent surveys.
	Bathymetric mapping occurs at peak greenness, during Bio Bout 2 or within ± 2 weeks of aquatic plant sampling.
Riparian habitat assessment	Riparian habitat assessment must occur during peak greenness.
Groundwater Chemistry	Completed within ± 1 day of water chemistry (contingency situations may necessitate 2 days).
Well redevelopment	Must not occur in the 2 weeks prior to groundwater chemistry sampling.

4 SAMPLING DATES

The surface water sampling strategy for the D05 lake sites (Crampton Lake and Little Rock Lake) is based on annual air temperature data collected from NOAA National Climatic Data Center (NCDC) and from the near-real time NEON data collected at the meteorological stations. Because these sites are ice-covered throughout parts of the year, surface water samples can be taken on a semi-monthly basis combined with more intensive sampling around ice-on and ice-off, while organismal sampling is based on accumulation of growing degree days throughout the season.



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

The following Tables and Figures indicate proposed sampling dates for all sample protocols to be undertaken at Domain 05 over the course of a year.

4.1 Sensor Maintenance

Sensor preventative maintenance for in-lake sensors and the meteorological station is scheduled every other week. Groundwater well maintenance includes light sensor maintenance every other week (confirm that the cables have not slipped, check for ice accumulation on the solar panel, remotely monitor the data stream), full maintenance quarterly (visually inspect the sensor, check the desiccant, check water clarity with bailers, check for roots in wells known to have that issue), and well redevelopment once per year. Additional details may be found in the preventative maintenance documents for each sensor and the lake buoys (RD[17]).

4.2 Water Chemistry Sampling Dates

Water chemistry includes sampling for water chemistry, aquatic stable isotopes, and dissolved gas in surface waters. These protocols should be completed on the same day as each other at each site.

Alkalinity and ANC titrations: Following a minimum of a year of alkalinity and ANC titrations at lake inflow, center (buoy), and outflow, it was determined that no significant difference existed between the three lake locations. Thus, we will only complete alkalinity and ANC titrations from the buoy location.

Standard recurrent sampling should take place 12 times per year on every first Tuesday of the month starting on the first Tuesday of the year, in coordination with TIS chemistry sampling and other national programs to enable standardization. If you cannot sample all sites on the same day, prioritize the core site for Tuesday sampling and sample the other site the following day or, if necessary, the following Tuesday.

Because these sites experience sustained winter temperatures below 0 °C, sampling will be less frequent than once monthly during the winter months and more frequently than once a month around the shoulder periods when turnover occurs in the lake coinciding with ice-on and ice-off dates (Table 3). Ice-off can be evaluated remotely by monitoring the staff gauge camera feed at the domain support facility. Ice-off in lakes is defined by the first loss of ice from the center of the lake in the spring. Ice-on in lakes is defined by the first ice coverage of the central part of the lake in the fall. Stratification can be determined by remotely evaluating temperature chain data.

Ice-off sampling strategy: One sample bout should occur one month prior to the long-term average of ice off conditions. The following sampling should occur within 1 week (maximum 2) of ice-off conditions assuming safe conditions allow access to the water body.

Ice-on sampling strategy: One sample should occur 2 weeks prior to the long-term ice-on averages for the region. Safe conditions for access to the lake must be met. The following sampling bout should occur



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

2 months after long-term ice-on averages for the region. Safe conditions require a minimum of 6" of ice to be able to safely access the lake for sampling.

Table 3. Proposed water chemistry sampling dates for D05 Crampton and Little Rock Lakes. Dates are estimated based on available local data and may shift based on actual site conditions. Please note that dates are suggested, but should be adjusted as necessary, following the guidelines above. Although Tuesdays are the target, sampling may be shifted so that water chemistry sampling at LIRO occurs on Mondays while sampling at CRAM occurs on Tuesdays.

CRAM/LIRO Bout	Date
1	First Tuesday of February (assuming safe conditions)
2	First Tuesday of March (assuming safe conditions)
3	First day of ice off (approx. 3 rd week of April, assuming safe conditions)
4	First Tuesday of May (flexible)
5	First Tuesday of June
6	First Tuesday of July
7	First Tuesday of August
8	First Tuesday of September
9	First Tuesday of October
10	Third Tuesday of October
11	First Tuesday after destratification (approx. 1 st week of November)
12	Third Tuesday of November (assuming safe conditions)

4.3 Groundwater Chemistry Sampling Dates

Groundwater chemistry includes sampling for water chemistry and aquatic stable isotopes (²H and ¹⁸O-H₂O only). These protocols should be completed on the same day as each other at each site.

Groundwater samples will be collected twice per year from a subset of 4 wells per site (Table 4). The wells will be specified prior to sampling and will remain the same between bouts. Two wells are sampled on the inlet side, and two on the outlet side of the lake. This will allow for chemical comparisons at opposite ends of the regional flow paths in addition to lake surface water samples.

The range of groundwater sampling dates, shown in Table 5 has been selected to target one sampling event in the spring and one in the autumn conditions. Groundwater chemistry should be sampled on or near the target date if at all possible. Groundwater sampling should be timed to occur on the same day (preferred) or within 1-2 days (preferably 1 day) of the surface water collection. This constraint aims to clarify origin of chemical fluxes between upstream sources versus local groundwater sources to the surface water by tightly coupling in time the two sampling bouts. The date range is provided to allow flexibility for Field Ops in selecting a time within the sampling event window where both sampling bouts can be performed in a maximum of a three-day period. Dates will be refined after the first few years of site-specific water table data are available for analysis.



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

Table 4. Groundwater Observation Wells at D05 Crampton and Little Rock Lakes. **Wells for groundwater chemistry sampling are denoted in bold text.**

CRAM Well ID	Latitude	Longitude
D05-CRAM-OW-01	46.21281776	-89.47709279
D05-CRAM-OW-02	46.21289759	-89.47754043
D05-CRAM-OW-03	46.21115824	-89.47855
D05-CRAM-OW-04	46.20826637	-89.47925944
D05-CRAM-OW-05	46.20759081	-89.47878602
D05-CRAM-OW-06	46.20562645	-89.47707533
D05-CRAM-OW-07	46.20449567	-89.47090402
D05-CRAM-OW-08	46.21145807	-89.47832631
LIRO Well ID	Latitude	Longitude
D05-LIRO-OW-01	45.994754	-89.707967
D05-LIRO-OW-02	45.994405	-89.707149
D05-LIRO-OW-03	45.996602	-89.699274
D05-LIRO-OW-04	46.000149	-89.701039
D05-LIRO-OW-05	46.000535	-89.702349
D05-LIRO-OW-06	46.001848	-89.703917
D05-LIRO-OW-07	46.001007	-89.705512

Table 5. Proposed groundwater chemistry sampling dates for D05 Crampton Lake.

CRAM Well Bout	Start Date	End Date
1	April 10	May 10
2	October 10	November 10
LIRO Well Bout	Start Date	End Date
1	April 10	May 10
2	October 10	November 10

4.4 Biology Bout, Sediment Chemistry Sampling, and Riparian Assessment Dates

The biology bout windows for lakes are based on a combination of parameters at each site. Using mean daily air temperature (NOAA NCDC datasets) to calculate growing degree days (centering around 10%, 50%, and 90% gdd) and the MODIS dataset to estimate riparian greenness (green-up and brown-down; Figure 1, Figure 2), 1 month sampling windows were pre-determined for all sites. Sampling windows may be adjusted for ice off dates as all biological sampling (except for surface water microbes) is based on actual conditions at the site. The riparian assessment will be conducted during the site-specific peak greenness window defined by the MODIS dataset which may or may not coincide with the biological and sediment sampling bout dates (Table 6).

- Surface water microbes in lakes should be sampled in conjunction with the standard recurrent water chemistry sampling, every other month, 6 times per year. If water chemistry sample timing fluctuates due to ice on/ice off dates, sample on every other water chemistry sampling date.



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

- Surface water DNA microbe samples collected during July or August should be marked for metagenomics analysis.
- Sampling for all other biological modules (aquatic plants/macroalgae, macroinvertebrates, zooplankton, phytoplankton/periphyton) as well as sediment chemistry, follow pre-determined sampling windows presented in Table 6. Sediment chemistry and Fish are sampled twice per year during Bouts 1 and 3.
 - The biology/sediment chemistry bout windows may be adjusted to start 3 days earlier or and/or end 3 days later than the dates listed in Table 6 to allow for more flexibility in scheduling. Any sampling outside of the bout window plus the 3-day buffer will require an entry in NEON’s problem-tracking system.
 - Sampling for each module at a site must occur within one day, with the exception of bathymetry and fish which may take up to 5 days at a site.
 - Bout 3 fish sampling may occur up to 30 days past the end of the Bout 3 window, if conditions allow (i.e., flowing water is present and it is safe to sample). Fish sampling should be scheduled within the bout windows, however sampling may be pushed later using this contingency as weather dictates.
- The riparian habitat assessment will occur within the dates provided in Table 6.

At northern lakes where ice on/ice off dates are a consideration, consider the suggested sampling bouts a guideline. Sample within the windows provided below if possible. On years where the ice off date is later than the start of Bout 1, adjust the window to 1 month after the date of actual ice off for sampling.

Table 6. Proposed Biological sampling windows for D05 Crampton Lake and Little Rock Lake. Fish sampling and Sediment Chemistry will take place during Bouts 1 and 3. The riparian habitat assessment peak greenness window may not coincide with the bout windows.

CRAM Bio Bout	Start Date	End Date
1	April 20	May 18
2	July 5	August 2
3	September 13	October 11
Riparian	May 29	September 11
LIRO Bio Bout	Start Date	End Date
1	April 14	May 12
2	July 9	August 6
3	September 9	October 7
Riparian	May 29	September 11



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

4.4.1 Suggested Biology and Sediment Chemistry Bout

1. Aquatic plants, macroinvertebrates, periphyton/phytoplankton, zooplankton (in any order)
 - a. Secchi/Depth profile data collection must also occur on days when phytoplankton and zooplankton are sampled.
2. Sediment chemistry (Bouts 1 and 3 only)
3. Fish (Bouts 1 and 3 only)

4.4.2 Other Biology Sampling

- Surface water microbes – 1st water chemistry bout of every-other month (likely Tuesday)
- Bathymetry – schedule within ± 2 weeks of Bout 2 aquatic plant sampling
 - Occurs every 5 years unless morphology changes significantly due to an extreme event
- The riparian habitat assessment can be scheduled anytime within the peak greenness window




Figure 1. Proposed bouts for biological sampling in D05 Crampton Lake. Sediment chemistry and fish sampling occur during Bouts 1 and 3.




Figure 2. Proposed bouts for biological sampling in D05 Little Rock Lake. Sediment chemistry and fish sampling occur during Bouts 1 and 3.



5 PROTOCOL DISTURBANCE AND PRIORITIZATION

5.1 Disturbance Criteria

Each aquatic protocol has its own unique sensitivity to disturbance and perturbations (Table 7). These sensitivities should dictate the order in which protocols are completed.

Table 7 Disturbance Criteria for lake sampling. Impact level: high (4), medium/high (3), medium/low (2), low (1), none (0). Bathymetry/morphology spans the entire permitted area. Sensors are located at the deepest point in the lake, and near the lake inlet and outlet.

Sample	Requirements	Impact Level	Disturbance
Sensor maintenance	None	1	Boat activity near sensor locations
Bathymetry	None	1	Boat activity throughout lake
Aquatic plants	None	3	Benthic collection at randomized points throughout the lake
Invertebrates	None	3	Benthic collection near water chemistry sampling sites and wading and substrate disturbance near shore
Zooplankton	6 hours- no disturbance that causes turbid conditions	2	Boat activity near sensor locations
Periphyton and phytoplankton	6 hours- no disturbance that causes turbid conditions	2	Wading and substrate disturbance near shore, boat activity near sensor locations
Sediment chemistry	None	4	Boat activity and benthic disturbance near sensor locations
Fish	6 hours- no disturbance that causes turbid conditions	4	Boat activity and wading nearshore.
Surface water chemistry, dissolved gas, isotopes, and surface microbes	None	1	Boat activity near sensor locations
Groundwater chemistry	None	0	Groundwater Removal
Riparian habitat assessment	None	2	Boat activity and substrate disturbance nearshore



6 SPATIAL SAMPLING STRATEGY

6.1 General Site Sampling Locations


Lake sampling protocols reference several locations within the lake, including the buoy (or center) location, the inlet sensor, and the outlet sensor (i.e., water chemistry, isotopes, dissolved gas, surface water microbes, phytoplankton, zooplankton protocols). Lake sites are also divided into 10 sections divided by the riparian bank locations (Figure 3), which are used when sampling macroinvertebrates, periphyton, and the riparian habitat assessment. Locations provided in the “proposed” columns in Table 9 and Table 10, are estimates. Specific coordinates at each site will be used for the life of the site.

Table 8. Module-specific sampling locations.

Sampling module	Location
Surface water chemistry	Buoy and groundwater wells
Dissolved gas	Buoy
Isotopes	Buoy and groundwater wells
Surface water microbes	Buoy
Phytoplankton	Buoy, lake inlet sensor (adjusted in-lake location for bio sampling), lake outlet sensor (adjusted in-lake location for bio sampling)
Riparian habitat assessment	Sections determined by HQ and provided to domain staff
Periphyton	In riparian sections, exact location determined by field ecologists
Aquatic plants, bryophytes, and macroalgae	10 randomized points
Macroinvertebrates (ponar)	Buoy, lake inlet sensor (adjusted in-lake location for bio sampling), lake outlet sensor (adjusted in-lake location for bio sampling)
Macroinvertebrates (sweep)	In riparian sections, exact location determined by field ecologists
Zooplankton	Buoy, lake inlet sensor (adjusted in-lake location for bio sampling), lake outlet sensor (adjusted in-lake location for bio sampling)
Fish	In riparian sections, exact location determined by field ecologists
Sediment chemistry	Buoy, lake inlet sensor (adjusted in-lake location for bio sampling)
Groundwater wells	Locations determined by HQ
Bathymetry	Whole lake



LAKES



Observational Sampling

- Water Chemistry, Isotopes, Dissolved Gas, Surface Microbes
- Riparian Assessments
- Sediment Chemistry
- Biology and Fish

Automated Instrument Measurements

- Meteorological Station
- Groundwater Wells
- Buoy with sensors
- Pressure/temperature sensors

Figure 3. General diagram for an AQU site showing sampling locations in a seepage lake system.



Title: Aquatic Site Sampling Design - NEON Domain 05		Date: 05/23/2019
NEON Doc. #: NEON.DOC.003604	Author: S. Parker	Revision: A

Table 9. CRAM Sampling Locations. Proposed coordinates are determined prior to sampling at HQ. Coordinates are groundtruthed by Field Science in the field and reported to Science. If available in the table, Field Science coordinates should be used for sampling.

<i>Location ID</i>	<i>Description</i>	<i>Proposed Latitude</i>	<i>Proposed Longitude</i>	<i>Field Sci Latitude</i>	<i>Field Sci longitude</i>
01 - Riparian	Riparian coordinates*	46.211195	-89.47836	46.211210	-89.478273
02 - Riparian	Riparian coordinates*	46.211906	-89.474979	46.211931	-89.475000
03 - Riparian	Riparian coordinates*	46.211687	-89.470889	46.211728	-89.470910
04 - Riparian	Riparian coordinates*	46.209663	-89.468757	46.209673	-89.468749
05 - Riparian	Riparian coordinates*	46.208185	-89.47128	46.208227	-89.471270
06 - Riparian	Riparian coordinates*	46.208945	-89.472853	46.209004	-89.472849
07 - Riparian	Riparian coordinates*	46.207468	-89.475203	46.207430	-89.475203
08 - Riparian	Riparian coordinates*	46.205972	-89.472684	46.205971	-89.472701
09 - Riparian	Riparian coordinates*	46.206522	-89.47649	46.206543	-89.476401
10 - Riparian	Riparian coordinates*	46.209147	-89.477303	46.209170	-89.477358
Inlet	Inlet location from SCR	46.21067	-89.478566	46.208042	-89.476412
Outlet	Outlet location from SCR	46.211028	-89.469083	46.212120	-89.474871
C0 (buoy)	S1 sensor location from SCR	46.210592	-89.476862	46.210556	-89.476651

* Riparian coordinates should be approximately evenly spaced throughout the sampling area

Table 10. LIRO Sampling Locations. Proposed coordinates are determined prior to sampling at HQ. Coordinates are groundtruthed by Field Science in the field and reported to Science. If available in the table, Field Science coordinates should be used for sampling.

<i>Location ID</i>	<i>Description</i>	<i>Proposed Latitude</i>	<i>Proposed Longitude</i>	<i>Field Sci Latitude</i>	<i>Field Sci longitude</i>
01 - Riparian	Riparian coordinates*	46.000749	-89.704033	46.000769	-89.704042
02 - Riparian	Riparian coordinates*	45.998981	-89.702475	45.998969	-89.702494
03 - Riparian	Riparian coordinates*	45.996382	-89.70407	45.996377	-89.704100
04 - Riparian	Riparian coordinates*	45.997103	-89.699997	45.997032	-89.700316
05 - Riparian	Riparian coordinates*	45.995238	-89.698262	45.995340	-89.698223
06 - Riparian	Riparian coordinates*	45.995006	-89.70133	45.994884	-89.701363
07 - Riparian	Riparian coordinates*	45.995566	-89.704287	45.995426	-89.704279
08 - Riparian	Riparian coordinates*	45.99479	-89.706522	45.994751	-89.706463
09 - Riparian	Riparian coordinates*	45.99701	-89.705653	45.997075	-89.705715
10 - Riparian	Riparian coordinates*	45.998981	-89.706191	45.998983	-89.706328
Inlet	Inlet location from SCR	45.995178	-89.706478	45.995229	-89.706476
Outlet	Outlet location from SCR	46.000606	-89.704064	46.000596	-89.704067
C0 (buoy)	S1 sensor location from SCR	45.998269	-89.704767		

* Riparian coordinates should be approximately evenly spaced throughout the sampling area



6.2 Site-Access and Instrument Locations

NEON sites will be visited by field ecologists on a regular basis. To protect the environment near sites, several access points have been established to minimize local disturbance over the life of the project for locations (e.g., sensors and boat launches) that are accessed frequently (Figure 4, Figure 5). Field ecologists must use established paths and access points when possible to avoid causing disturbance to the site.




Figure 4. Site access and instrument locations at D05 Crampton Lake.



NEON is sponsored by the National Science Foundation and operated under a cooperative agreement by Battelle

LITTLE ROCK LAKE DOMAIN 05 GREAT LAKES



Legend

- Buoy
- Benchmark
- ⊗ Camera
- ⊕ Groundwater Well
- ▲ Inlet
- ▼ Outlet
- △ Met Station
- ⊙ Portal
- ⊥ Staff Gauge
- Access



0 37.5 75 150 225 300 Meters

WGS 1984 UTM Zone 16N
December 2017



Figure 5. Site access and instrument locations at D05 Little Rock Lake.



6.3 Riparian Sampling Locations

Riparian coordinates are determined prior to sampling at HQ. Coordinates at most sites are groundtruthed by Field Science in and reported back to Science to update Table 9 and Table 10. If available in the table, Field Science coordinates should be used for riparian sampling. Lake riparian sections are numbered from 1-10 clockwise around the lake (Figure 6, Figure 7).



Figure 6. CRAM ideal riparian sampling design




Figure 7. LIRO ideal riparian sampling design




7 REFERENCES

- Keller, M., D. S. Schimel, W. W. Hargrove, and F. M. Hoffman. 2008. A continental strategy for the National Ecological Observatory Network. *Frontiers in Ecology and the Environment*. 6: 282-284.
- Hargrove, W. W. and J. Pickering. 1992. Pseudoreplication: a sine qua non for regional ecology. *Landscape Ecology* 6: 251-258.
- Hargrove, W. W., and F. M. Hoffman. 1999. Using multivariate clustering to characterize ecoregion borders. *Computers in Science and Engineering*. 1: 18-25.
- Hargrove, W. W., and F. M. Hoffman. 2004. Potential of Multivariate Quantitative Methods for Delineation and Visualization of Ecoregions. *Environmental Management*, 34: S39–S60.
- Steele, J. H. 1978. Some comments on plankton patches. *In Spatial pattern in plankton communities*. Edited by J.H. Steele. Plenum Press, New York. pp. 1–20.



APPENDIX A LAKES

LAKE TEMPORAL SAMPLING STRATEGY



Bio Bout
 4a. Aquatic Plant Clip Harvest (Bout 2)
 4b. Aquatic Plant Point Pres/Abs (Bouts 1 & 3)
 4c. Macroinvertebrates
 4d. Periphyton, Phytoplankton
 4e. Zooplankton
 5. Sediment Chemistry (Bouts 1 & 3)
 6. Fish (Bouts 1 & 3)

SAMPLING BOUT	BOUTS/YEAR	FIELD HRS	LAB HRS
1 Sensor Maintenance	26	1 to 2	0
2 Water Chemistry, Isotopes, Dissolved Gas	12	1 to 3	1 to 3
3 Surface Microbes	12	2 to 4	0 to 1
4 Biology			
a. Aquatic Plant Clip Harvest	1	3 to 8	4 to 8, 4, 2*
b. Aquatic Plant Point Pres/Abs	2	3 to 8	0
c. Macroinvertebrates	3	4	1 to 2
d. Periphyton, Phytoplankton	3	3	4 to 8, 2, 2*
e. Zooplankton	3	4	1 to 2
5 Sediment Chemistry	2	4 to 8	1 to 4
6 Fish	2	8 to 40*	1 to 4
7 Peak Greenness			
a. Riparian Habitat Assessment	1	4 to 8	0
b. Bathymetry	<1	8 to 40*	0
8 Groundwater Chemistry	2	6 to 20*	3 to 8

Peak Greenness
 7a. Riparian Habitat Assessment
 7b. Bathymetry (every 5 yrs)

* indicates completion over multiple days



<i>Title:</i> Aquatic Site Sampling Design - NEON Domain 05		<i>Date:</i> 05/23/2019
<i>NEON Doc. #:</i> NEON.DOC.003604	<i>Author:</i> S. Parker	<i>Revision:</i> A

APPENDIX B OBSOLETE LOCATIONS, CRAM

We initially collected water chemistry samples at lake inlet, outlet, and buoy locations. After an analysis of data and discussions with external community members, we concluded that the water chemistry in lakes without true inlets and outlets were not significantly different at inlet and outlet locations relative to the buoy. Thus, to optimize OS sampling funds, we have removed lake inlet and outlet water chemistry sampling at lakes without true inlets and outlets and will only sample water chemistry at the lake buoy location.

Groundwater chemistry samples were initially collected at wells widely distributed around the site, and have since narrowed the strategy to sample two wells on the inlet side and two on the outlet side of the lake.