

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

## AOS PROTOCOL AND PROCEDURE: RIPARIAN HABITAT ASSESSMENT

<b>PREPARED BY</b>	<b>ORGANIZATION</b>	<b>DATE</b>
Brandon Jensen	AQU	03/03/2016
Kaelin Cawley	AQU	06/01/2016
Charlotte Roehm	AQU	03/14/2013
Glenn Patterson	AQU	11/04/2011

<b>APPROVALS</b>	<b>ORGANIZATION</b>	<b>APPROVAL DATE</b>
Andrea Thorpe	SCI	09/14/2016
Mike Stewart	SE	09/06/2016

<b>RELEASED BY</b>	<b>ORGANIZATION</b>	<b>RELEASE DATE</b>
Jennifer DeNicholas	CM	09/15/2016

See configuration management system for approval history.

The National Ecological Observatory Network is a project solely funded by the National Science Foundation and managed under cooperative agreement by Battelle. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.



<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

## Change Record

<b>REVISION</b>	<b>DATE</b>	<b>ECO #</b>	<b>DESCRIPTION OF CHANGE</b>
A	09/15/2016	ECO-03994	Initial release, merging streams and lakes Riparian Habitat Assessment P&Ps (NEON.DOC.001195 and NEON.DOC.001196).

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

**TABLE OF CONTENTS**

**1 OVERVIEW .....1**

1.1 Background ..... 1

1.2 Scope..... 3

1.3 NEON Science Requirements and Data Products ..... 3

1.4 Acknowledgments..... 4

**2 RELATED DOCUMENTS AND ACRONYMS .....4**

2.1 Applicable Documents ..... 4

2.2 Reference Documents..... 4

2.3 Acronyms ..... 4

2.4 Definitions..... 5

**3 METHOD .....8**

**4 DATA COLLECTION SCHEDULE .....11**

4.1 Frequency and Timing..... 11

4.2 Criteria for Determining Onset and Cessation of Field Data Collection ..... 12

4.3 Timing for Laboratory Processing and Analysis ..... 12

4.4 Data Collection Timing Contingencies ..... 12

**5 SAFETY .....13**

**6 PERSONNEL AND EQUIPMENT.....15**

6.1 Equipment..... 15

6.2 Training Requirements..... 19

6.3 Specialized Skills..... 19

6.4 Estimated Time ..... 19

**7 STANDARD OPERATING PROCEDURES:.....20**

SOP A Preparing for Field Activities ..... 20

SOP B Field Data Collection..... 22

B.1 Data Collected per Date..... 22

B.2 Data Collected per Transect..... 22

B.3 Stream Canopy Cover ..... 22

B.4 Physical Habitat Characterization ..... 24

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

B.5 Riparian Bank Characteristics ..... 28

B.6 Ending the Riparian Assessment Day..... 31

SOP C Data Entry and Verification ..... 31

SOP D Sample Shipment ..... 31

**8 REFERENCES .....32**

**LIST OF TABLES AND FIGURES**

**Table 1.** Contingent decisions..... 12

**Table 2.** Equipment list – Field equipment for wadeable stream riparian habitat assessment..... 15

**Table 3.** Equipment list – Lake and non-wadeable riparian habitat assessment procedure ..... 17

**Table 4.** Equipment list – General boating equipment..... 18

**Table 5.** Datasheets associated with this protocol..... 34

**Figure 1.** The riparian zone represents the land closest to the shoreline. .... 1

**Figure 2.** Examples of riparian habitats in different eco-climatic regions. a) D10 Central Plains; b) D18 Tundra; c) D11 Southern Plains; and d) D03 Southeast..... 2

**Figure 3.** Sketch of riparian habitat assessment set-up and scheme for a) lakes; and b) non-wadeable streams..... 9

**Figure 4.** A generic wadeable stream site layout with riparian habitat assessment locations ..... 10

**Figure 5.** An example of (a) a tandem clinometer used both as a compass and for measuring angles and (b) a laser rangefinder which uses a laser beam to determine the distance to an object and a clinometer for determining height; and (c) a convex spherical densiometer used for determining % canopy cover. 10

**Figure 6.** Preparing and reading a densiometer ..... 21

**Figure 7.** Location of 10 transects in the stream and sub-location of data collection points within each transect (Modified from Kaufmann, 2001). ..... 23

**Figure 8.** Determining the transects for riparian vegetation structure observations at each transect. .... 25

**Figure 9.** Positioning of boat/buoy for riparian assessment at each location. The distance to the shoreline and the approximate location size of the riparian zone are also shown..... 26

**Figure 10.** Location of adjacent areas around riparian assessment zone. .... 26

**Figure 11.** Determining shoreline bank angle for different bank types using a meter stick and clinometer. (A) Typical shoreline, (B) incised channel, (C) undercut stream bank, and (D) overhanging stream bank.29

**Figure 12.** Examples of shoreline revetments a) rip-rap with rock, b) gabion baskets filled with rock, c) soft armoring using root wads and logs, and d) in-stream rock J-hook for modifying flow..... 30

**Figure 13.** Location of 10 transects in the stream and sub-location of data collection points within each cross-section transect: ..... 36


**Figure 14.** Riparian vegetation structure observation transects: ..... 37

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

# 1 OVERVIEW

## 1.1 Background

Riparian areas are ecosystems adjacent to streams, rivers, lakes, wetlands, and floodplains that form complex and interrelated hydrologic systems (Verry et al., 2000; Figure 1). While riparian areas are determined by hydrologic conditions and topographic relief, they are mapped independently from either wetlands or uplands. Riparian ecosystems are transition zones between aquatic and terrestrial systems and are, hence, unique in their high biological diversity. These areas are characterized by the presence of plant communities contiguous to and affected by surface or ground water of perennial or ephemeral water bodies. Riparian areas lack the amount or duration of water usually present in wetlands, yet display wetter conditions than adjacent uplands (U.S. Fish and Wildlife Service, 1997 & 2009). These areas are characterized by vegetation comprising trees, shrubs, wildflowers, grasses, and other plants, but have distinctly different vegetation than adjacent areas or have species similar to surrounding areas that exhibit a more vigorous or robust growth form (Cowardin et al., 1979; Figure 1). High habitat complexity and diverse biological communities are formed as a result of frequent disturbances related to, for example, water movement and fluctuation, ice abrasion and sediment transport (Verry et al., 2000; Figure 2).



**Figure 1.** The riparian zone represents the land closest to the shoreline.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A



**Figure 2.** Examples of riparian habitats in different eco-climatic regions. a) D10 Central Plains; b) D18 Tundra; c) D11 Southern Plains; and d) D03 Southeast.

Water level fluctuations result in the establishment and presence of vegetation able to withstand wet and dry conditions and are an important buffer between the aquatic and terrestrial ecosystems. The riparian zone is characterized by vegetation comprising trees, shrubs, wildflowers, grasses, and other plants.

Riparian areas play a critical role in water quality by filtering or absorbing contaminants from the upland landscape, or from floodwaters, through the provision of shade in shorelines that keeps water temperatures low, as a carbon source to streams through litterfall, for creating soil stability and sediment control, and by supporting wildlife habitat and food for fish and other aquatic organisms. These areas also provide nearshore habitat structural elements, such as large woody debris (LWD), that create ideal conditions for shelter for many land and aquatic organisms and are a sink for sediments. Through infiltration, absorption, uptake, filtering, and deposition processes, riparian zones significantly reduce the amount of runoff and pollutants that reach both surface and underground water sources (Narumalani et al., 1997).

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

The overall objective of riparian habitat assessment is to provide accurate and up-to-date data of riparian cover composition, distribution, and interspersions of riparian habitats within the study area and enable identification of potential change over time. Since the remote sensing images, light detection and ranging (LIDAR), and other imagery available for interpretation may be dated and disturbances in the watershed can be rapid, it is important to consistently verify and update the riparian maps through annual site visits. Riparian habitat assessment will take place once per year during times of maximum canopy cover to match AOP flights. The time window for performing the riparian habitat assessment was chosen to match the window for AOP flights of the aquatic sites, which were determined by the time window of peak greenness for the closest terrestrial site. Peak greenness dates are defined as the range of dates where Moderate-resolution Imaging Spectroradiometer (MODIS) normalized difference vegetation index (NDVI) is within 90% of the site maximum. Most aquatic sites are co-located with terrestrial sites, therefore the peak greenness dates for these aquatic sites match exactly with the co-located terrestrial site and the flight time window. However, there are several aquatic sites that are not co-located with terrestrial sites, so by matching with the flight time window of the closest terrestrial site, there may be a slight mismatch with the actual peak greenness window at these aquatic sites (HOPB, CUPE, GUIL, MCDI, ARIK, PRIN, WLOU, REDB, BIGC, and OKSR). In addition, there are 3 aquatic sites that are too far away from a terrestrial site to be flown by AOP. For these 3 sites (BLUE, SYCA and MCRA), peak greenness was determined using the same algorithm for the aquatic site directly. The approach follows a modified version of the U.S. Environmental Protection Agency’s Environmental Monitoring and Assessment Plan (EMAP) protocol (Kaufmann, 2001), employing a randomized, systematic spatial sampling design that minimizes bias in the placement of measurements. This approach is met by making measurements at pre-defined areas that are spaced equidistant along a stream/river corridor or around a lake. Data for the riparian canopy cover extent as well as the vegetation composition and physical habitat observations are recorded at 10 equally spaced lateral transects across the channel and up to 50 m into the riparian zone within the permitted site boundaries.

**1.2 Scope**

This document provides a change-controlled version of Observatory protocols and procedures. Documentation of content changes (i.e. changes in particular tasks or safety practices) will occur via this change-controlled document, not through field manuals or training materials.

**1.3 NEON Science Requirements and Data Products**

This protocol fulfills Observatory science requirements that reside in NEON’s Dynamic Object-Oriented Requirements System (DOORS). Copies of approved science requirements have been exported from DOORS and are available in NEON’s document repository, or upon request.

Execution of this protocol generates raw data satisfying NEON Observatory scientific requirements. These data are used to create NEON data products, and are documented in the NEON Scientific Data Products Catalog (RD[03]).



Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## 1.4 Acknowledgments

This protocol is based on modified versions of the U.S. Environmental Protection Agency (2008), U.S. Environmental Protection Agency (2008, 2007), and U.S. Fish and Wildlife Service (2009, 1997).

## 2 RELATED DOCUMENTS AND ACRONYMS

### 2.1 Applicable Documents

Applicable documents contain higher-level information that is implemented in the current document. Examples include designs, plans, or standards.

AD[01]	NEON.DOC.004300	EHS Safety Policy and Program Manual
AD[02]	NEON.DOC.004316	Operations Field Safety and Security Plan
AD[03]	NEON.DOC.000724	Chemical Hygiene Plan and Biosafety Manual
AD[04]	NEON.DOC.050005	Field Operations Job Instruction Training Plan
AD[05]	NEON.DOC.014051	NEON FSU Field and Laboratory Procedures Quality Assurance Plan

### 2.2 Reference Documents

Reference documents contain information that supports or complements the current document. Examples include related protocols, datasheets, or general-information references.

RD[01]	NEON.DOC.000008	NEON Acronym List
RD[02]	NEON.DOC.000243	NEON Glossary of Terms
RD[03]	NEON.DOC.002652	NEON Level 1, Level 2, and Level 3 Data Products Catalog
RD[04]	NEON.DOC.001271	NEON Protocol and Procedure: Manual Data Transcription
RD[05]	NEON.DOC.001646	General AQU Field Metadata Sheet
RD[06]	NEON.DOC.001152	NEON Aquatic Sample Strategy Document
RD[07]	NEON.DOC.001154	AOS Protocol and Procedure: Aquatic Decontamination
RD[08]	NEON.DOC.003162	AOS Protocol and Procedure: Wadeable Stream Morphology
RD[09]	NEON.DOC.001197	AOS Protocol and Procedure: Bathymetry and Morphology of Lakes and Non-Wadeable Streams
RD[10]	NEON.DOC.003156	Datasheets for AOS Protocol and Procedure: Riparian Habitat Assessment in Wadeable Streams
RD[11]	NEON.DOC.002764	Datasheets for AOS Protocol and Procedure: Riparian Habitat Assessment in Lakes and Non-Wadeable Streams

### 2.3 Acronyms

Acronym	Definition
DBH	Diameter at Breast Height
EMAP	Environmental Monitoring and Assessment Plan (US EPA)
EPA	Environmental Protection Agency
GDD	Growing Degree Days



Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

LWD	Large Woody Debris
MODIS	Moderate Resolution Imaging Spectroradiometer
SOP	Standard Operating Procedure
USDA NRCS	United States Department of Agriculture Natural Resource Conservation Service
USGS	United States Geological Survey

## 2.4 Definitions

**Algal:** In these Aquatic Beds, algae have the greatest areal coverage. Algal Beds are widespread and diverse in the Marine and Estuarine Systems, where they occupy substrates characterized by a wide range of sediment depths and textures.

**Aquatic Bed:** The Class Aquatic Bed includes habitats where plants that grow principally on or below the surface of the water (i.e., surface plants or submergents) are the uppermost life form layer with at least 30 percent areal coverage

**Aquatic Moss:** In this Subclass, aquatic mosses have the greatest areal coverage, which are far less common than algae or vascular plants, and occur primarily in the Riverine System and Permanently Flooded systems

**Bedrock:** Where bedrock is covering 75 percent or more of the surface and less than 30 percent areal coverage of macrophytes

**Canopy Closure:** The amount of forest overstory measured with a densiometer from the center of the bankfull channel.

**Cobble-Gravel:** The unconsolidated particles smaller than stones are predominantly cobbles and gravel. Shell fragments, sand, and silt often fill the spaces between the larger particles

**Dominant:** The principal vegetative species prevalent in the area. Usually considered >30% coverage.

**Emergent:** Growing above the water’s surface. Rooted with an herbaceous stem.

**Floating Vascular:** In this Subclass, vascular plants that float freely on or below the water surface have the greatest areal coverage. Floating Vascular Beds occur mainly in the Lacustrine, Palustrine, and Riverine Systems

**Forested:** Woody vegetation more than 6 meters in height.

**Growth form:** Pertaining to the health, compactness, crowding and/or number of individuals.

**Hydric Soil:** Soils that are sufficiently wet in the upper part to develop anaerobic conditions during the growing season.

**Intermittent:** An area where moisture is only prevalent for part of the year, or where substrate is exposed. No detectable seasonal pattern in surface water.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

**Lentic:** Biotic and/or abiotic interactions in standing water.

**Lichen:** The areal coverage of lichens exceeds that of mosses. Lichen Wetlands also are a Northern Subclass.

**Lotic:** Biotic and/or abiotic interactions in moving water.

**Moss Lichen:** The Moss-Lichen Class includes areas where mosses or lichens cover at least 30 percent of substrates other than rock and where emergents, shrubs, or trees alone or in combination cover less than 30 percent

**Moss:** The areal coverage of mosses exceeds that of lichens; Moss dominated wetlands are most abundant in the far northern boreal forests and Arctic tundra

**Mud:** The unconsolidated particles smaller than stones are predominantly silt and clay, although coarser sediments or organic material may be intermixed

**Nonpersistent:** Nonpersistent emergents are emergent hydrophytes whose stems and leaves are evident above the water surface or above the soil surface if surface water is absent, only during the growing season or shortly thereafter

**Organic:** The unconsolidated material smaller than stones is predominantly organic; there is no minimum depth requirement. The organic material is dead plant tissue in varying stages of decomposition

**Perennial:** Present in all seasons of the year and usually over multiple years.

**Persistent:** Persistent emergents are emergent hydrophytes whose stems and leaves are evident all year above the surface of the water or above the soil surface if water is absent

**Phragmites australis:** Wetlands in this subclass are dominated by common reed (*Phragmites australis*)

**Revetment:** Structures built to preserve the existing uses of the shoreline, to protect the slope, and act as defense against erosion. These include rocks, concrete, trees, and in-stream structures such as vanes or J hooks to modify stream flow.

**Riparian:** Transitional areas between terrestrial and aquatic ecosystems, distinguished by gradients in biophysical conditions, ecological processes, and biota.

**Rock Bottom:** The Class Rock Bottom includes habitats with substrates having an aerial cover of stones, boulders, or bedrock 75 percent or greater and vegetative cover of less than 30 percent

**Rooted Vascular:** In this Subclass, rooted vascular plants have the greatest areal coverage, they are referred to by others as temperate grass flats

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

**Rubble:** Where there is less than 75 percent areal cover of bedrock, but stones and boulders alone or in combination with bedrock cover 75 percent or more of the area. The areal coverage of macrophytes is less than 30 percent

**Sand:** The unconsolidated particles smaller than stones are predominantly sand, although finer or coarser sediments may be intermixed

**Scrub/Shrub:** Woody vegetation <6 meters in height.

**Thalweg:** Deepest portion of a stream or river channel that carries the greatest volume of water flow

**Unconsolidated Bottom:** The Class Unconsolidated Bottom includes habitats with at least 25 percent cover of particles smaller than stones and a vegetative cover less than 30 percent

**Vegetated:** Some Streambeds, with at least 30 % plant cover, are exposed long enough to be colonized by pioneer plants that, unlike Emergent Wetland plants or Scrub-Shrub Wetland plants, are usually killed by rising water levels

**Wetland:** Areas transitional between terrestrial and aquatic environments where the water table is usually at or near the surface or the land is covered by shallow water.

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

### 3 METHOD


This protocol is intended to provide a rapid estimate of the riparian vegetation, human impacts, and bank characteristics, which buffers the permitted banks of lakes, rivers, and streams. Field data collection undertaken as part of this protocol consists of the following steps:

1. Walk or boat around the site within the permitted boundaries and conduct vegetation cover and composition observations at specified transect GPS locations.
2. Record human activities that are present in the riparian and adjacent areas.
3. Conduct lake, river, or stream bank observations.
4. At stream sites, collect canopy density measurements using a densitometer.

Any observations of the riparian areas outside of the specified Riparian Assessment locations should be recorded in the General AQU Field Metadata Sheet (RD[05]). Provide a reasonable estimate of all surveyed parameters. Lake and non-wadeable stream based riparian habitat assessment occurs by identifying 10 primary equidistant transects round the perimeter of the lake or 5 transects on both shores (left and right banks) of the non-wadeable stream for a total of 10 transects (**Figure 3**). Wadeable stream riparian habitat assessment occurs by identifying 10 transects spread equally throughout the permitted reach (**Figure 4**). The approximate locations of the 10 transects are determined prior to the riparian assessment visit and marked on the Site-Specific Riparian Locations figure in the Domain-specific Aquatic Site Sampling Strategy document. Once at the field site, technicians travel to an identified transect and establish the dimensions of the survey area. Riparian habitat assessment surveys are conducted off shore using a boat at lakes and rivers or from the center of the channel in streams. A photo is taken at this point at each transect location for future reference and archived appropriately.

The riparian portion of the plot extends up to 50 m from the shoreline/bank towards the terrestrial systems (less if the plot is visibly obstructed) and 20 m (10 m each side of central point) along the shoreline. The dimensions of the plot (up to 50 m by 20 m inland) are determined using a compass and angle referencing clinometer (**Figure 5**). In this zone, technicians record information about the vegetation type and, the height and areal coverage of trees, shrubs, and grasses using an inclinometer or rangefinder. In addition, observations of shoreline substratum (e.g., gravel, sand), the high-water mark, and bank slope characteristics are noted on the appropriate field datasheets. Field Datasheets. Anthropogenic activities and other features (e.g., buildings, land use, docks, trash, vegetation disturbances) are noted and any striking features seen adjacent to the plot. A photo is taken at each transect location for future reference and archived appropriately.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A



**Figure 3.** Sketch of riparian habitat assessment set-up and scheme for a) lakes; and b) non-wadeable streams.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A




Figure 4. A generic wadeable stream site layout with riparian habitat assessment locations



Figure 5. An example of (a) a tandem clinometer used both as a compass and for measuring angles and (b) a laser rangefinder which uses a laser beam to determine the distance to an object and a clinometer for determining height; and (c) a convex spherical densiometer used for determining % canopy cover.

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

The activities outlined in this protocol are based on modified methods developed by the U.S. Environmental Protection Agency (2008) and U.S. Fish and Wildlife Service (1997, 2009).

The following modifications have been made:

- The determination of the 20 meter horizontal widths of the riparian areas in the field are determined using a compass and angle referencing clinometer rather than by estimation.
- This method follows Ode (2007) which uses the Strickler modification (17-point) of a convex spherical densiometer to correct for overestimation of canopy density (thickness and consistency of plant foliage) that occurs with unmodified readings (Strickler, 1959).
- The maximum distance into the riparian zone that may be used as the identification area has been extended to 50 m to enable more representative identification in areas with large floodplains and shallow terrains.
- The determination of distance and height can be made using a laser rangefinder.

Standard Operating Procedures (SOPs), in Section 7 of this document, provide detailed step-by-step directions, contingency plans, data collecting and recording tips, and best practices for implementing this data collection procedure. For more details on how to operate the laser rangefinder refer to the NEON Rangefinder protocol (NEON.DOC.001717). To properly collect data, field technicians **must** follow the protocol and associated SOPs. Use NEON’s problem reporting system to resolve any field issues associated with implementing this protocol.

The value of NEON data hinges on consistent implementation of this protocol across all NEON domains, for the life of the project. It is therefore essential that field personnel carry out this protocol as outlined in this document. In the event that local conditions create uncertainty about carrying out these steps, it is critical that technicians document the problem and enter it in NEON’s problem tracking system.

The procedures described in this protocol will be audited according to the Field Audit Plan (AD[05]). Additional quality assurance will be performed on data collected via these procedures according to the NEON Data and Data Product Quality Assurance and Control Plan (AD[06]).

**4 DATA COLLECTION SCHEDULE**

**4.1 Frequency and Timing**

The riparian habitat assessment in wadeable streams shall occur once per year. Data collection timing is provided annually by Science Operations and shall be outlined in the NEON Aquatic Sampling Strategy Document (RD[06]). The time window for performing the riparian habitat assessment was chosen to match the window for AOP flights of the aquatic sites, which were determined by the time window of peak greenness for the closest terrestrial site. Peak greenness dates are defined as the range of dates where MODIS NDVI is within 90% of the site maximum. Most aquatic sites are co-located with terrestrial sites, therefore the peak greenness dates for these aquatic sites match exactly with the co-



Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

located terrestrial site and the flight time window. However, there are several aquatic sites that are not co-located with terrestrial sites, so by matching with the flight time window of the closest terrestrial site, there may be a slight mismatch with the actual peak greenness window at these aquatic sites (HOPB, CUPE, GUIL, MCDI, ARIK, PRIN, WLOU, REDB, BIGC, and OKSR). In addition, there are 3 aquatic sites that are too far away from a terrestrial site to be flown by AOP. For these 3 sites (BLUE, SYCA and MCRA), peak greenness was determined using the same algorithm for the aquatic site directly.

**4.2 Criteria for Determining Onset and Cessation of Field Data Collection**

Field data collection for riparian habitat occurs during the period of maximum vegetation cover and during safe wading conditions (see Section 5 Safety). Refer to the Site Specific Sampling Strategy Document for the data collection timeline (Appendix D and E). Maximum vegetation cover date ranges are determined using MODIS data and growing degree days (GDD). The specific times are determined using multivariate statistics and site specific historical information (see RD[06]).

**4.3 Timing for Laboratory Processing and Analysis**

There is no domain lab processing for this protocol.

**4.4 Data Collection Timing Contingencies**

Table 1. Contingent decisions

Delay/Situation	Action	Outcome for Data Products
Hours	If equipment stops functioning during data collection continue from where the data collection left off as soon as possible or restart if more than one week has passed.	No adverse outcome.
	If there is no access to the shore (due to safety issues, dense vegetation, sensitive ecosystem, unsafe ground conditions or because part of the segment is on private property), record conditions and comments on the field data sheet. Proceed by walking, driving, or boating along the edge of the water body and make observations where possible.	No adverse outcome.
	If the depth of water in the thalweg of the stream is too deep or the flow is too fast for safe wading, record conditions and comments on your field datasheet. If the edge of the lake or non- wadeable stream is not accessible within 10 meters for the riparian assessment due to, for example, thick emergent aquatic vegetation, estimate the closest distance to the lake shore. Then continue as detailed in the protocol.	No adverse outcome.
Days	If the weather and wading creates unsafe conditions	No adverse outcome.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

	become unsafe during field activities, stop and resume as soon as safe conditions for wading are met. Unless physical change or more than one week has passed as a result of the change in weather and flow (i.e. bank erosion, movement of large woody debris etc.) data collection should resume from where it was left off and completed within the window provided for the Riparian Habitat Assessment protocol.	
7 Days or More	Restart assessment if 7 or more days have passed since the previous measurements were started or if the vegetation cover has changed significantly (loss of leaves or leaf on).	No adverse outcome.

**5 SAFETY**

This document identifies procedure-specific safety hazards and associated safety requirements. It does not describe general safety practices or site-specific safety practices.

Personnel working at a NEON site must be compliant with safe field work practices as outlined in the Operations Field Safety and Security Plan (AD[02]) and EHS Safety Policy and Program Manual (AD[01]). Additional safety issues associated with this field procedure are outlined below. The Field Operations Manager and the Lead Field Technician have primary authority to stop work activities based on unsafe field conditions; however, all employees have the responsibility and right to stop their work in unsafe conditions.

Activities in streams should only be performed when flow conditions are safe. Do not attempt to wade a stream where velocity x depth is  $\geq 10 \text{ ft}^2/\text{s}$  ( $0.93 \text{ m}^2/\text{s}$ ).

In addition, the following safety guidelines are provided:

1. Due to site-specific hazards that may be encountered in accessing the shoreline technicians may complete observations without dismounting from the vessel. In addition, the technicians are required not to put hands and feet in waters where alligators are present and to make sure a safe distance from hazards is maintained.
2. All personnel must be wearing a personal flotation device (PFD) prior to entering and while in the boat.
3. Personnel in wadeable streams shall work in teams of 2, at a minimum, and shall wear personal flotation devices (PFDs) when wading where velocity, depth or obstructions may create a hazardous work environment (e.g. stream is fast moving, the depth is greater than knee-high in slow moving stream, or the stream may present slip or fall hazards).
4. All employees shall have access to a form of communication with other team members such as a two-way radio.

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 09/15/2016
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

5. Technicians should be aware of any site-specific hazards and the waters of that particular location (i.e. current status, tidal charts, etc.)

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## 6 PERSONNEL AND EQUIPMENT

### 6.1 Equipment

The following equipment is needed to implement the procedures in this document. Equipment lists are organized by task. They do not include standard field and laboratory supplies such as charging transects, first aid kits, drying ovens, ultra-low refrigerators, etc.

**Table 2.** Equipment list – Field equipment for wadeable stream riparian habitat assessment

Item No.	R/S	Description	Purpose	Quantity	Special Handling
<b>Durable items</b>					
	S	Personal flotation devices	Safe wading	2	N
	R	GPS unit	Navigating to sampling locations and establishing transects	1	N
	R	Surveyor's tape measure 50 m or 100 m	Establishing transects	1	N
	R	Permanent transect marker rods	Establishing transects	22	N
MX102980	R	Suunto Clinometer	Determining shoreline bank angle	1	N
MX100348	R	Spherical Densiometer	Determining canopy cover	1	N
MX104742	R/S	Rangefinder (shared with FSU)	Determining angles, distance and height (alternate to using the clinometer)	1	N

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

Item No.	R/S	Description	Purpose	Quantity	Special Handling
	R	Folding meter stick	Measuring stream water depth	1	N
	R	Stream base map	Navigating to sampling locations	2	N
	R	Quick References (Appendix B)	Protocol field reference	1	N
	R	Camera	Photographing riparian locations	1	N
	R	Calculator	Calculating stream cover density	1	N
	R	First aid kit		1	N
	R	Waders	Safe wading	2	N
<b>Consumable items</b>					
	R	Canopy Cover Datasheets	Recording data	1	N
	R	Physical Habitat Characterization Datasheets	Recording data	10	N
	S	Field notebook	Recording field notes	2	N

R/S=Required/Suggested

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

**Table 3.** Equipment list – Lake and non-wadeable riparian habitat assessment procedure

Item No.	R/S	Description	Purpose	Quantity	Special Handling
<b>Durable items</b>					
	R	Meter stick (1 m)	Use with clinometer to measure bank angle	1	N
	R	Suunto Clinometer	Determining shoreline bank angle	1	N
	R	Lake or Non-wadeable stream map	Navigating to sampling locations	1	N
	R	Quick reference field operations handbook	References	1	N
	R	Camera	Photographing riparian locations	1	N
	R	Sonar with GPS antenna	Navigating to sampling locations	1	N
	R	Memory card	For camera and sonar	3	N
	R	Handheld GPS unit	Navigating to sampling locations	1	N
<b>Consumable items</b>					
	R	Physical Habitat Characterization Forms	Recording physical habitat characterization data	12	N
	R	Field notebook	Recording field notes	2	N
	R	12 V small batteries	For GPS unit	1	N

R/S=Required/Suggested

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 09/15/2016
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

**Table 4.** Equipment list – General boating equipment

Item No.	R/S	Description	Purpose	Quantity	Special Handling
<b>Durable items</b>					
	R	Boat		1	Y
	R	Anchor with rope		1	N
	R	Oars		2	N
	R	Trolling Electric Motor		1	Y
	R	Battery (12 volt)		1	Y
	R	Safety kit for boat (e.g., flares, bailer, float with rope)		1	Y
	R	First Aid Kit		1	N
	R	Personal Flotation Devices (PFDs)		1 per person	N
<b>Consumable items</b>					
		(none)			

R/S=Required/Suggested



<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 06/22/16
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

**6.2 Training Requirements**

All technicians must complete required safety training and protocol-specific training for safety and implementation of this protocol as required in Field Operations Job Instruction Training Plan (AD[04]).

All personnel required to operate a boat shall be trained through an approved program. All others shall be aware of boating safety procedures.

Personnel are to be trained in stream riparian habitat assessment measurements and safe working practices for stream work, to include use of waders and PFDs, where applicable.

**6.3 Specialized Skills**

Where applicable, personnel will be licensed to operate a boat and able to safely handle a motor and drive a boat safely.

Personnel should be able to use a clinometer to compute angles and distances.

**6.4 Estimated Time**

The time required to implement a protocol will vary depending on a number of factors, such as skill level, system diversity, environmental conditions, and distance between measurement locations. The timeframe provided below is an estimate based on completion of a task by a skilled two-person team (i.e., not the time it takes at the beginning of the field season). Use this estimate as framework for assessing progress. If a task is taking significantly longer than the estimated time, a problem ticket should be submitted.

We estimate that riparian habitat assessment for all transects or points at a site requires 2 technicians for 6-8 hours plus travel to and from the site.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## 7 STANDARD OPERATING PROCEDURES:

See Appendix B for Quick References and Appendix C for Reminders.

### SOP A Preparing for Field Activities

*(Modified from U.S. EPA 2007)*

1. Ensure memory cards are blank. If files are present, confirm data and photos have been uploaded prior to deleting files.
2. Verify all equipment is available and functioning properly.
3. Fully charge all batteries and electronic equipment. **IMPORTANT:** Ensure all batteries and electronic equipment are fully charged, even if new.
4. Ensure all equipment that will touch the water is decontaminated with bleach (see AOS Protocol and Procedure: Aquatic Decontamination Protocol RD[07]).
5. Print applicable Datasheets (1 canopy cover datasheet for streams sites and 10 Physical Habitat Characterization Datasheets for all types of sites).
6. The cross-section transect locations will be established at NEON headquarters for lakes, river, and stream sites (**Figure 3, Figure 4, Figure 7**). Before heading out to the field, confirm the location of cross-section transects. Ten cross-section transects are placed at equal intervals along the full permitted reach length, but not greater than 100 m apart within a 1 km NEON aquatic reach at stream and river sites. For lake sites the transects will be spaced as evenly as possible around the lake perimeter. These transects shall not be located on the same transects as the biological sampling cross-sections. Install and monument the transect locations with the permanent monument markers on either the right or left bank above the high water mark.
7. **For stream sites** (For lake sites, skip to step 8), where canopy cover will be estimated using a densiometer:
  - a. Prepare the densiometer before heading out into the field (Figure 6). Place tape over the bottom 2/3 of the densiometer. A “V” shape is outlined to help constrain readings in the upper 1/3 of the densiometer that are otherwise representative of reflection of the vegetation to the side as well as directly above. This increases accuracy of readings to vegetation density only directly above the viewer (Figure 6). Dots should be added with a permanent marker where the lines intersect on the **convex** densiometer.
  - b. The densitometer should be semi-permanently mounted to a monopod. When reading from the densitometer in the center of the stream the mono-pod, in conjunction with the bubble level, will allow for the technician to maintain the densitometer position as they move around it to face all four directions. The monopod can be marked to ensure a height of 30 cm above the water surface, when taking measurements in the center of the stream, or ground surface, when taking measurements at either bank.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A




Figure 6. Preparing and reading a densiometer

8. Familiarize yourself with the most common species of plants in your domain/sites.
  - a. Resources available include provided reference guides and the USDA NRCS online guide "The PLANTS Database."
    - 1) Under the 'I want to...' tab, select **"See list of the plants in my state."**
    - 2) Under the 'I want to...' tab, select **"Learn about all the endangered plants of the U.S."**
9. Familiarize yourself with the clinometer, rangefinder, and spherical densiometer, if applicable (Appendix G). Refer to the Forestry-Suppliers Technical Bulletin M0003 Appendix F.



Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## SOP B Field Data Collection

### B.1 Data Collected per Date

Each day that riparian assessment field measurements are made the stage of the stream, lake, or river should be recorded. This is done by reading the level of water on a staff gage.

### B.2 Data Collected per Transect

For each stream or river transect indicate whether the measurements are being recorded for the left of right bank. At stream sites only, record the water depth in the thalweg where the transect crosses the stream. For lake sites, no transect specific information, other than the actual transect ID, needs to be recorded.


### B.3 Stream Canopy Cover

Stream canopy cover measurements are only collected at wadeable stream sites. If you are performing riparian assessment at a lake or river site, please proceed to SOP B.4 Physical Habitat Characterization. This method is based on the Water Quality Monitoring Technical Guidebook from Oregon’s Watershed Enhancement Board (OWEB 1999) and California Department of Pesticide Regulation (2004).



Canopy cover over the stream is measured at each of the 10 stream cross-section transects using a monopod mounted **convex** spherical densiometer. Six measurements are obtained at each transect: 4 measurements in the four directions (upstream, right bank, left bank, and downstream) from the mid-channel (stream center; equal distance from left and right banks); 1 measurement from the left bank and 1 from the right bank, both bank measurements are made facing away from the stream, perpendicular to shore, into the canopy for a total of 6 measurements.

1. Using the Canopy Cover datasheet, enter the appropriate header data (site ID, technician ID, collection date, time, protocol number, and revision number).
2. Begin at the farthest downstream location. Affix the tape measure along the 1<sup>st</sup> transect from the left to the right bank.
3. Stand in the 1<sup>st</sup> transect (Figure 7) at mid-channel facing upstream.



**Figure 7.** Location of 10 transects in the stream and sub-location of data collection points within each transect (Modified from Kaufmann, 2001).



4. Measure and record overstory canopy cover from the stream center (mid-channel).
  - a. Set up the densiometer 0.3 meters above the water surface. This avoids errors stemming from people of different heights and water of varying depth.
  - b. Tilt the monopod such that the densiometer is level, use the level bubble indicator and hold so the reflection of the top of your head just touches the point of the “V”, as in **(Figure 6)**.
  - c. Count the number of points covered by vegetation and enter this value in the datasheet under “# of points covered”. Values will be between 0 for completely open and 17 for completely covered canopy.
  - d. Take four densiometer readings from the stream center at each of the 10 transects, once each while facing upstream, left bank, downstream, and right bank.
  - e. As you change direction be sure to keep the base of the monopod in the same location and pivot around the densitometer so that it does not change location. Be sure to read the densitometer when it is level after pivoting.
  - f. Record the values on the Canopy Cover Datasheet under “Center-Upstream”, “Center-River Left”, “Center-Downstream”, and “Center-River Right”.


Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

5. Measure and record canopy cover percent standing 0.3 m from the water’s edge on the LEFT bank facing into the terrestrial portion of the riparian transect, away from the stream. Take only one reading at this bank. Set up the monopod such that the densiometer is 0.3 m above the land surface and it is level.
  - a. Record the values on the Canopy Cover Datasheet under “Left Bank-River Left”.
6. Measure and record canopy cover percent standing 0.3 m from the water’s edge on the RIGHT bank facing into the terrestrial portion of the riparian transect, away from the stream. Take only one reading at this bank. Set up the monopod such that the densiometer is 0.3 m above the land surface and it is level.
  - a. Record the values on the Canopy Cover Datasheet under “Right Bank-River Right”.

#### B.4 Physical Habitat Characterization

1. Record observations on the site-specific riparian habitat assessment datasheet. The focus of this riparian vegetation habitat assessment is to record observations of riparian vegetation and habitat; do not record littoral vegetation observations.
2. Using the *Physical Habitat Characterization Datasheet*, enter the appropriate header data (site ID, technician ID, collection date and time, and transect number).
3. **In streams**, stand in the middle of the stream (or where safe and both banks are in view) on the transect. Take a photo of the riparian zone facing toward the shoreline at each transect location; one facing the left bank and one facing the right bank. In the photo include the top of the current waterline and as much of the riparian habitat as possible (widest angle and in landscape orientation); record the photo number.
4. **In lakes and rivers**, Position the boat at a distance of 10 m (~30 ft., offshore), anchor if necessary. Drop the float overboard in the spot where data collection is occurring. Use this as a mid-point marker for the 20 m horizontal distance for measuring the littoral areas and the perpendicular point from the shoreline, from where to delineate the littoral and riparian habitats (**Figure 9** and **Figure 10**).
5. In Part A of the datasheet, enter the full classification code following the Hierarchical Determination for Dominant System and USDA PLANTS symbol. Refer to Appendix E, Riparian Classification System for detailed instructions to properly document the dominant plants.


Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A




**Figure 8.** Determining the transects for riparian vegetation structure observations at each transect.



Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A



**Figure 9.** Positioning of boat/buoy for riparian assessment at each location. The distance to the shoreline and the approximate location size of the riparian zone are also shown.



**Figure 10.** Location of adjacent areas around riparian assessment zone.

- Using Part B of the datasheet, record estimates of riparian vegetation structure and composition. Include observations at three levels: the canopy layer (> 5 m high), understory (0.5 – 5.0 m high), and the ground cover layer (< 0.5 m high). Additionally, observations of human land use activities and disturbances shall be recorded. Observations of riparian cover may total

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

up to 300 %, 100 % each for canopy, understory, and ground cover (e.g. canopy cover is near 100% and there is a very dense understory).

7. For each of the upper two vegetation layers (canopy and understory only), record the type of vegetation (including trees, shrubs, herbs, grasses, and forbs) that occur by indicating the presence of Deciduous (D), Coniferous (C), Broadleaf Evergreen (E), Mixed (M), or None (N). A mixed vegetation layer is considered mixed if the more than 10% of the areal coverage is composed of an alternate vegetation type. Record the semi-quantitative (observational) measurements on the Physical Habitat Characterization datasheet, for each of the three vegetation layers and human influences.

8. Vegetation Structure and Composition

- a. Layers of Riparian Vegetation:

- 1) **Canopy** >5 m height

- a) Big Trees >0.3m DBH
      - b) Small Trees < 0.3m DBH

- 2) **Understory** 0.5 to 5 m height

- a) Woody shrubs and saplings
      - b) Tall herbs, grasses and forbs

- 3) **Ground Cover** <0.5 m height

- a) Woody shrubs
      - b) Herbs, grasses, and forbs
      - c) Standing water and inundated vegetation
      - d) Bare dirt or duff

- 4) **Human**

- a) Buildings, roads, boat ramps, lawns/parks, agriculture, industry, agricultural, industrial, cleared lot/pavement, or trash.

- b. Categorization of Cover:

- 1) Absent = 0%
    - 2) Sparse > 0 - < 10 %
    - 3) Moderate > 10 - < 40 %
    - 4) Heavy > 40 - < 75 %
    - 5) Very Heavy > 75 %



- c. Complete a separate *Physical Habitat Characterization Datasheet* for each transect.

9. Record, on the datasheet, other observations made regarding areas that are *adjacent* (outside of the 10 m upstream and downstream transect) to those horizontal stretch being assessed. This is particularly important where human disturbance is evident (i.e. house, road, boat ramp, lawn, power lines, garbage etc.).

10. Make every reasonable attempt to record physical habitat observations and measurements for all 10 riparian transects. Where this is impossible, indicate this information in the metadata and make a comment field on the Physical Habitat Characterization Datasheets (RD[09]) to clearly


Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

indicate that no observations could be made for a given feature at that particular riparian transect (EPA 2007; Field Operations Manual).

11. If you are unsure of or unable to determine the canopy cover type, take a photo of the site and specimen and note this in the field metadata sheet.

### B.5 Riparian Bank Characteristics

1. Using Part C of the Physical Habitat Characterization datasheet enter data from the following observations.
  - a. With the clinometer and the meter stick, determine and record the (approximate) *Bank Angle* in Part C (1<sup>st</sup> box) of the *Physical Habitat Characterization Datasheet* (Appendix A).
  - b. Lay the meter ruler against the bank with one end towards the water. Half of the meter stick (0.5 m) should be resting on the bank to determine the angle. Place the clinometer on the meter stick; read and record the bank angle in degrees (**Figure 11**).
  - c. There may be a variety of angles. If two predominate, tick both and estimate which one is dominant. If a boulder or log is present along the bank being measured, record a bank angle from an adjacent and representative slope.
    - 1) Vertical/undercut > 75°
    - 2) Steep 30 – 75°
    - 3) Gradual < 30°



**Figure 11.** Determining shoreline bank angle for different bank types using a meter stick and clinometer. (A) Typical shoreline, (B) incised channel, (C) undercut stream bank, and (D) overhanging stream bank.

2. Determine the *Bank Revetment* (artificial bank reinforcement structures) through observations and record in Part C (2<sup>nd</sup> box) of the *Physical Habitat Characterization Datasheet* (Appendix A).
  - 1) Hard Bank – Rip-rap (**Figure 12**), rock covering over the bank, usually large angular boulders, concrete blocks or rectangular gabion wire baskets (**Figure 12**)
  - 2) In-Stream – Human-placed large woody debris, tree root wads (**Figure 12**), rock vanes, or J-hooks (**Figure 12**) in the stream channel to modify flow
  - 3) Other – Soft armoring of the shore using planted trees, shrubs, and human-placed logs intended to stop erosion
  - 4) None

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A



**Figure 12.** Examples of shoreline revetments a) rip-rap with rock, b) gabion baskets filled with rock, c) soft armoring using root wads and logs, and d) in-stream rock J-hook for modifying flow.

3. Determine the Dominant and Subdominant *Bank Texture* through observations and note this in Part C (3<sup>rd</sup> box) of the *Physical Habitat Characterization Datasheet* (Appendix A).
  - a. Bedrock – very resistant to erosion
  - b. Boulder/Cobble – boulders >10”/cobble 2.5 to 10”. Moderate resistance to erosion
  - c. Gravel – 0.08-2.5 “. Moderate to high erodibility when dominant component.
  - d. Sand – 0.004 – 0.08”. High bank erodibility when dominant component
  - e. Silt – 0.00008 – 0.004 “. Non-cohesive with high to very high erodibility
  - f. Clay – < 0.00008”. Cohesive clays are relatively resistant to erosion
  - g. Mix – Variety of particle sizes. i.e. glacial till

<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 06/22/16
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

## **B.6 Ending the Riparian Assessment Day**

Equipment maintenance, cleaning, and storage

- a. Recharge all batteries.
- b. Ensure all equipment is properly decontaminated and dry prior to storage according to the NEON Aquatic Decontamination Protocol (RD[07]).

### **SOP C Data Entry and Verification**

As a best practice, field data collected on paper datasheets should be digitally transcribed within 7 days of collection or the end of a bout (where applicable). However, given logistical constraints, the maximum timeline for entering data is within 14 days of collection or the end of a bout (where applicable). See TOS Protocol and Procedure: Manual Data Transcription (RD[04]) for complete instructions regarding manual data transcription.

### **SOP D Sample Shipment**

There is no sample shipment for this protocol.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## 8 REFERENCES

- Bushnell Outdoor Products. 2011. Bushnell Laser Rangefinder G-Force 1300 ARC (Model 201965, 201966) Product Manual. <http://www.bushnell.com/getmedia/32804b08-ac38-4bc2-a70a-2b3330b431e1/gforce1300arc-201965-201966.pdf?ext=.pdf> (accessed January 7, 2016).
- California Department of Pesticide Regulation: Environmental Monitoring Branch. 2004. SOP Number: FSOT.002.01. Standard Operating Procedure, Instructions for the Calibration and Use of a Spherical Densimeter. California Department of Pesticide Regulation, Environmental Monitoring Branch. 4pp
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service, Washington, D.C. FWS/OBS-79/31.
- Forestry Suppliers. 2006. How to use a Clinometer. Technical Bulletin M0003. pp 2. [http://www.forestry-suppliers.com/t01\\_pages/pdfs/M0003.pdf](http://www.forestry-suppliers.com/t01_pages/pdfs/M0003.pdf) (accessed January 7, 2016).
- Forestry Suppliers. 2013. Using Forest Densimeters. [http://www.forestry-suppliers.com/Documents/1450\\_msdms.pdf](http://www.forestry-suppliers.com/Documents/1450_msdms.pdf) (accessed January 7, 2016).
- Kaufmann, P.R, 2001. Physical Habitat Characterization. Section 7 EMAP, Western Pilot Field Operations Manual for Wadeable Streams., Section 7 (Physical Habitat Characterization). Pp. 1-58.
- Narumalani, S., Y. Zhou, and J.R. Jensen. 1997. Application of Remote Sensing and Geographic Information Systems to the Delineation and Analysis of Riparian Buffer Zones. *Aquatic Botany*, 58 (3-4): 393-409
- Ode, P. 2007 Standard Operating Procedures for Collecting Benthic Macroinvertebrate Samples and Associated Physical and Chemical Data for Ambient Bioassessments in California. State Water Resources Control Board, Surface Water, Ambient Monitoring Program. 48pp.
- Oregon's Watershed Enhancement Board. 1999. Water Quality Monitoring Technical Guidebook. Chapter 14: Stream shade and canopy cover monitoring methods.
- Strickler, Gerald S. 1959. Use of the densimeter to estimate density of forest canopy on permanent sample plots. USDA Forest Service, Pacific Northwest Forest and Range Exp. Sta. Research Note 180, Portland, Oregon, 5 pp.
- U.S. Department of Agriculture, Natural Resource Conservation Service. 2015. The PLANTS Database. National Plant Data Team, Greensboro, NC 27401-4901 USA. <http://plants.usda.gov> (accessed January 7, 2016).
- U.S. Environmental Protection Agency. 2007. Survey of the Nation's Lakes. Field Operations Manual. EPA 841-B-06-xxx. U.S. Environmental Protection Agency, Washington, DC. , 121pp.



<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 06/22/16
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

U.S. Environmental Protection Agency. 2008. Development for a Statewide Wetland and Riparian Mapping, Assessment and Monitoring Program 2009-2015. 29 pp.

U.S. Fish and Wildlife Service. 1997. A system for mapping riparian areas in the western United States. U. S. Fish and Wildlife Service., Washington, DC. 15pp.

U.S. Geological Survey. 2016. The National Map. <http://viewer.nationalmap.gov/viewer> (accessed January 7, 2016).

Verry, E.S., J.W. Hornbeck, and A. Dolloff. 2000. Riparian Management in Forests of the Continental Eastern United States. Eds. Verry, Hornbeck & Dolloff. Lewis Publishers, Boca Raton.



<i>Title:</i> AOS Protocol and Procedure: Riparian Habitat Assessment		<i>Date:</i> 06/22/16
<i>NEON Doc. #:</i> NEON.DOC.003826	<i>Author:</i> B. Jensen	<i>Revision:</i> A

**APPENDIX A DATASHEETS**

The following datasheets are associated with this protocol:

**Table 5.** Datasheets associated with this protocol

<b>NEON Doc. #</b>	<b>Title</b>
NEON.DOC.001646	General AQU Field Metadata Sheet
NEON.DOC.003156	Datasheets for AOS Protocol and Procedure: Riparian Habitat Assessment in Wadeable Streams
NEON.DOC.002764	Datasheets for AOS Protocol and Procedure: Riparian Habitat Assessment in Lakes and Non-Wadeable Streams

These datasheets can be found in the NEON Document Warehouse.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## APPENDIX B QUICK REFERENCES

### B.1 Steps for Riparian Habitat Assessment

**Step 1** – Check the riparian habitat assessment kit to make sure all supplies are packed.

**Step 2** – Print out Physical Habitat Characterization Datasheets (RD[09])

**Step 3** – Ensure the General AQU Field Metadata Sheet (RD[06]) is completed on each field site visit.

**Step 4** – At stream sites, measure Stream Canopy Cover Density:

1. Measure canopy cover over the stream at each of the 10 transects using a convex spherical densiometer.
2. Take measurements in the four directions at mid-channel, and 1 measurement from the left bank and right bank facing away from the stream into the terrestrial portion of the riparian transect for a total of 6 measurements.
3. Place tape over the bottom 2/3 of the densiometer and measure and record the number of filled points.

**Step 4** – After returning to the boat access point, navigate to the first riparian assessment location.

**Step 5** – Riparian Vegetation Structure:

Stand in the center of the stream and make qualitative (observational) measurements


- a. Layers of Riparian Vegetation
- b. Categorization of Cover

**Step 6** – Riparian Bank Characteristics:


1. Shoreline Bank Angle
2. Bank Revetment
3. Bank Texture
4. Estimate and record the vertical distance between present water level and the high water mark
5. Take photos at all transect locations

## B.2 Locations for Riparian Habitat Assessment

**Figure 13.** Location of 10 transects in the stream and sub-location of data collection points within each cross-section transect:




**Figure 14.** Riparian vegetation structure observation transects:




Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

### 1. Lakes



### 2. Non-Wadeable Streams



Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## APPENDIX C REMINDERS

### Before heading into the field: Make sure you...

- Collect and prepare all equipment and ensure batteries are charged.
- Pre-print the *Canopy Cover* and *Physical Habitat Characterization Datasheets* and the Riparian Locations map from the AOS Site Specific Sampling Strategies document.
- Familiarize yourself with the most common and sensitive species of plants in your domain/sites.

### While collecting data: Be sure to...

- Measure canopy cover at wadeable stream sites.
- Using a convex spherical densiometer collect six measurements at each transect: 4 measurements in the four directions at mid-channel (upstream, left bank, downstream, and right bank), and 1 measurement from the left bank (looking away from the stream into the terrestrial portion of the riparian transect) and right bank (looking away from the stream into the terrestrial portion of the riparian transect) for a total of 6 measurements. REMINDER left and right banks are defined while looking downstream.
- Hold the densiometer 0.3 meters above the water surface.
- At all sites complete a separate *Physical Habitat Characterization Datasheet* for each transect.
- If unsure of or unable to determine the plant canopy cover type or encounter other issues, take a photo of the site and/or specimen and note this in the field metadata; include a comment on the Physical Habitat Characterization Datasheet as well.

### Ending the day: Be sure to...

- Recharge all batteries.
- Ensure all equipment is properly decontaminated and dry prior to storage.

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

## APPENDIX D ESTIMATED DATES FOR ONSET AND CESSATION OF DATA COLLECTION

See the DXX Site Specific Sampling Strategy Document on AQU's NEON intranet site.

## APPENDIX E SITE-SPECIFIC INFORMATION

### Hierarchical Determination of Dominant System

This classification follows the USFWS (2009) "A system for mapping riparian areas in the western United States" and incorporates the plant identification system developed by the USDA PLANTS Database (2015).

- 1) **Subsystem:** Defines three categories reflecting the water source for the riparian area
  - a) stream (moving water, shallow enough to safely wade)
  - b) river (moving water, too deep or swift to safely wade)
  - c) lake (standing water)
- 2) **Class:** Describes the dominant non-hydrophytic life form of riparian vegetation. **Subclass:** Further describes the Class, defined under each applicable class.
  - a) rock bottom - The Class Rock Bottom includes habitats with substrates having an aerial cover of stones, boulders, or bedrock 75 percent or greater and vegetative cover of less than 30 percent
    - i) bedrock - Where bedrock is covering 75 percent or more of the surface and less than 30 percent areal coverage of macrophytes
    - ii) rubble - Where there is less than 75 percent areal cover of bedrock, but stones and boulders alone or in combination with bedrock cover 75 percent or more of the area. The areal coverage of macrophytes is less than 30 percent
  - b) unconsolidated bottom - The Class Unconsolidated Bottom includes habitats with at least 25 percent cover of particles smaller than stones and a vegetative cover less than 30 percent
    - i) Cobble-Gravel - The unconsolidated particles smaller than stones are predominantly cobbles and gravel. Shell fragments, sand, and silt often fill the spaces between the larger particles
    - ii) Sand - The unconsolidated particles smaller than stones are predominantly sand, although finer or coarser sediments may be intermixed
    - iii) Mud - The unconsolidated particles smaller than stones are predominantly silt and clay, although coarser sediments or organic material may be intermixed
    - iv) Organic - The unconsolidated material smaller than stones is predominantly organic; there is no minimum depth requirement. The organic material is dead plant tissue in varying stages of decomposition
    - v) Vegetated - Some Streambeds, with at least 30 % plant cover, are exposed long enough to be colonized by pioneer plants that, unlike Emergent Wetland plants or Scrub-Shrub Wetland plants, are usually killed by rising water levels
  - c) aquatic bed - The Class Aquatic Bed includes habitats where plants that grow principally on or below the surface of the water (i.e., surface plants or submergents) are the uppermost life form layer with at least 30 percent areal coverage

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

- i) Algal - In these Aquatic Beds, algae have the greatest areal coverage. Algal Beds are widespread and diverse in the Marine and Estuarine Systems, where they occupy substrates characterized by a wide range of sediment depths and textures.
- ii) Aquatic Moss - In this Subclass, aquatic mosses have the greatest areal coverage, which are far less common than algae or vascular plants, and occur primarily in the Riverine System and Permanently Flooded systems
- iii) Rooted Vascular - In this Subclass, rooted vascular plants have the greatest areal coverage, they are referred to by others as temperate grass flats
- iv) Floating Vascular - In this Subclass, vascular plants that float freely on or below the water surface have the greatest areal coverage. Floating Vascular Beds occur mainly in the Lacustrine, Palustrine, and Riverine Systems
- d) moss lichen - The Moss-Lichen Class includes areas where mosses or lichens cover at least 30 percent of substrates other than rock and where emergents, shrubs, or trees alone or in combination cover less than 30 percent
  - i) Moss - The areal coverage of mosses exceeds that of lichens; Moss dominated wetlands are most abundant in the far northern boreal forests and Arctic tundra
  - ii) Lichen - The areal coverage of lichens exceeds that of mosses. Lichen Wetlands also are a Northern Subclass.
- e) forested - woody vegetation usually > 6 m in height
  - i) Broad-leaved Deciduous - In this Subclass, broad-leaved deciduous species have the greatest areal coverage in the tree layer
  - ii) Needle-leaved Deciduous - In this Subclass, needle-leaved deciduous species have the greatest areal coverage in the tree layer
  - iii) Broad-leaved Evergreen - In this Subclass, broad-leaved evergreen species have the greatest areal coverage in the tree layer
  - iv) Needle-leaved Evergreen - In this Subclass, needle-leaved evergreen species have the greatest areal coverage in the tree layer
  - v) Dead - This Subclass includes stands of dead woody plants 6 m in height or taller, regardless of their density, with less than 30 percent cover of living vegetation
- f) scrub/shrub - woody vegetation usually < 6 m in height
  - i) Broad-leaved Deciduous - In this Subclass, broad-leaved deciduous species have the greatest areal coverage in the scrub/shrub layer
  - ii) Needle-leaved Deciduous - In this Subclass, needle-leaved deciduous species have the greatest areal coverage in the scrub/shrub layer
  - iii) Broad-leaved Evergreen - In this Subclass, broad-leaved evergreen species have the greatest areal coverage in the scrub/shrub layer
  - iv) Needle-leaved Evergreen - In this Subclass, needle-leaved evergreen species have the greatest areal coverage in the scrub/shrub layer
  - v) Dead - This Subclass includes stands of dead woody plants < 6 m in height or taller, regardless of their density, with less than 30 percent cover of living vegetation



Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

- g) emergent - erect, rooted vegetation with herbaceous stems
  - i) Persistent - Persistent emergents are emergent hydrophytes whose stems and leaves are evident all year above the surface of the water or above the soil surface if water is absent
  - ii) Nonpersistent - Nonpersistent emergents are emergent hydrophytes whose stems and leaves are evident above the water surface or above the soil surface if surface water is absent, only during the growing season or shortly thereafter
  - iii) Phragmites australis - Wetlands in this subclass are dominated by common reed (*Phragmites australis*)
- 3) **Dominance Type:** Refers to vegetative species within the mapping unit. The description for the dominant species type will follow the USDA PLANTS Database (2015) plant symbology, e.g.:
  - a) **(PODE3)** *Populus deltoids* W. Eastern cottonwood
  - b) **(ALRU2)** *Alnus rubra* Bong. Red alder

### Classification of Riparian Vegetation

- 1) Layers of Riparian Vegetation:
  - a) **Canopy** > 5 m height
    - i) Big Trees > 0.3 m DBH
    - ii) Small Trees < 0.3 m DBH
  - b) **Understory** 0.5 to 5 m height
    - i) Woody shrubs and saplings
    - ii) Tall herbs, grasses and forbs
  - c) **Ground Shrubs** < 0.5 m height
    - i) Woody shrubs
    - ii) Herbs, grasses and forbs
    - iii) Standing water and inundated vegetation
    - iv) Bare dirt or duff
  - d) **Human**
    - i) Buildings, roads, ramps, lawns, agricultural, industrial etc.
- 2) Categorization of Cover:
  - a) Absent 0%
  - b) Sparse > 0 - < 10 %
  - c) Moderate > 10 - < 40 %
  - d) Heavy > 40 - < 75 %
  - e) Very Heavy > 75 %

### Bank Features

- 1) Bank angle
  - a) Vertical/Undercut >75° vertical
  - b) Steep 30 – 75°

Title: AOS Protocol and Procedure: Riparian Habitat Assessment		Date: 06/22/16
NEON Doc. #: NEON.DOC.003826	Author: B. Jensen	Revision: A

- c) Gradual <math><30^\circ</math>
- 2) Bank Revetment
  - a) Hard Bank – Rock covering over the bank, usually large angular boulders, concrete blocks or rectangular gabion wire baskets
  - b) In-Stream – Human-placed large woody debris, tree root wads, rock vanes, or J-hooks in the stream channel to modify flow
  - c) Other – Soft armoring of the shore using planted trees, shrubs, and human-placed logs intended to stop erosion
  - d) None
- 3) Bank Texture
  - a) Bedrock – very resistant to erosion
  - b) Boulder/Coble – boulders >10”/cobbles 2.5 to 10” Moderate resistance to erosion
  - c) Gravel – 0.08-2.5” Moderate to high erodibility when dominant component.
  - d) Sand – 0.004 – 0.08” High bank erodibility when dominant component
  - e) Silt – 0.00008 – 0.004” Non-cohesive with high to very high erodibility
  - f) Clay – < 0.00008” Cohesive clays are relatively resistant to erosion
  - g) Mix – Variety of particle sizes. i.e. glacial till
- 4) Water Level Fluctuations
  - a) High mark
  - b) Current gage height

## APPENDIX F EQUIPMENT MANUALS

**Clinometer Manual** (Forestry Suppliers 2006)

**Laser Rangefinder Manual** (Bushnell 2011)

**Spherical Densiometer** (Forestry Suppliers 2013)