



<i>Title:</i> TOS Standard Operating Procedure: Cactus Biomass and Handling		<i>Date:</i> 03/10/2022
<i>NEON Doc. #:</i> NEON.DOC.001715	<i>Author:</i> S. Hiebert	<i>Revision:</i> C

TOS STANDARD OPERATING PROCEDURE: CACTUS BIOMASS AND HANDLING

PREPARED BY	ORGANIZATION	DATE
Katie Jones	FSU	02/13/2018
Sienna Hiebert	FOPS	06/29/2016
Courtney Meier	FSU	06/29/2016

APPROVALS	ORGANIZATION	APPROVAL DATE
Kate Thibault	SCI	03/10/2022

RELEASED BY	ORGANIZATION	RELEASE DATE
Tanisha Waters	CM	03/10/2022

See configuration management system for approval history.



Change Record

REVISION	DATE	ECO #	DESCRIPTION OF CHANGE
A	08/04/2016	ECO-03914	Initial Release
B	03/30/2018	ECO-05527	<ul style="list-style-type: none">• Added reference to mobile applications and generalized app instructions throughout• Added tagStatus• Updated variable names to align with ingest workbook and mobile application• Added mapping option for D14
C	03/10/2022	ECO-06787	<ul style="list-style-type: none">• Revised logo• Fixed formatting



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

TABLE OF CONTENTS

1 DESCRIPTION..... 1

1.1 Purpose..... 1

1.2 Scope 1

1.3 Applies To..... 1

1.4 Acknowledgments..... 1

2 RELATED DOCUMENTS AND ACRONYMS..... 2

2.1 Applicable Documents 2

2.2 Reference Documents 2

2.3 Acronyms 2

2.4 Definitions 2

3 SAFETY..... 3

3.1 Laser Rangefinder Safety..... 3

3.2 Cactus Safety..... 3

4 PERSONNEL AND EQUIPMENT 4

4.1 Equipment 4

4.2 Training Requirements..... 7

4.3 Specialized Skills 7

4.4 Estimated Time..... 7

5 CONTINGENCIES AND NOTES 8

6 STANDARD OPERATING PROCEDURES 9

SOPA PREPARING FOR THE FIELD 9

SOPB BIOMASS/PRODUCTIVITY MEASUREMENTS.....10

7 REFERENCES25

APPENDIX A QUICK REFERENCES26

APPENDIX B REMINDERS.....27

APPENDIX C ESTIMATED DATES FOR ONSET AND CESSATION OF SAMPLING.....28



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

LIST OF TABLES AND FIGURES

Table 1. Equipment list – Preparing for sampling 4

Table 2. Equipment list – Structure measurements at one plot..... 4

Table 3. Cactus shape descriptions and datasheet codes.15

Table 4. Structure Measurements. Shape determines which measurements are required.18

Table 5. Identification qualifier codes (idQ) to designate unknown species or those species with uncertain identification in the field or after identification in the lab.19

Table 6. List of potential structural measurements made on cactus individuals; note that not all measurements are made on a given cactus shape.....19

Table 7. plantStatus value definitions. For damaged individuals (codes 4-7), if the source of the damage cannot be determined, record as Damaged – other, include remarks describing damage.24

Table 8. Site-specific sampling start and end dates for cacti biomass measurements. See the Vegetation Structure Protocol (RD[04]) for start and end dates if they are not included in this table.28

Figure 1. Illustration of a 20 m × 20 m base plot (left; larger destructive sampling portion of the plot not shown), a 40 m × 40 m base plot (right), and associated nested subplots used for measuring woody stem vegetation11

Figure 2. *Cylindropuntia fulgida* in the elipsoid shape category illustrating where to take measurements.21

Figure 3. *Echinocereus* sp. illustrating a multi-stem columnar cacti in which one measures the height (in this case, the mean height of all stems), meanBasalDiameter, and stemCount.....22

Figure 4. *Opuntia* sp. with new vegetative pads growing off of an old pad.22

Figure 5. Branch length measurement (measure along the stem) in *Carnegiea gigantea*23



<i>Title:</i> TOS Standard Operating Procedure: Cactus Biomass and Handling		<i>Date:</i> 03/10/2022
<i>NEON Doc. #:</i> NEON.DOC.001715	<i>Author:</i> S. Hiebert	<i>Revision:</i> C

1 DESCRIPTION

1.1 Purpose

This document outlines the NEON Standard Operating Procedure (SOP) for acquiring structural measurements from members of the family Cactaceae (cacti). Structural data collected from cactus individuals may then be used to estimate volume, and in some cases, standing biomass. This SOP complements standardized structural measurements for woody plants described in the Vegetation Structure Protocol (RD[04]), and provides techniques specific to cactus growth forms. This SOP is only required at sites where qualifying cacti are present in sampling plots, and should be implemented concurrently with vegetation structure measurements.

1.2 Scope

This document provides a change-controlled version of an Observatory procedure. Documentation of content changes (i.e. changes in particular tasks or safety practices) will occur via this change-controlled document, not through field manuals or training materials.

1.3 Applies To

The procedure described in this document is used in the Measurement of Vegetation Structure (RD[04]) protocol and relies on the SOP B: Mapping and Tagging datasheet and the Biomass and Productivity Measurements- Other datasheet (RD[05]).

1.4 Acknowledgments



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

2 RELATED DOCUMENTS AND ACRONYMS

2.1 Applicable Documents

Applicable documents contain higher-level information that is implemented in the current document. Examples include designs, plans, or standards.

AD[01]	NEON.DOC.004300	EHS Safety Policy and Program Manual
AD[02]	NEON.DOC.004316	Operations Field Safety and Security Plan
AD[03]	NEON.DOC.050005	Field Operations Job Instruction Training Plan

2.2 Reference Documents

Reference documents contain information that supports or complements the current document. Examples include related protocols, datasheets, or general-information references.

RD[01]	NEON.DOC.000008	NEON Acronym List
RD[02]	NEON.DOC.000243	NEON Glossary of Terms
RD[03]	NEON.DOC.005003	NEON Scientific Data Products Catalog
RD[04]	NEON.DOC.000987	TOS Protocol and Procedure: Measurement of Vegetation Structure
RD[05]	NEON.DOC.001573	Datasheets for TOS Protocol and Procedure: Measurement of Vegetation Structure
RD[06]	NEON.DOC.001025	TOS Protocol and Procedure: Plot Establishment
RD[07]	NEON.DOC.001271	TOS Protocol and Procedure: Manual Data Transcription
RD[08]	NEON.DOC.001717	TOS Standard Operating Procedure: TruPulse Rangefinder Use and Calibration
RD[09]	NEON.DOC.001792	NEON Herbarium Specimen Label and Annotation Generation
RD[10]	NEON.DOC.014042	TOS Protocol and Procedure: Plant Diversity Sampling

2.3 Acronyms

All acronyms used in this document are defined in RD[01].

2.4 Definitions

Allometry: The study of the relationship between the relative size of plant structures and biomass.

Areole: An areola, especially a small area bearing spines or hairs on a cactus.

Cladode: A flattened organ of a plant; in *Opuntia* species, these are generally referred to as pads.

Glochid: A barbed bristle on the areole of cacti. Generally small and hair-like.

Pad: See cladode definition.



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

3 SAFETY

This document identifies procedure-specific safety hazards and associated safety requirements. It does not describe general safety practices or site-specific safety practices.

Personnel working at a NEON site must be compliant with safe field work practices as outlined in the Operations Field Safety and Security Plan (AD[02]) and EHS Safety Policy and Program Manual (AD[01]). Additional safety issues associated with this field procedure are outlined below. The Field Operations Manager and the Lead Field Technician have primary authority to stop work activities based on unsafe field conditions; however, all employees have the responsibility and right to stop their work in unsafe conditions.

3.1 Laser Rangefinder Safety

A laser rangefinder/hypsometer/compass instrument is used to map individual cacti (D14 only) and to measure various structural attributes (i.e., height, crown dimensions). Safety considerations for this instrument include:

- Avoid staring directly at the laser beam for prolonged periods. The rangefinder is classified as eye-safe to Class 1 limits, which means that virtually no hazard is associated with directly viewing the laser output under normal conditions. As with any laser device, however, reasonable precautions should be taken in its operation. It is recommended that you avoid staring into the transmit aperture while firing the laser.
- Never attempt to view the sun through the scope. Looking at the sun through the scope may permanently damage the eyes.

3.2 Cactus Safety

The potential for injury exists when working with cacti due to the prevalence of spines. As a simple precautionary measure, avoid making contact with cacti. Always wear personal protective equipment such as leather boots, long pants, long sleeves, and gloves, and remain cognizant of where you walk.

Most cactus-inflicted wounds should be treated like any other scratch or puncture, that is, cleaned then bandaged if necessary. Use a comb, tweezers, or adhesive to pick out spines that break off in the skin prior to treating the affected area.

Glochids that penetrate the skin may be difficult to extract because of their barbed shafts. To remove glochids, use tweezers or adhesives. Treat any remaining glochids as a wound, keep the area clean and covered to prevent dermatitis.

Segments of *Cylindropuntia* species break off easily and may become attached to footwear, clothing, or skin. To remove segments, use a comb or other solid object (i.e. Leatherman, rocks, etc.) to dislodge the segment. Symbiotic bacteria living on cactus spines may cause inflammation, which typically subsides without treatment within a few days.



4 PERSONNEL AND EQUIPMENT

4.1 Equipment

The following equipment is needed to implement the procedures in this document. Equipment lists are organized by task. They do not include standard field and laboratory supplies such as charging stations, first aid kits, drying ovens, ultra-low refrigerators, etc.

Table 1. Equipment list – Preparing for sampling.

Supplier	Supplier Number	R/S	Description	Purpose	Quantity	Special Handling
Durable Items						
Amazon Cabela's REI	IK270217 895022	S	GPS receiver, recreational accuracy	Navigate to sampling location at sites with plot markers	1	N
		S	USB Cable	Transfer data to GPS unit	1	N
Consumable Items						
		S	Rite in the Rain paper	Printing field datasheets	1	N

R/S=Required/Suggested

Table 2. Equipment list – Structure measurements at one plot.

Supplier	Supplier Number	R/S	Description	Purpose	Quantity	Special Handling
Durable Items						
Compass Tools	88180-04	S	GPS receiver, decimeter accuracy (e.g. GEO XH 6000, 7X)	Navigate to sampling locations at sites where markers are absent	1	N



Supplier	Supplier Number	R/S	Description	Purpose	Quantity	Special Handling
Amazon Cabela's REI	IK270217 895022	S	GPS receiver, recreational accuracy	Navigate to sampling location at sites with plot markers	1	N
		R	Backup mobile data recorder battery, fully charged	Continue data collect in the event tablet loses charge	1	N
		S	Regional flora reference guide and/or key	Identify unknown species	1	N
ForestrySuppliers	91567	S	Laser Rangefinder, ± 30 cm accuracy	Measure height	1	N
ForestrySuppliers	71112	R	Measuring stick, 2 m folding	Measure heights and widths of cacti	1	N
Ben Meadows ForestrySuppliers	122732 39945	R	Measuring tape, minimum 50 m	Delineate plot and subplot boundary	3	N
Ben Meadows ForestrySuppliers	100952 39167	R	Chaining pins or other suitable anchor	Anchor measuring tapes to delineate plot and subplot boundary	9	N
Ben Meadows ForestrySuppliers	213379 37184	R	Compass with mirror and declination adjustment	Determine plot boundary	1	N
Consumable Items						
RD[05]		R	Datasheets	Recording Data	1	N
		R	AA battery	Spare battery for GPS receiver	4	N



Supplier	Supplier Number	R/S	Description	Purpose	Quantity	Special Handling
		R	CR123A battery	Spare battery for laser rangefinder	2	N
		S	Survey marking flag, PVC or fiberglass stake	Delineate sampling area and temporarily mark stems before/after measurement	50	N
Ben Meadows Forestry Supplier	Varies depending on numerical sequence	R	Round numbered aluminum tag, silver; 0001-6000 and 8001-9999	Tag new qualifying stems	As needed	N
Paragon Spring	Pig Tail Stakes	R	Pigtail stake	Mark each specimen	As needed	N

R/S=Required/Suggested



<i>Title:</i> TOS Standard Operating Procedure: Cactus Biomass and Handling		<i>Date:</i> 03/10/2022
<i>NEON Doc. #:</i> NEON.DOC.001715	<i>Author:</i> S. Hiebert	<i>Revision:</i> C

4.2 Training Requirements

All technicians must complete required safety training and procedure-specific training for safety and implementation of this procedure as required in Field Operations Job Instruction Training Plan (AD[04]).

Technicians must be trained in the proper care of the laser rangefinder. Although this tool is resistant to dust and water, it is important to seal open ports and use lens caps when applicable. Care must also be taken to avoid scratching lenses. Finally, technicians should be trained to carefully measure the heights of cacti using the laser rangefinder.

4.3 Specialized Skills

At least one of the technicians executing this protocol must be able to identify regionally-specific cacti to species via visual inspection and use of a dichotomous key.

4.4 Estimated Time

The amount of time required to implement this protocol will vary depending on a number of factors, such as cactus density, skill level, species diversity, environmental conditions, and distance between plots. For a skilled two-person team, it should require no more than 4 hours to complete a single plot. If a task is taking significantly longer than the estimated time, a problem ticket should be submitted.



<i>Title:</i> TOS Standard Operating Procedure: Cactus Biomass and Handling		<i>Date:</i> 03/10/2022
<i>NEON Doc. #:</i> NEON.DOC.001715	<i>Author:</i> S. Hiebert	<i>Revision:</i> C

5 CONTINGENCIES AND NOTES

N/A



6 STANDARD OPERATING PROCEDURES

SOP A Preparing for the field

Measurements to estimate cactus biomass will be made on an individual plant basis, and the techniques used are consistent with those described in the Vegetation Structure protocol [RD(04)].

A.1 Configure GPS

Transfer all required files containing plot locations to the GPS receiver.

A.2 Check the laser rangefinder

1. Make sure the lenses on the laser rangefinder are free of dirt and debris. Clean with a lens cloth or lens tissue if necessary.
2. Declination is different at each site, and also changes with time. Due to these factors declination should be set when moving between sites, and per site values should be looked up annually at <http://www.ngdc.noaa.gov/geomag-web/>
3. *Declination Offset.* Check the current declination against what is entered in the laser rangefinder unit. See RD[08] for details.
4. *Tilt-sensor Calibration.* In the rare instance that the laser rangefinder has suffered a severe drop shock, the tilt-sensor requires re-calibration prior to continued field work. See RD[08] for details. To accurately perform tilt-sensor calibration, a surface known to be perfectly flat is required.
5. *Compass Calibration.* The compass should be calibrated after the batteries are changed. Be aware that interference from local magnetic fields may prevent accurate calibration, and can cause the calibration routine to fail.

A.3 Print field datasheets

Print Datasheets for TOS Protocol and Procedure: Measurement of Vegetation Structure RD[05]. The preferred method of data collection through use of a mobile device running the '(TOS) VST: Apparent Individuals [PROD]' mobile application, but datasheets should always be available in the field as a backup data collection method should the mobile device fail.

A.4 Sync Mobile Device

Check that mobile devices have the required applications loaded, are fully charged and synced before leaving the Domain Support Facility.



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

SOP B Biomass/Productivity Measurements

B.1 Introduction

This SOP is implemented at all Tower Plots and a subset of Distributed Plots, and in a manner consistent with the temporal and spatial sampling strategy outlined in TOS Protocol and Procedure: Measurement of Vegetation Structure (RD[04]).

B.2 Temporal Strategy

Cactus will typically be sampled concurrently with all other vegetation structure measurements, that is, during the dormant season after annual growth is complete. Tower plots are re-measured on either an annual or once every 3 year interval. Distributed plots are re-measured once every 3 years. Sampling should be scheduled to begin within ± 2 weeks of the date specified in Appendix H of the Vegetation Structure protocol. Tower plot sampling interval is annual at typical, temperate sites and once every 3 years at cold or dry continental sites. Appendix C summarizes the sampling window for sites where cactus are expected.

B.3 Spatial Strategy

For 20 m x 20 m Tower Plots, the entire 20 m x 20 m area is sampled. At sites with 40 m x 40 m Tower Plots, only two randomly selected 20 m x 20 m subplots are sampled (see **Figure 1**). Lists of randomly selected subplots are provided by Science Operations. In Distributed Plots, Cactaceae are sampled only in the inner 20 m x 20 m 'core' of the plot. To standardize sampling effort across plots, use nested subplots to constrain the measurement area, following the guidelines outlined in the Vegetation Structure protocol (see RD[04] for details). Briefly:

- In 20 m x 20 m plots: If nested subplots are employed, a minimum of 20 individuals, tallied across all nested subplots, must be measured in the entire 400 m² plot.
- In 40 m x 40 m plots: If nested subplots are employed, a minimum of 20 individuals must be measured in *at least one* of the assigned 20 m x 20 m subplots.

All cacti in the designated measurement area are identified and measured. As few established allometries are available to relate non-destructive field measurements to biomass of an individual, most cactus measurements will be used to calculate volume and will be treated in a similar fashion to shrub volume calculations described in RD[04]. In addition, structural measurements from cacti that NEON staff collects will provide a platform for potential community development of allometric biomass equations.




Figure 1. Illustration of a 20 m × 20 m base plot (left; larger destructive sampling portion of the plot not shown), a 40 m × 40 m base plot (right), and associated nested subplots used for measuring woodystem vegetation. The 20 m × 20 m plot size may be used for either Distributed or Tower Plots, and the 40 m × 40 m plot size is only for Tower Plots (at specific sites). Numbers in plain grey text indicate subplotIDs and numbers in italic black text indicate nested subplotIDs.

B.4 Qualifying Individuals

In order to qualify for measurement using this protocol, each specimen must be:

- In the Cactaceae family
- Alive **or** have a primary stem that is:
 - Woody and upright (with an angle that is greater than 45 degrees from the ground), **or**
 - Woody and with a naturally decumbent growth form
- Greater than or equal to 0.1 m stem length
 - Stem length minimum measurement applies to both upright and prostrate growth habits.
 - A minimum size is specified to constrain level of effort (LOE) for sampling activities. Cacti < 0.1 m stem length will be accounted for in plant diversity sampling.

B.5 Preparing for Data Capture

Mobile applications are the preferred mechanism for data entry.

Data recording procedures are more fully described in the Vegetation Structure protocol and in the Vegetation structure Fulcrum Manual. This SOP covers information that will be recorded in three different mobile applications as part of the comprehensive Vegetation Structure sampling bout:



- (TOS) VST: Plot Meta-Data [PROD]: presence/absence by growth form, nested subplot sizes, random subplot selection
- (TOS) VST: Mapping and Tagging [PROD]: tagID, taxonID and location information (mapping cacti only occurs in D14)
- (TOS) VST: Apparent Individuals [PROD]: field measurement of qualifying individual

Before going to the field:

- ***Double check that mobile devices are fully charged and synced***

Given the potential for mobile devices to fail under field conditions, it is imperative that paper datasheets are always available to record data. Paper datasheets should be carried along with the mobile devices to sampling locations at all times.

B.6 Procedure

1. Navigate to the selected plot.
2. Delineate the plot. Use existing plot markers, the 50 m tapes, and chaining pins to carefully delineate the plot and subplots.
 - It is not necessary to pay attention to whether the plot is sloped or flat; the tape is used only to help determine the plot edges and orient 1 m² plot frames.
 - Refer to the Plot Establishment Protocol (RD[06]) for a review of tape wrapping techniques that can be used to delineate subplots and nested subplots.
3. Plot metadata: Determine nested subplot size. Group all forms / species of cacti together for determining nested subplot sizes. Consult the Vegetation Structure protocol (RD[04]) for a thorough explanation of requirements for nested subplot use. Record value in the (TOS) VST: Plot Meta-Data [PROD] mobile application or on the plot metadata datasheet.
 - If executing this protocol in a 20 m x 20 m plot, a sum of, at minimum, 20 cacti are required across the entire plot if nested subplot are used.
 - If executing this protocol in a 40 m x 40 m plot, then a minimum of 20 cacti are required in at least one of the two randomly selected subplots.
4. For each plot, record in the Plot Metadata application:
 - **plotID**: SITE_###
 - **date**: YYYYMMDD
 - **nestedSubplotAreaOther**: This is the nested subplot area as defined in SOP C in RD[04], ### m²
 - **totalAreaSampledOther**: This is the subplot or nestedSubplotArea multiplied by the number of subplots surveyed. Auto-generated in mobile application.



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

- **Ex.** If plotSize = 40m x 40m and no nestedSubplotArea is used, all of the two randomly selected subplot are surveyed, totalAreaSampled = 800
- **measuredBy:** Name of the technician making measurements, email address
- **recordedBy:** Name of the technician recording data, email address
- Presence/absence of all growthForms,
- List of subplot/nested subplots where growthForm is absent (all growth forms listed as present within the plot). This provides a record of verified zeroes for end users.

5. Mapping and Tagging new individuals: Tag all qualifying cacti encountered within the selected measurement area except specimens in the `pads` shape category (see **Table 3**). Domain 14 will tag all qualifying cacti, including `pad` shape category.

- Tag with a unique aluminum ID tag for repeat measurements
- Attach tag to a pigtail stake
- Place the pigtail stake in the ground within 5 cm of a cactus stem. For the ease of finding the tag at a later time, place pigtails on the same side of the cactus for every specimen
- Be consistent with tag placement for cacti. It may be useful to tag consistent with vegetation structure methods of tagging for your site.



Example: D14 staff place the pigtail stake on the south side of cacti to avoid staring into the sun when locating a tag. Also be consistent in the spatial method you use for tagging; for example, begin in the Southwest corner of the plot and add tags as you move Northward within the plot.

- Record new tagIDs in the `(TOS) VST: Mapping and Tagging [PROD]` or SOP B: Mapping and Tagging datasheet
- Cacti in D14 may, optionally, be mapped according to guidelines in RD[05]. All other domains will not map tagged individuals.

6. Biomass measurements: For each qualifying cactus encountered, record:

- a. **subplotID:** number of the subplot in which the plant is located, ##, (see **Figure 1**)
- b. **nestedSubplotID:** unique identifier of the nested subplot - assigned on a per-subplot basis, #, (see **Figure 1**)
- c. **tagID:** domain level unique identifier, #####, null for untagged individuals.
- d. **tagStatus;** description of conditions of the tag for a previously measured individual. Choose one of the following options:
 - **ok;** tag is attached, and consistent with previously entered value
 - **replaced;** tag no longer attached, and has been re-made with same number
 - **mislabelled;** tagID was incorrectly recorded, and has been re-assigned






Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

- **untagged**; previously measured individual that no longer has a unique tagID assigned; cannot determine previous tagID. Assign new tagID.
 - **untagged**; no tag attached, not required for the selected individual
- e. **taxonID** and **identificationQualifier** (Table 5) code if needed: USDA plant species code, e.g., *OPPO* (*Opuntia polyacantha*), *CYFU10* (*Cylindropuntia fulgida*), etc. If the individual had been tagged and identified during a previous bout, this field will prepopulate in the mobile application.
- The **taxonID** should be a code from the NEON master list of plant species codes (found in the 'taxonTables' supplied by NEON Science Operations, and qualified according to technician confidence (Table 5).
 - A searchable list of all taxonID for the selected domain is available in the mobile applications. If a datasheet is being used, technicians are strongly encouraged to use ONLY the NEON master code. In the event that a code different from the NEON master code is used on a datasheet, the full **scientificName** associated with that code must be provided with each datasheet on which the non-NEON code is used, either via annotation or by attaching a key to the datasheet.
 - The NEON master taxon lists provide codes for instances when identification below a given taxonomic rank (e.g., family or genus) cannot be made. These are indicated by a 'sp.' or 'spp.' in the scientific name, where the former is used when only one unknown species is involved, and the latter when a shrub group may contain multiple species from a given genus. When one of these genus-level codes is selected, an identification qualifier is not needed, unless for example, the genus is uncertain.
 - Example: If you record **taxonID** = "CYLIN2" (*Cylindropuntia sp.*), do NOT record **idQ** = "cf. species"; it is already clear that you do not know the species based on the fact that a Genus-level code was reported in **taxonID**.
 - Morphologically challenging species:
 - It is expected that, during the course of this work, crews will encounter species that cannot confidently be identified in the field. In these cases, record morphospeciesID according to the guidelines in the Plant Diversity protocol (RD[10]).
- f. **identificationReferences**: If a taxonomic key is used to identify the individual, record, last name(s) of author(s)
- If no references are used for identification, leave this field blank
- g. **shape**: the form that the cactus takes that determines the required measurements.
- To determine the appropriate shape, consult **Table 3**, then match selected shape and required fields in **Table 4**.







- Make a determination of shape based on the individual being measured; shape is not necessarily consistent within a particular species.
- Shape is approximate, and is used to estimate volume. It is possible that multiple shapes may be appropriate for a given individual. Be consistent and do not spend more than 30 seconds assessing shape.



Table 3. Cactus shape descriptions and datasheet codes.

shape	Code*	Definition	Example shapes [†]	Example Genera
Columnar	col	Non-segmented cacti that are slender in shape and erect in growth habit, may have 1 or more stems.		<i>Stenocereus sp.</i> <i>Lophocereus sp.</i> <i>Echinocereus sp.</i> <i>Echinocactus sp.</i> <i>Mammillaria sp.</i>
Columnar taxonID = CAG110	col	Species-specific measurements for Saguaro.		<i>Carnegiea gigantea</i>
Oblate Half Sphere	ohs	Single stems depressed globose in shape OR cluster of segmented stems that, as a group, form an oblate half sphere.		<i>Astrophytum sp.</i> <i>Escobaria sp.</i>



shape	Code*	Definition	Example shapes [†]	Example Genera
Vine	vin	Cacti with vine-like, non-self supporting growth forms.		<i>Peniocereus sp.</i> <i>Hylocereus sp.</i>
Sphere or ellipsoid	eph	Single stems spherical in shape OR cluster of segmented stems elevated on central stalk		<i>Cylindropuntia sp.</i> <i>Ferocactus sp.</i> <i>Mammillaria sp.</i>
Half Sphere	hsp	Groups of clustered stems from a single individual that, as a group, approximately form a half sphere.		<i>Cylindropuntia sp.</i>
Inverted Cone	icn	Cluster of segmented stems that, as a group, approximately form an inverted cone shape.		<i>Cylindropuntia sp.</i>



shape	Code*	Definition	Example shapes [†]	Example Genera
Cone	cne	Cluster of segmented stems that, as a group, approximately form a cone shape.		<i>Cylindropuntia sp.</i>
Cylinder	cyl	Cacti that grow in long cylindrical segments and approximate the form of a cylinder shape.		<i>Cylindropuntia sp.</i>
Pad	pad	Cacti that have flat, round cladodes (<i>Opuntia</i>).		<i>Opuntia sp.</i>

**Cylindropuntia* and other segmented cacti may occur in a variety of shapes, assess each cacti individually consistent with shapes used to calculate volume of shrubs (half-sphere, cone, cylinder...)

[†] image credits: col, eph, icn, pad, and vin modified from Dimmitt 2000, ohs Molina-Freaner et al. 1998, cne, hsp, cyl modified from Engelmann and Bigelow 1856.



Table 4. Structure Measurements. Shape determines which measurements are required.

Shape	plantStatus	height [‡]	maxCrown Diameter	ninetyCrown Diameter	baseCrown Height	basalStem Diameter	measurement Height	maxBaseCrown Diameter	ninetyBase CrownDiameter	newPadCount	oldPadCount	stemCount	branchCount	meanBranch Length	meanBasal Diameter
Columnar	X	X										X			X
Columnar Saguaro [§]	X	X				X							X	X	
Oblate half sphere	X	X	X	X											
Vine	X												X	X	
Sphere or ellipsoid	X	X	X	X	X	X	X								
Half sphere	X	X	X	X											
Inverted cone	X	X	X	X				X	X						
Cone	X	X	X	X											
Cylinder	X	X	X	X											
Pad (<i>Opuntia</i>)	X									X	X				

[‡]For multi-stem columnar species, this is the mean stem length of the cluster

[§] Required measurements are based on (Steenbergh 1972, Niklas 2002)



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

Table 5. Identification qualifier codes (idQ) to designate unknown species or those species with uncertain identification in the field or after identification in the lab.

idqCode	identificationQualifier	Description
CS	cf. species	Roughly equals but “not sure” about the species
AS	aff. species	“Similar to, but is not” the species
CG	cf. genus	Roughly equals but “not sure” about the genus
AG	aff. genus	“Similar to, but is not” the genus
CB	cf. subspecies	Roughly equals but “not sure” about the subspecies
AB	aff. subspecies	“Similar to, but is not” the subspecies
CF	cf. family	Roughly equals but “not sure” about the family
AF	aff. family	“Similar to, but is not” the family
CV	cf. variety	Roughly equals but “not sure” about the variety
AV	aff. variety	“Similar to, but is not” the variety

- After **shape** has been assigned, determine which of the measurements listed in **Table 6** must be recorded for the current individual:

Table 6. List of potential structural measurements made on cactus individuals; note that not all measurements are made on a given cactus shape.

Variable	Detailed description
Height*	Maximum height of single stem individuals, mean height for multi-stem individuals, measured using folding ruler, laser rangefinder, or measuring tape, 0.1 m, (see Figure 5)
maxCrownDiameter*	Largest diameter of canopy or stem, depending on which is larger, measured using calipers or folding ruler, 0.1 m, (see Figure 2)
ninetyCrownDiameter*	Diameter at 90° to maxDiameter measured using calipers or folding ruler, 0.1 m, (see Figure 2)
baseCrownHeight*	Use a rigid, collapsible meter stick to measure the average height above the ground for the lowest portion of the crown, 0.1 m
basalStemDiameter	Cross-sectional stem diameter at soil surface, measured using calipers or folding ruler, 0.01m
measurementHeight	The height at which the diameter is measured, soil surface = 0 cm, 1 cm
maxBaseCrownDiameter	Maximum diameter of the base of individuals in the inverted cone shape category, measuring tape, folding ruler or laser rangefinder, 1 cm, (see Figure 2)
baseCrownNinetyDiameter	90 degrees measurement of the maxBasalDiameter for individuals in the inverted cone shape category, 1 cm, (see Figure 2)
newPadCount	Number of cladodes, of any size, produced in the current growing season (see Figure 4) Count vegetative, non-flowering, pads only.
oldPadCount	Number of cladodes produced in previous growing seasons (see Figure 4)
stemCount	Number of stems in a multi-stem columnar cluster (see Figure 3)



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

Variable	Detailed description
meanBranchLength	For cacti in the columnar or vine shape that also have branches. To account for biomass in the branches in columnar or vine cacti (i.e. <i>Carnegiea gigantea</i> , <i>Peniocereus sp.</i> , etc.) measure the length of each branch using a folding ruler, laser rangefinder, or measuring tape. Measure along the length of branches, including bends if present. If there is more than one branch length, calculate the mean of all branch lengths, and report to the nearest 0.1 m; see Figure 5 for examples of how to measure complicated branch lengths
meanBasalDiameter	Use calipers to measure the basal diameter of each stem at ground level. Calculate the mean of all stems to the nearest 0.1 cm. For densely clustered multi-stem columnar individuals where basal diameter of interior stems cannot be measured directly, estimate basal diameter based on diameter of visible portion of the stem, assume columnar shape.
branchCount	For cacti in the columnar or vine shape that also have branch(es), the number of branches on individual cactus, (see Figure 5)
plantStatus	Health status of individual, if known, note the factors associated with death or damage in the notes section of the datasheet along with the corresponding tagID, #, (see Table 7)

* small individuals may be measured to cm accuracy (0.01 m)




Figure 2. *Cylindropuntia fulgida* in the elipsoid shape category illustrating where to take measurements. Measure basalStemDiameter (yellow line) at the ground level; maxCrownDiameter (black line) and ninetyDiameter across the canopy (black dotted circle); baseCrownHeight (green line) from the ground to the base of the crown and total height (purple line). Photo by Sienna Hiebert.




Figure 3. Echinocereus sp. illustrating a multi-stem columnar cacti in which one measures the height (in this case, the mean height of all stems), meanBasalDiameter, and stemCount. Photo by Sienna Hiebert.



Figure 4. *Opuntia* sp. with new vegetative pads growing off of an old pad. In this picture newPadCount = 3 and oldPadCount = 1.




Figure 5. Branch length measurement (measure along the stem) in *Carnegieia gigantea*. Measure each branch then average the length of all branches. Enter that value in the meanBranchLength data field. For cacti that have become cristate like in this picture, note “crested” in the **remarks** field. Photo by Sienna Hiebert.



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

Table 7. plantStatus value definitions. For damaged individuals (codes 4-7), if the source of the damage cannot be determined, record as Damaged – other, include remarks describing damage.

Code	Description
1	Live– any live cacti (new, re-measured or ingrowth).
2	Standing dead – either new or re-measured, regardless of factors associated with death. The entire cacti must be dead.
3	Removed – a cactus that has been cut and removed by direct activity to harvesting or land clearing (re-measurement plots only).
4	Live, insect damaged - Visible damage caused by insects; note 'crown' or 'bole' damage in remarks, and indicate type of insect causing damage if possible
5	Live, disease damaged - Visible damage cause by disease; indicate location and type of disease, in remarks, if possible
6	Live, physically damaged – Visible damage not caused by disease or insects; indicate location and type of physical damage, in remarks, if possible
7	Live, other damage – Note location and type of damage, in remarks, if possible
8	No longer qualifies – Note reason in remarks
11	Lost, burned - A previously measured individual that is not measured in the current bout because the plot has been burned and the individual could not be located for remeasurement
12	Lost, herbivory - A previously measured individual that is not measured in the current bout because the individual could not be located for remeasurement. Cause of loss is presumed to be herbivory
13	Lost, presumed dead - A previously measured individual that is not measured in the current bout because the individual could not be located for remeasurement, the individual is presumed dead based on evidence within the plot. Note in remarks reason (i.e. blowdown event)
14	Lost, tag damaged - A previously measured individual that is not measured in the current bout due to inability to re-locate because of widespread damage to tags within a plot
15	Lost, fate unknown - A previously measured individual that is not measured in the current bout note because the individual could not be relocated
16	Downed - Part or all of previously measured individual is down, and in contact with the ground.



7 REFERENCES

- Dimmitt, M. A. 2000. Cactaceae (cactus family). Page 628 *in* S. J. Phillips and Comus, editors. A Natural History of the Sonoran Desert. Arizona-Sonora Desert Museum. Arizona-Sonora Desert Museum Press, Tucson, Arizona, and University of California Press, Berkeley, California.
- Engelmann, G., and J. M. Bigelow. 1856. The Botany of the Expedition: Description of the Cactaceae. Pages 1–58. Explorations and surveys for a railroad route from the Mississippi River to the Pacific Ocean. Route near the twenty-fifth parallel, Explored by Lieutenant A. W. Whipple, Topographical Engineers, in 1853 and 1854., War Department, Washington, D. C.
- Molina-Freaner, F., C. Tinoco-Ojanguren, and K. Niklas. 1998. Stem biomechanics of three columnar cacti from the Sonoran Desert. *American Journal of Botany* 85:1082–1082.
- Niklas, K. J. 2002. On the allometry of biomass partitioning and light harvesting for plants with leafless stems. *Journal of theoretical biology* 217:47–52.
- Steenbergh, W. F. 1972. Lightning-caused destruction in a desert plant community. *The Southwestern Naturalist*:419–429.



APPENDIX A QUICK REFERENCES

Quick Reference: Making Quality Measurements of Cactaceae

Step 1 – Calibrate laser rangefinder compass.

Step 2 – Delineate measurement area.

Step 3 - Assess need for nested subplots (new plots only).

Step 4 – Identify qualifying species.

Step 5 - Record metadata.

Step 6 – Tag qualifying species.

Step 7 – Take and record biomass measurements.

Step 8 – Remove temporary flagging.

For directions on using the laser rangefinder, see TOS Standard Operating Procedure: TruPulse Rangefinder Use and Calibration (RD[09]).



APPENDIX B REMINDERS

Making quality measurements of vegetation structure

Measurement Area: Make sure you know ...

- ✓ Size of plot and subplots.
- ✓ Number of subplots in the plot.
- ✓ Size of nested subplots (if any) for plots previously measured.
- ✓ How to determine whether nested subplots are needed for new plots.

Taking measurements: Remember to...

- ✓ Carefully record all metadata, measurements, and observations in mobile application or on datasheet.

Using the laser rangefinder: Pay close attention to ...

- ✓ Declination – Is it set for your current location?
- ✓ Battery charge – Replace when low-charge indicated and bring back-up batteries.
- ✓ Transcription of measurements onto datasheet.
- ✓ Metal objects – Keep them at least 2 feet away from instrument when using internal compass.



Title: TOS Standard Operating Procedure: Cactus Biomass and Handling		Date: 03/10/2022
NEON Doc. #: NEON.DOC.001715	Author: S. Hiebert	Revision: C

APPENDIX C ESTIMATED DATES FOR ONSET AND CESSATION OF SAMPLING

The dates in **Table 8** below are based on historic records and estimated from MODIS-EVI phenology data averaged from 2001-2009, and are therefore estimates for the average start and stop dates of sampling. It is essential that domain staff monitor real-time conditions to determine when to start (and stop) sampling. Unless indicated otherwise, “End Date” values are in the next calendar year. The sites included in **Table 8** are only sites where cacti are expected to be. If cacti exist at a site that is not included on **Table 8**, see the Vegetation Structure Protocol (RD[04]) for start and end dates, and issue a problem ticket.

“Start Date” definition: Below, values in the “Start Date” field correspond to the average day of year at which greenness begins to decrease.

“Start Date” and “End Date” fields are relevant to cacti measurement windows in both Distributed and Tower Plots, and represent the period of time during which vegetation photosynthetic activity is minimal following a growing season. If you feel this assessment is inaccurate for your site, please submit a problem ticket to Science Operations. If provided measurement windows are not logistically feasible, changes to “Start Date” may be made in consultation with Science Operations.

Table 8. Site-specific sampling start and end dates for cacti biomass measurements. See the Vegetation Structure Protocol (RD[04]) for start and end dates if they are not included in this table.

Domain Number	Site ID	Start Date	End Date	Tower plot sampling interval	Additional Sampling Information
03	DSNY	07/09	03/01	Annual	Target taxa present.
	JERC	08/08	03/31	Annual	Target taxa present.
	OSBS	07/09	03/11	Annual	Target taxa present.
04	GUAN	12/01	03/01	Annual	Dates correspond to the dry season, and are derived from Ensenada precipitation data (1980-2015).
10	CPER	07/29	03/31	Annual	Target taxa present.
	RMNP	07/29	04/30	Annual	
11	CLBJ	06/29	03/01	Annual	Target taxa present.
	OAES	NA	NA	Annual	
13	MOAB	08/13	03/26	3-year	Target taxa present.
14	JORN	09/02	03/21	3-year	Target taxa present.
	SRER	08/28	05/31	3-year	Target taxa present.
15	ONAQ	06/19	03/16	3-year	Target taxa present.