

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

TOS SCIENCE DESIGN FOR MOSQUITO ABUNDANCE, DIVERSITY, AND PHENOLOGY

PREPARED BY	ORGANIZATION	DATE
Katherine LeVan	SCI	04/20/2018
David Hoekman	SCI	09/10/2014
Cara Gibson	SCI	09/07/2011

APPROVALS	ORGANIZATION	APPROVAL DATE
Kate Thibault	SCI	06/14/2018
Mike Stewart	SYE	06/14/2018

RELEASED BY	ORGANIZATION	RELEASE DATE
Judy Salazar	CM	06/14/2018

See configuration management system for approval history.

The National Ecological Observatory Network is a project solely funded by the National Science Foundation and managed under cooperative agreement by Battelle. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

Change Record

REVISION	DATE	ECO #	DESCRIPTION OF CHANGE
A	10/10/2014	ECO-02020	Initial release
B	06/14/2018	ECO-05584	<ul style="list-style-type: none"> • Minor format and text edits • Corrected protocol names in reference documents and updated NEON TOS site numbers • Updated DNA barcoding section to include project names, site-specific deviations from the design

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

TABLE OF CONTENTS

1 DESCRIPTION.....1

1.1 Purpose 1

1.2 Scope..... 1

1.3 Acknowledgments..... 1

2 RELATED DOCUMENTS AND ACRONYMS1

2.1 Applicable Documents 1

2.2 Reference Documents..... 2

2.3 External References 2

2.4 Acronyms 2

3 INTRODUCTION3

3.1 Overview of the Observatory..... 3

3.2 Components of the Observatory 4

3.3 The Terrestrial Observation System (TOS)..... 4

4 INTRODUCTION TO THE MOSQUITO ABUNDANCE, DIVERSITY, AND PHENOLOGY SAMPLING DESIGN5

4.1 Background 5

4.1.1 Mosquitoes as a Sentinel Taxon 5

4.2 NEON’s Contribution..... 6

4.3 Purpose and Scope..... 6

5 SAMPLING FRAMEWORK7

5.1 Science Requirements..... 7

5.2 Data Products..... 7

5.3 Priorities and Challenges for Mosquito Abundance, Diversity, and Phenology Sampling 7

6 SAMPLING DESIGN FOR MOSQUITO ABUNDANCE, DIVERSITY, AND PHENOLOGY8

6.1 Sampling Methods 8

6.2 Sample Processing 10

6.2.1 Sample Handling 11

6.3 Taxonomic verifications and reference collections 11

6.3.1 Archive 12

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

6.4	Spatial Distribution of Sampling.....	12
6.4.1	Selecting Plot Locations within Sites.....	12
6.5	Temporal Distribution of Sampling.....	12
6.5.1	Sampling Session.....	12
6.5.2	Temperature Threshold	13
6.5.3	Field Season Sampling.....	15
6.5.4	Seasonal Timing	15
6.6	Logistics and Adaptability	17
7	REFERENCES	19

LIST OF TABLES AND FIGURES

Table 1.	Data summary of prototype field mosquito collection from Domain 3 in the summer of 2012..	10
Figure 1.	The seven Grand Challenges defined by the National Research Council (Committee on Grand Challenges in Environmental Sciences 2001).....	3
Figure 2.	Field season sample session timing of mosquito trap deployment	13
Figure 3.	Field season mosquito sampling schedule for a representative domain. Sampling typically occurs at one site each week, alternating between the core and relocatable sites. The number of weeks in the field season varies among domains.....	15
Figure 4.	Annual mosquito sampling timeline for a representative domain. Trapping occurs all year at the core site, with more traps during the warm part of the year when mosquitoes are most active. Note: from 2013 to 2016, a 4C threshold was used to determine the implementation of bouts; in 2017 and later, a revised threshold of 10C was used based on observed captures during the first four years of sampling.	16
Figure 5.	An example of mosquito sampling at a core site. The example begins with and ends with weekly off-season sampling and shows a brief field season. Sampling at relocatable sites is not shown but would occur in the intervening weeks (4-14).....	17

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

1 DESCRIPTION

1.1 Purpose

NEON design documents are required to define the scientific strategy leading to high-level protocols for NEON subsystem components, linking NEON Grand Challenges and science questions to specific measurements. Many NEON *in situ* measurements can be made in specific ways to enable continental-scale science rather than in ways that limit their use to more local or ecosystem-specific questions. NEON strives to make measurements in ways that enable continental-scale science to address the Grand Challenges. Design Documents flow from questions and goals defined in the NEON Science Strategy document, and inform the more detailed procedures described in Level 0 (L0; raw data) protocol and procedure documents, algorithm specifications, and Calibration/Validation (CalVal) and maintenance plans.

1.2 Scope

This document defines the rationale and requirements for mosquito abundance, diversity and phenology sampling in the NEON Science Design.

1.3 Acknowledgments

The design was reviewed by and refined with input from a technical working group consisting of researchers with relevant expertise. The mosquito technical working group includes Drs. Chris Barker, Roberto Barrera, Mark Blackmore, William Bradshaw, Desmond Foley, Howard Ginsberg, Mary Hayden, Chris Holzapfel, Steve Juliano, Laura Kramer, Shannon LaDeau, Chet Moore, Roger Nasci, Bill Reisen, and Harry Savage. Tanya Chesney also contributed to this document.

2 RELATED DOCUMENTS AND ACRONYMS

2.1 Applicable Documents

Applicable documents contain information that shall be applied in the current document. Examples are higher level requirements documents, standards, rules and regulations.

AD[01]	NEON.DOC.000001	NEON Observatory Design
AD[02]	NEON.DOC.000913	TOS Science Design for Spatial Sampling
AD[03]	NEON.DOC.000911	TOS Science Design for Vectors and Pathogens

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

2.2 Reference Documents

Reference documents contain information complementing, explaining, detailing, or otherwise supporting the information included in the current document.

RD[01]	NEON.DOC.000008	NEON Acronym List
RD[02]	NEON.DOC.000243	NEON Glossary of Terms
RD[03]	NEON.DOC.002652	NEON Level 1, Level 2 and Level 3 Data Products Catalog
RD[04]	NEON.DOC.014049	TOS Protocol and Procedure: Mosquito Sampling

2.3 External References

N/A

2.4 Acronyms

NOAA	National Oceanic and Atmospheric Administration
NCDC	National Climatic Data Center

All other acronyms used in this document are defined in RD[01].

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

3 INTRODUCTION

3.1 Overview of the Observatory

The National Ecological Observatory Network (NEON) is a continental-scale ecological observation platform for understanding and forecasting the impacts of climate change, land use change, and invasive species on ecology. NEON is designed to enable users, including scientists, planners and policy makers, educators, and the general public, to address the major areas in environmental sciences, known as the Grand Challenges (Fig. 1). NEON infrastructure and data products are strategically aimed at those aspects of the Grand Challenges for which a coordinated national program of standardized observations and experiments is particularly effective. The open access approach to the Observatory’s data and information products will enable users to explore NEON data in order to map, understand, and predict the effects of humans on the earth and understand and effectively address critical ecological questions and issues. Detailed information on the NEON design can be found in AD[01], AD[02].




Figure 1. The seven Grand Challenges defined by the National Research Council (Committee on Grand Challenges in Environmental Sciences 2001)

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

3.2 Components of the Observatory

There are five components of the Observatory, the Airborne Observation Platform (AOP), Terrestrial Instrument System (TIS), Aquatic Observation System (AOS), Aquatic Instrument System (AIS), and Terrestrial Observation System (TOS). Collocation of measurements associated with each of these components will allow for linkage and comparison of data products. For example, remote sensing data provided by the Airborne Observation Platform (AOP) will link diversity and productivity data collected on individual plants and stands by the Terrestrial Observation System (TOS) and flux data captured by instruments on the tower (TIS) to that of satellite-based remote sensing. For additional information on these systems, see Keller et al. (2008) and Schimel et al. (2011b).

3.3 The Terrestrial Observation System (TOS)

The NEON TOS will quantify the impacts of climate change, land use, and biological invasions on terrestrial populations and processes by sampling key groups of organisms (sentinel taxa), infectious disease, soil, and nutrient fluxes across system interfaces (air, land, and water) (AD[01], AD[02]). The sentinel taxa were selected to include organisms with varying life spans and generation times, and wide geographic distributions to allow for standardized comparisons across the continent. Many of the biological measurements will enable inference at regional and continental scales using statistical or process-based modeling approaches. The TOS sampling design captures heterogeneity representative of each site to facilitate this inference when possible. Plot and organism-scale measurements will also be coordinated with the larger-scale airborne measurements, which provide a set of synergistic biological data products at the regional scale. Details of these design elements and algorithms can be found in individual design documents available through the NEON website (www.neonscience.org).

The standardization of protocols across all sites is key to the success of NEON (and its novelty) and must be maintained at all sites through time. Thus, although specific techniques may be required at some sites (e.g., due to different vegetation types), protocols have been developed to ensure data comparability. These details can also be found in individual design documents available through the NEON website (www.neonscience.org).

The TOS Science Designs define the scientific strategies leading to high-level sampling designs for NEON sentinel taxa, terrestrial biogeochemistry, and infectious disease, linking NEON Grand Challenges and science questions to specific measurements. The TOS Spatial Sampling Design document describes the sampling design that collocates observations of the components of the TOS (AD[02]). TOS Science Design documents were developed following input from the scientific community, including module-specific Technical Working Groups, and the National Science Foundation (AD[01]). Science Designs will be reviewed periodically to ensure that the data collected by NEON are those best suited to meet the requirements of the observatory (AD[01]), are (to the extent possible) consistent with standards used by the scientific community, and fit within the scope of NEON.

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

4 INTRODUCTION TO THE MOSQUITO ABUNDANCE, DIVERSITY, AND PHENOLOGY SAMPLING DESIGN

4.1 Background

4.1.1 Mosquitoes as a Sentinel Taxon

The Terrestrial Observation System (TOS) at NEON is charged with monitoring the responses of biodiversity and ecosystems to environmental change. A NEON design committee (AIBSnews 2007) selected mosquitoes (Diptera: Culicidae) as a focal taxon for measurement. Mosquitoes are a diverse and widespread family of insects with aquatic larval and pupal forms and flying adults that have been extensively studied because of their ecological and epidemiological significance. As a dominant taxon in aquatic food webs, mosquitoes compose a sizable proportion of invertebrate biomass in aquatic systems and act as a key food source for aquatic and terrestrial predators (e.g., fish, amphibians, spiders, birds). Mosquitoes also act as vectors for numerous parasites and pathogens of humans, livestock, and wildlife and their biology and ecology have been extensively studied to characterize and mitigate impacts of associated diseases. Most female mosquitoes collect blood meals from vertebrates in order to provide energy for their developing eggs. Due to their potential impacts on human health, mosquito populations have been and continue to be monitored by national, state and local agencies. Mosquito-borne diseases can also influence the health of livestock [e.g., Rift Valley Fever (Daubney and Hudson 1931), Eastern equine encephalitis (Kissling et al. 1954)] and wildlife populations [e.g., avian malaria (Van Riper III et al. 1986), West Nile Virus (Marra et al. 2004)]. For example, the emergence of West Nile virus in North America has resulted in widespread population declines of several common birds (e.g., crows, robins, wrens, chickadees, blue jays; LaDeau et al. 2007) with important potential consequences for ecosystem services like seed dispersal, carrion scavenging and insect regulation (LaDeau et al. 2008).

Because of their sensitivity to environmental gradients and perturbations, mosquitoes represent an ideal sentinel taxon for evaluating the ecological effects of global change phenomena. The distribution of mosquito populations and seasonal phenology are influenced by many landscape factors including climate, vegetation and host availability (Buckner et al. 2010, Reisen 2010). Their short generation time and high fecundity allow mosquitoes to respond quickly to environmental change, but because of the group’s high diversity and varied ecological niches, the nature and magnitude of these changes can differ markedly among species.

Changes in global climate are predicted to affect the distribution, demography, and seasonal phenology of many mosquitoes; associated effects on disease transmission cycles have also been posited (Epstein et al. 1998). For example, as the climate warms mosquito populations are expanding their geographical ranges (Hongoh et al. 2011) and transmission rates may increase, though climate is not the only driver of mosquito-borne diseases dynamics (Reiter 2001). Because of their frequent association with humans and ability to thrive in human-modified environments, mosquito ecology is also likely to be significantly affected by land use changes. Mosquitoes are highly mobile and able to move into new areas as climatic

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

conditions change, often aided by unintentional human transport (Lounibos 2002). Invasions of mosquitoes that carry human diseases from tropical into sub-tropical and temperate areas are more likely under warmer conditions. Climate conditions influence not only mosquito distributions but also the life cycles of pathogens and the transmission of disease (Gage et al. 2008). Higher temperatures can shorten the life cycle of viruses and mosquitoes, increase blood meal and oviposition rates and thereby the efficiency of transmission (Reisen 2010). Together, these climate effects may expand the biogeographic ranges of mosquitoes and the diseases they carry into temperate areas. Climate may also indirectly influence mosquitoes and disease dynamics by affecting the phenology of their hosts (Burkett-Cadena et al. 2011). In addition, mosquitoes have tremendous genetic variability and an extensive body of genetic information is available for these insects. NEON will measure mosquito populations across a broad spatial extent for the next 3 decades and will be able to detect changes in their distribution and phenology.

4.2 NEON’s Contribution

Earth’s environment is changing rapidly. Data at the temporal and spatial scales that NEON will provide are necessary to understand, forecast and manage our changing biosphere (Keller et al. 2008, Schimel et al. 2011a).

NEON’s mosquito sampling will provide a cost effective and informative measure of a biological response to environmental, climate and land-use change. NEON sampling will augment state and local mosquito collection efforts and will enable us to characterize the response of mosquito populations, including abundance, phenology, and range to land use and other ecosystem drivers.

4.3 Purpose and Scope

This document defines the rationale and requirements for mosquito (Diptera: Culicidae) abundance, diversity and phenology in the NEON Science Design. Details about protocols including educational materials to conduct protocols, quality assurance and quality control or calibration and validation procedures are addressed in protocol specific documents.

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

5 SAMPLING FRAMEWORK

5.1 Science Requirements

This science design is based on Observatory science requirements that reside in NEON’s Dynamic Object-Oriented Requirements System (DOORS). Copies of approved science requirements have been exported from DOORS and are available in NEON’s document repository, or upon request.

5.2 Data Products

Execution of the protocols that stem from this science design procures samples and/or generates raw data satisfying NEON Observatory scientific requirements. These data and samples are used to create NEON data products, and are documented in the NEON Level 1, Level 2 and Level 3 Data Products Catalog (RD[03]).

5.3 Priorities and Challenges for Mosquito Abundance, Diversity, and Phenology Sampling

Standardized, well established and widely used sampling methods were selected to maximize comparability across time and between domains within NEON and be comparable to other mosquito collection programs. To enhance utility to the scientific community, Center for Disease Control and Prevention (CDC)-CO₂ light traps, standard mosquito collection tools, were selected.

NEON’s general strategy for the allocation of mosquito sampling effort is to sample core sites more frequently than relocatable sites. Core sites are prioritized because of the long-term nature of core site sampling (expected to continue for the entire 30-year lifespan on the observatory in contrast to relocatable sites, which are expected to be sampled for 7-10 years). Relocatable sites, which will be sampled less frequently than core sites, will serve to expand the spatial extent of inference.

Two primary constraints govern the spatial distribution of plots within a site. The first is collocation of measurements in order to promote comparability and the characterization of relationships between disparate TOS measurements across the observatory. Second, sampling logistics limit the amount of travel time that can be allocated to checking mosquito traps. Due to these dual constraints, this design places traps within 30m of roads while following the stratified-random sampling design used to spatially allocate sampling for other TOS protocols and co-locating the mosquito traps with distributed plots (where plant, beetle, and soil sampling may also occur; see AD[02] for more details) where possible.

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

6 SAMPLING DESIGN FOR MOSQUITO ABUNDANCE, DIVERSITY, AND PHENOLOGY

In keeping with NEON’s broad mission, this design must be able to be deployed across a wide range of sites in a standardized way (e.g., methods, sampling frequencies) such that data are comparable across time and space. The design must be relatively uncomplicated so that it can be deployed consistently by disparate field crews over multiple years with minimal chance of alteration.

NEON is intended to be an integrated measurement system. Coordinating measurements made for multiple TOS data products, e.g., mosquito abundance and diversity (the primary focus of this design) with pathogen presence and prevalence (AD[04]), is therefore a key component of the NEON design. A potential consequence of an integrated measurement design is sub-optimal sampling for individual data products at individual sites. Coordinating mosquito abundance and diversity sampling with sampling for mosquito-borne pathogens results in a number of attractive efficiencies, including saving considerable time and money because the same mosquito samples can be counted and identified as well as tested for pathogens. However, abundance and diversity sampling aims to survey a broad cross section of the mosquito community, while pathogen testing targets particular vector species and requires as many individuals as possible for testing. Thus, the two sampling efforts differ fundamentally in their foci and objectives and would be optimized using different sampling strategies, resulting in a trade-off between data quality at the site level and comparability utilizing a combined sampling design. The sampling approach adopted and described here is optimized for mosquito abundance and diversity sampling by using taxonomically general sampling spread broadly across time and space. Collected mosquitoes will subsequently be tested for pathogen presence/prevalence and additional trapping for specific vector mosquito species may be implemented at certain sites based on sampling efficacy (AD[04]). In this document, 'mosquito sampling' is used to describe the sampling for the coordinated sampling activities for abundance and diversity and pathogen sampling, unless otherwise specified.

6.1 Sampling Methods

Mosquitoes will be sampled at NEON sites using CDC CO₂ light traps, a standard and widely used method used by the CDC and other mosquito researchers for public health monitoring for mosquito-borne diseases for a half-century (Sudia and Chamberlain 1962). Although there are many variations on this basic trap, all traps use CO₂ as bait because CO₂ is a component of vertebrate breath that mosquitoes use to locate hosts. The sublimating CO₂ attracts mosquitoes to the vicinity of the trap and a fan that pulls the mosquitoes into a net bag for live storage until the trap is collected by a field technician. Some traps also include a light source, but light bulbs will be disconnected for NEON sampling because, though they do help attract some mosquitoes, light also attracts a wide variety of bycatch organisms that complicate sample processing and ultimately reduce sample quality.

CDC CO₂ light traps arguably collect the greatest diversity of all common species and are regularly used in mosquito-borne disease surveillance, thus maximizing comparability with other data sets (Sudia and Chamberlain 1962, Service 1993). However, they have known limitations and biases (e.g., do not

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

effectively sample *Aedes* spp. or blood-fed mosquitoes generally targeted for pathogen testing) and ideally should be paired with at least one additional trap type to ensure taxonomically broad sampling. Gravid traps are an attractive option, in part because they effectively sample blood-fed mosquitoes (Reiter 1983), but logistic challenges associated with standardization and transport of the fetid water into remote locations limit their usability within NEON. Resting box traps (Komar et al. 1995, 2011, Williams and Gingrich 2007, Burkett-Cadena et al. 2008) and BG-sentinel traps (Biogents; Krockel et al. 2006, Meeraus et al. 2008) are likely to capture mosquitoes especially useful for pathogen testing at some sites. Therefore resting box and BG-sentinel traps may be implemented along with CDC CO₂ light traps to augment coverage of important vector species (see the TOS Science Design for Vectors and Pathogens (AD[03]) for more details).

Mosquitoes exhibit diel activity patterns; some species are most active during crepuscular periods while others are most active during the night (Silver 2008). Traps are typically set in the late afternoon and allowed to run through the night until morning the following day. However, some mosquitoes are also day flyers (Hoel et al. 2009), and, in order to maximize coverage of mosquito activity, some daytime sampling will be conducted. NEON will trap during the night for comparison with other mosquito monitoring efforts and re-set the traps in the morning and trap during the day to sample the entire mosquito activity period and capture species that may be missed during evening sampling.

Mosquitoes exhibit strong seasonal abundance patterns. At many sites there may be months when mosquito activity is non-existent due to low temperatures. We will conduct spatially constrained low intensity sampling during the “off season” to monitor mosquito absence and will resume normal field season sampling upon the detection of mosquito presence (specific thresholds described in section 6.5.4). Off season samples will be used to define the “shoulders” of the annual mosquito season at each site.

From 2010-2013, NEON prototyped CDC CO₂ light traps at 16 sites with broad geographical coverage (including sites in Massachusetts, New Hampshire, Michigan, North Dakota, Utah, Colorado, Virginia, Tennessee, Alabama and Florida). The methods detailed above were implemented in summer 2012 at the three sites in Domain 03, Jones Ecological Research Center, Ordway-Swisher Biological Station and Disney Wilderness Preserve. During the D03 prototype, mosquito traps were deployed for a total of 22 nights. On average, over 600 mosquitos were captured per trap and therefore 10 traps per site yielded over 6000 mosquitoes for a trapping night. Overall, a total of 141,892 mosquitoes from 43 different species were collected (Table 1).

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

Table 1. Data summary of prototype field mosquito collection from Domain 3 in the summer of 2012

Site	Total mosquito abundance	Number of sampling sessions conducted	Total number of trap nights	Plots per site	Average number of mosquitoes captured per trap	Average total abundance for one night of trapping at a site	Number of mosquito species recorded
Jones	30944	4	8	10	380	3868	22
Ordway	64390	4	8	8-10	933	8049	26
Disney	46558	3	6	10	776	6651	31
Domain 3 total	141892	11	22		682	6169	43

Prototype sampling efforts have contributed to a DNA barcode library, site-specific species lists (with indications of nativity sourced from the literature), and a test dataset to use for optimizing NEON’s data ingest and quality control processes. In addition, logistical challenges associated with deploying traps in disparate environments served to inform the current sampling design. Specific issues and lessons learned include:

- Light bulbs, which are included as part of the CO₂ traps were observed to result in a higher amount of bycatch (e.g., moths) compared to when lights were turned off. Because “cleaner” samples with less bycatch are much easier to process and result in better data quality, light bulbs will be disabled in the traps used in operations.
- Grazing livestock caused damage to traps at the Central Plains Experimental Range. To mitigate this damage, aluminum foil was wrapped around wires and battery leads. This technique will be useful at other grazed sites.
- At sites where trees are not available to hang mosquito traps (e.g. Central Plains Experimental Range, Sterling), NEON tested shepherd’s hooks as an alternative structure from which to hang traps. Some models were found to be too thin and tended to break or bend or have too wide of a hook that the trap could swing off from. In loose or sandy soils, hooks required stabilization (e.g., posts were sunk into a plastic bucket with concrete). By trial and error, NEON identified a robust shepherd’s hook deployment strategy that can accommodate mosquito traps at any sites where appropriate structures are not present.

6.2 Sample Processing

Minimal processing of mosquito samples will occur within each Domain lab; mosquitoes will be identified at external facilities. This approach will minimize identification errors due to the expertise

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

needed to accurately identify mosquitos to species. NEON will also outsource all molecular, genetic, and pathogenic analyses of samples.

6.2.1 Sample Handling

After being live-trapped in the field, mosquitoes will be frozen, transferred to and stored in sample vials (dry) at -80°C until being sent to identification and pathogen testing facilities. These facilities will identify mosquitoes to species based on visual examination of external morphology. From among mosquitoes collected in each sampling event at each trap, either a set proportion or a fixed number of mosquitoes will be identified, based on catch rates during the first few years of operations. Initially, up to 200 mosquitoes will be identified and enumerated by species and sex from each collected trap. When more than 200 mosquitoes are collected, a representative subsample of ~200 individuals will be identified to estimate species composition. The uncounted proportion will be quantified by weight to estimate abundance.

After identification, individuals of target species will be destructively tested for pathogens, as detailed in the TOS Science Design for Vectors and Pathogens (AD[03]). In addition, ten individuals of each species per domain per year will be removed before testing to serve as vouchers. These vouchers and all enumerated, identified mosquitoes will be sent to an archive facility.

6.3 Taxonomic verifications and reference collections

Some mosquitoes sent for morphology-based taxonomic analysis will also have tissue sent for DNA sequencing of the Folmer region of the CO1 gene (aka DNA barcoding; Folmer et al. 1994). Exact numbers of specimens expertly identified versus identified through DNA sequencing will depend on the abundance and diversity of collected mosquitoes and annual funding (with up to 95 individuals sequenced per domain per year). However, mosquitoes that are rare, particularly difficult to identify, or poorly represented in previous collection events will be prioritized for DNA sequencing. DNA sequence data will supplement expert identifications and provide greater resolution in cases of poorly resolved taxonomy (e.g., the *Anopheles crucians* complex) or cryptic species.

The combined use of expert identifications and sequencing will also improve the ability of the broader scientific community to make accurate identifications. As NEON accumulates and publishes sequence data on specimens that have also been identified by mosquito experts, the quality and quantity of sequence information available for many mosquito species will grow. Publically available DNA reference sequences will aid in understanding the inter- and intra-specific variation within mosquito populations, support accurate identification of mosquito specimens by non-experts and reveal the presence of cryptic species. The specimens for the NEON DNA sequence reference library, to date, were collected during field prototype campaigns or obtained from museum archives (Gibson et al. 2012). All assembled resources for each specimen - sequence data, photos, and other ecological information - can be publicly accessed online from the Barcode of Life Datasystem (BOLD, <http://boldsystems.org/>). Two BOLD

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

projects have been established; a project for operational data ('Mosquito sequences DNA barcode' or project code MOSN) and a project for prototype data ('NEON Prototype Mosquito sequences DNA barcode' or project code MOSP).

6.3.1 Archive

All identified specimens and any extracted DNA will be archived in existing facilities and available for use by the scientific community.

6.4 Spatial Distribution of Sampling

6.4.1 Selecting Plot Locations within Sites

Mosquito trapping plots will be chosen using a stratified random approach, maintaining stratification across dominant vegetation types ($\geq 5\%$ of total cover) at each site (AD[02]). Ten plots will be selected for long-term sampling such that the number of plots per vegetation type is proportional to the percent cover at the site. Plot locations will be constrained to fall within 30 m of roads in order to reduce travel time associated with each sampling event. Though constraining plot locations may reduce the statistical area of inference, the benefit of maintaining adequate sample size given budget constraints is deemed worth this cost. However, traps will be placed a minimum of 300 meters apart to maintain independence, even at the cost of reducing the total number of traps at small sites.

Mosquito species differ substantially in their ability to fly long distances, and species might travel anywhere from 150 meters to over a kilometer during an 8-hour sampling window. A study seeking to characterize the population of a single species of mosquito would likely choose minimum inter-trap distances scaled to the flight ability of the particular species under consideration. However, the NEON Mosquito Technical Working Group selected 300 meters as a minimum distance between traps because NEON seeks to characterize diversity of the whole mosquito community. If time allows in initial operations, the NEON Mosquito Technical Working Group will analyze the diversity data generated by the mosquito program and evaluate whether the 300-meter threshold is appropriate given the known dispersal distances of the captured mosquito community.

6.5 Temporal Distribution of Sampling

6.5.1 Sampling Session

A single session (referred to as a "bout") of mosquito sampling will consist of three CO₂ light trapping periods - two consecutive trapping nights and the intervening day (Figure 2) - at up to 10 plots per site (one trap per plot) during field sampling. This 40-hour sampling period will catch both day- and night-active mosquitoes and thus maximize community representation by covering the full spectrum of mosquito activity. Samples will collectively be used to characterize mosquito abundance, diversity and phenology at the site level. The samples collected during the three trapping events within a bout will be

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

kept separate to determine the unique species composition of day-time vs. night-time sampling. In addition, data at the resolution of a trap-night is useful for comparison with other mosquito monitoring efforts. In 2018, the second night of sampling was dropped for cost saving reasons.

Off-season sampling sessions will consist of one trap deployed at each of three plots at the core site only for a single night (see details in section 6.5.4).




Figure 2. Field season sample session timing of mosquito trap deployment

Data will be reported at the trap level (i.e., each trap day/night by location within a site). In addition to mosquito abundance and diversity, mosquito phenology data will be collected. Phenology metrics (e.g., peak abundance, dormant time, first emergence) could be derived from abundance and diversity data collected during field sampling.

6.5.2 Temperature Threshold

The goal of phenology sampling for mosquitoes is to register at least one zero capture before the start of each field season, such that data from early field season sampling are known to be the first dates of occurrence for a particular observed species (for sites that do not have mosquito activity year-round). Temperature plays an important role in determining whether mosquitoes are active and able to be captured by CO₂ trapping. Thus, traps will only be deployed if the average daily high temperature for the previous 5 days was >10 °C. In cases where this criterion is not met, the presumed mosquito catch is zero and off-season sampling will continue. This threshold will be applied to both field season and off season sampling, and, if the threshold is not met the day prior to the first night of sampling during a sampling session, the entire sampling session will be canceled.

Initial Temperature Thresholds

Prior to 2017, the temperature threshold for deployment during off-season was based on a review of the literature and required that the average daily high temperature for the previous 5 days was greater than 4 °C (e.g., Cossins and Bowler 1987). In 2017, NEON scientists took a data driven approach where

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

observed mosquito collections at core sites from 2016 were compared against NOAA NCDC data from nearby weather stations. Core sites represented a wide geographic range across the network and diversity of environments characterized by disparate temperature and moisture profiles (D01: HARV, D02:SCBI, D05:UNDE, D06:KONZ, D09:WOOD, D10:CPER, D14:SRER, D15:ONAQ). Core sites were selected for this analysis if 1) they had a complete season of data for 2016, 2) they experienced an off season in 2016 (this criterion disqualified sites that conduct year-round sampling), and 3) had at least one observed ‘no capture’ during the 2016 off season.

For this effort, NEON collated ten years of observed NOAA NCDC data from the nearest weather station to each site (NOAA NCDC between 2006–2015; Menne *et al.* 2012). Eligible weather stations were the closest to the NEON site that met location and elevation criteria; weather stations were required to be closer than 150 km and within 300 m of the altitude of the NEON site to receive consideration. One weather station was selected for each site (the closest station that met the above criteria) and for each date the 5-day running average of observed daily maximum temperatures (TMAX) for the prior 5 days was calculated (for each date from 10 years of observed temperature data).

The goal of the analysis was to determine the optimal threshold where the number of off season mosquito absences were minimized but at least one mosquito absence would be documented. Thus, NEON evaluated a range of possible temperature thresholds from 4 to 25 °C. Using the mosquito collection data and weather data described above, NEON staff calculated the number of off-season bouts that would have been implemented at these eight core sites given a particular temperature threshold. In the 4 °C threshold scenario, eight core sites captured the absence of mosquitoes for about 7.5 weeks (1.7 SE) before documenting mosquito emergence. This analysis revealed that by raising the temperature threshold from 4 °C to 10 °C, each site would reduce their sampling windows by 3 weeks on average while still documenting the absence of mosquitoes during at least one collection at all but one of the core sites. Combined with the directive to initiate field sampling whenever mosquitoes are observed flying, NEON adopted the change in 2017 as a cost saving measure.

Sample Scheduling

As a separate effort, NEON staff also used a data driven approach to determine approximate site-specific sampling start dates (included in Appendix D of the Mosquito Protocol (RD[04])) to assist with domain staffing and scheduling. Selected data and weather stations followed the same criteria used above for the temperature threshold evaluation (above). One weather station was selected for each site (the closest station that met the criteria) and the 5-day running average of observed daily maximum temperatures (TMAX) was calculated for each year of observed temperature data. For every site, the provided estimated start of sampling was determined as the average Julian date that temperatures first met the 5-day running average criteria. Likewise, the estimated sampling end date is the average Julian date where the running average of TMAX drops below 10 °C.

6.5.3 Field Season Sampling

Mosquitoes will be sampled using the same schedule at all domains where mosquitoes are present, irrespective of local density. During the field season, mosquito sampling sessions will occur every other week at the core site and monthly at each of the relocatable sites (Figure 3). The number of weeks in a field season varies among domains and the criteria for the beginning and end of the field season are detailed below (Section 6.5.4). In addition, the core site will be the indicator site for off-season sampling.




Figure 3. Field season mosquito sampling schedule for a representative domain. Sampling typically occurs at one site each week, alternating between the core and relocatable sites. The number of weeks in the field season varies among domains.

6.5.4 Seasonal Timing

Mosquitoes display seasonal abundance and activity patterns that vary among species and regions. Most of these patterns exhibit some general level of consistency. For example, at many NEON sites, mosquitoes are absent during part of each year where winter months are associated with unfavorable climatic conditions and mosquitoes re-emerge in early/mid spring. At seasonal sites, mosquito sampling is unnecessary during the winter and will be suspended until mosquitoes become active in the spring. This strategy requires that sampling be initiated and maintained in concert with seasonal mosquito activity patterns at a site, such that the phenology of mosquito species is adequately captured. This is

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

especially important in the context of the mosquito pathogen program, as disease transmission patterns are likely to be impacted by climate change-mediated alterations in mosquito distribution and phenology. Determining appropriate staffing models to ensure that mosquito phenology is captured is complicated by the fact that the precise timing of these seasonal events can vary considerably among years. While logistically attractive, an approach that uses a fixed calendar date to determine when to stop and restart sampling each year is unacceptable because it will frequently result in the start/end of sampling being mistimed because of interannual phenological variation. As an alternative, NEON will employ an “off-season” sampling plan distinct from those employed during field season sampling in order to track mosquito activity (Figure 4).




Figure 4. Annual mosquito sampling timeline for a representative domain. Trapping occurs all year at the core site, with more traps during the warm part of the year when mosquitoes are most active. Note: from 2013 to 2016, a 4C threshold was used to determine the implementation of bouts; in 2017 and later, a revised threshold of 10C was used based on observed captures during the first four years of sampling.

Within a domain, the end of the regular sampling season will occur following three consecutive zero-catch sampling sessions at the core site (Figure 5). A field season zero-catch at the core site (no mosquitoes caught in all 10 traps) will trigger off-season sampling at the core site the following week (the intervening week before the next field season session at the core site). Note that field season sampling still occurs at the relocatable site during this intervening week. If the off-season sample is a zero-catch and the following sampling week at the core site is also a zero-catch (3 consecutive weeks of zero-catches at the core site, consisting of 2 field season sessions and 1 off season session), all field season sampling at the core and relocatable sites will stop and weekly off-season sampling at the core site will continue until a positive mosquito catch triggers the resumption of field season sampling (e.g., in the spring of the following year). Zero-catches at relocatables have no effect on the off-season/field-season transition.

Off-season sampling will continue once per week (so long as temperatures are above the threshold, see 6.5.2) until at least one mosquito is collected. This will initiate the resumption of field season sampling (Figure 5), including both the core site (10 traps every 2 weeks) and relocatable sites (10 traps every 4 weeks).




Figure 5. An example of mosquito sampling at a core site. The example begins with and ends with weekly off-season sampling and shows a brief field season. Sampling at relocatable sites is not shown but would occur in the intervening weeks (4-14).

6.6 Logistics and Adaptability

Once a set of long-term mosquito sampling plots is selected, NEON will generally maintain a fixed plot design. In some cases results may point to a plot being dropped in favor of sampling at a new and potentially more productive or informative location. Decisions about if and when to change the

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

sampling design will be made in consultation with an expert review committee and Observatory Director.

<i>Title:</i> TOS Science Design for Mosquito Abundance, Diversity, and Phenology		<i>Date:</i> 06/14/2018
<i>NEON Doc. #:</i> NEON.DOC.000910	<i>Author:</i> K. LeVan	<i>Revision:</i> B

7 REFERENCES

- AIBSnews. 2007. NEON design 2007. *BioScience* 57:198–200.
- Buckner, E. A., M. S. Blackmore, S. W. Golladay, and A. P. Covich. 2010. Weather and landscape factors associated with adult mosquito abundance in southwestern Georgia, U.S.A. *Journal of Vector Ecology* 36:269–278.
- Burkett-Cadena, N. D., M. D. Eubanks, and T. R. Unnasch. 2008. Preference of female mosquitoes for natural and artificial resting sites. *Journal of the American Mosquito Control Association* 24:228–235.
- Burkett-Cadena, N. D., C. J. W. McClure, R. A. Ligon, S. P. Graham, G. E. Hill, S. S. Ditchkoff, M. D. Eubanks, H. K. Hassan, and R. Thomas. 2011. Host Reproductive Phenology Drives Seasonal Patterns of Host Use in Mosquitoes. *PloS one* 6.
- Committee on Grand Challenges in Environmental Sciences, O. C. for the C. on G. C. in E. S. 2001. Grand Challenges in Environmental Sciences. Page 106. The National Academies Press, Washington, D.C.
- Cossins, A. R., and K. Bowler. 1987. *Temperature Biology of Animals*. Chapman & Hall, New York.
- Daubney, R., and J. R. Hudson. 1931. Enzootic Hepatitis or Rift Valley Fever. An Undescribed Virus Disease of Sheep Cattle and Man from East Africa. *Journal of Pathology and Bacteriology* 34:545–579.
- Epstein, P. R., H. F. Diaz, S. Elias, G. Grabherr, N. E. Graham, W. J. M. Martens, E. Mosley-Thompson, and J. Susskind. 1998. Biological and Physical Signs of Climate Change: Focus on Mosquito-borne Diseases. *Bulletin of the American Meteorological Society* 79:409–417. American Meteorological Society.
- Gage, K. L., T. R. Burkot, R. J. Eisen, and E. B. Hayes. 2008. Climate and vectorborne diseases. *American Journal of Preventive Medicine* 35:436–450.
- Gibson, C. M., R. H. Kao, K. K. Blevins, and P. D. Travers. 2012. Integrative taxonomy for continental-scale terrestrial insect observations. *PLoS ONE* 7:e37528. doi: 10.1371/journal.pone.0037528.
- Hoel, D. F., D. L. Kline, and S. A. Allan. 2009. Evaluation of six mosquito traps for collection of *Aedes albopictus* and associated mosquito species in a suburban setting in north central Florida. *Journal Of The American Mosquito Control Association* 25:47–57.
- Hongoh, V., L. Berrang-Ford, M. E. Scott, and L. R. Lindsay. 2011. Expanding geographical distribution of the mosquito, *Culex pipiens*, in Canada under climate change. *Applied Geography* 33:53–62. Elsevier Ltd.

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

- Keller, M., D. S. Schimel, W. W. Hargrove, and F. M. Hoffman. 2008. A continental strategy for the National Ecological Observatory Network. *Frontiers in Ecology and the Environment* 6:282–284. Eco Soc America. doi: 10.1890/1540-9295(2008)6[282:ACSFTN]2.0.CO;2.
- Kissling, R. E., R. W. Chamberlain, M. E. Eidson, R. K. Sikes, and M. A. Bucca. 1954. Studies on the North American arthropod-borne encephalitides. 2. Eastern Equine Encephalites in Horses. *American Journal of Hygiene* 60:237–250.
- Komar, N., B. J. Biggerstaff, N. A. Panella, and R. J. Kent Crockett. 2011. The Centers for Disease Control and Prevention Resting Trap: A Novel Device for Collecting Resting Mosquitoes. doi: 10.2987/09-5900.1.
- Komar, N., R. J. Pollack, and A. Spielman. 1995. A nestable fiber pot for sampling resting mosquitoes. *Journal of the American Mosquito Control Association* 11:463–467.
- Krockel, U., A. Rose, A. E. Eiras, and M. Geier. 2006. New tools for surveillance of adult yellow fever mosquitoes: comparison of trap catches with human landing rates in an urban environment. *Journal of the American Mosquito Control Association* 22:229–238. doi: 10.2987/8756-971X(2006)22[229:NTFSOA]2.0.CO;2.
- LaDeau, S. L., A. M. Kilpatrick, and P. P. Marra. 2007. West Nile virus emergence and large-scale declines of North American bird populations. *Nature* 447:710–713.
- LaDeau, S. L., P. P. Marra, A. M. Kilpatrick, and C. A. Calder. 2008. West Nile Virus Revisited: Consequences for North American Ecology. *BioScience* 58:937–946. University of California Press on behalf of the American Institute of Biological Sciences.
- Lounibos, L. P. 2002. Invasions by Insect Vectors of Human Disease. *Annual Review of Entomology* 47:233–266. Annual Reviews.
- Marra, P. P., S. Griffing, C. Caffrey, A. M. Kilpatrick, R. McLean, C. Brand, E. M. I. Saito, A. P. Dupuis, L. Kramer, and R. Novak. 2004. West Nile Virus and Wildlife. *BioScience* 54:393–402. University of California Press on behalf of the American Institute of Biological Sciences.
- Meeraus, W. H., J. S. Armistead, and J. R. Arias. 2008. Field comparison of novel and gold standard traps for collecting *Aedes albopictus* in Northern Virginia. *Journal of the American Mosquito Control Association* 24:244–248.
- Menne, M. J., I. Durre, B. Korzeniewski, S. McNeal, K. Thomas, X. Yin, S. Anthony, R. Ray, R. S. Vose, B. E. Gleason, and T. G. Houston (2012): Global Historical Climatology Network - Daily (GHCN-Daily), Version 3. [subset: 2006-01-01 to 2015-12-31]. NOAA National Climatic Data Center. doi:10.7289/V5D21VHZ

Title: TOS Science Design for Mosquito Abundance, Diversity, and Phenology		Date: 06/14/2018
NEON Doc. #: NEON.DOC.000910	Author: K. LeVan	Revision: B

Reisen, W. K. 2010. Landscape Epidemiology of Vector-Borne Diseases. *Annual Review of Entomology* 55:461–483.

Reiter, P. 1983. A Portable, Battery-Powered Trap for Collecting Gravid Culex Mosquitoes. *Mosquito News* 43:496–498.

Reiter, P. 2001. Climate change and mosquito-borne disease. *Environmental health perspectives* 109 Suppl:141–161.

Van Riper III, C., S. G. Van Riper, M. L. Goff, and M. Laird. 1986. The Epizootiology and Ecological Significance of Malaria in Hawaiian Land Birds. *Ecological Monographs* 56:327–344. Ecological Society of America. doi: 10.2307/1942550.

Schimel, D., M. Keller, S. Berukoff, R. Kao, H. Loescher, H. Powell, T. Kampe, D. Moore, and W. Gram. 2011a. 2011 Science Strategy: enabling continental-scale ecological forecasting.

Schimel, D., M. Keller, S. Berukoff, R. Kao, H. Loescher, H. Powell, T. Kampe, D. Moore, W. Gram, D. Barnett, R. Gallery, C. Gibson, K. Goodman, C. Meier, S. Parker, L. Pitelka, Y. Springer, and R. U. K Thibault. 2011b. NEON SCIENCE STRATEGY: Enabling Continental-Scale Ecological Forecasting. Page 56.

Service, M. W. 1993. *Mosquito ecology: Field sampling methods*, Second edition. Mosquito ecology: Field sampling methods, Second edition.

Silver, J. B. 2008. *Mosquito Ecology*. Nature. Springer.

Sudia, W. D., and R. W. Chamberlain. 1962. Battery-operated light trap, an improved model. *Mosquito News* 22:126–129.

Williams, G. M., and J. B. Gingrich. 2007. Comparison of light traps, gravid traps, and resting boxes for West Nile virus surveillance. *Journal of vector ecology : journal of the Society for Vector Ecology* 32:285–291. doi: 10.3376/1081-1710(2007)32.