



NEON USER GUIDE TO COARSE DOWNED WOOD LOG SURVEY (DP1.10010.001)

PREPARED BY	ORGANIZATION
Katherine Jones	SCI
Courtney Meier	SCI
Samuel Simkin	SCI



CHANGE RECORD

REVISION	DATE	DESCRIPTION OF CHANGE
A	12/15/2017	Initial Release
B	07/07/2020	Included general statement about usage of neonUtilities R package and statement about possible location changes. Updated taxonomy information. Included 'Sampling Design Changes' section. Updated 'Spatial Resolution and Extent' section. Made other clarifications within document.
C	04/11/2022	Added language in section 4 Taxonomy addressing RTE species obfuscation in the data. Updated section 5.3 Data Revision with latest information regarding data release. Updated information regarding the geoNEON package



<i>Title:</i> NEON User Guide to Coarse Downed Wood Log Survey (DP1.10010.001)	<i>Date:</i> 04/11/2022
<i>Author:</i> Katherine Jones	<i>Revision:</i> C

TABLE OF CONTENTS

1 DESCRIPTION	1
1.1 Purpose	1
1.2 Scope	1
2 RELATED DOCUMENTS AND ACRONYMS	2
2.1 Associated Documents	2
2.2 Acronyms	2
3 DATA PRODUCT DESCRIPTION	3
3.1 Spatial Sampling Design	3
3.2 Temporal Sampling Design	4
3.3 Sampling Design Changes	4
3.4 Variables Reported	5
3.5 Spatial Resolution and Extent	5
3.6 Temporal Resolution and Extent	6
3.7 Associated Data Streams	7
3.8 Product Instances	7
3.9 Data Relationships	7
3.10 Special Considerations	8
4 TAXONOMY	8
5 DATA QUALITY	9
5.1 Data Entry Constraint and Validation	9
5.2 Automated Data Processing Steps	9
5.3 Data Revision	10
5.4 Quality Flagging	10
6 REFERENCES	10

LIST OF TABLES AND FIGURES

Table 1	Descriptions of the dataQF codes for quality flagging	10
---------	---	----



<i>Title:</i> NEON User Guide to Coarse Downed Wood Log Survey (DP1.10010.001)	<i>Date:</i> 04/11/2022
<i>Author:</i> Katherine Jones	<i>Revision:</i> C

Figure 1 Illustration of three randomly oriented LIDS transects (arrow vectors) superimposed over a 20m x 20m plot (left), and a 40m x 40m plot (right). There is a 3m gap between the sampling reference point (i.e. the plot centroid) and the start of each transect, which minimizes the chance that CDW particles close to the plot centroid get double- or triple-counted. Gray shapes represent CDW particles, dashed lines show the location and orientation of cross-sectional area measurement(s) (ovals) for those particles that intersect a LIDS transect, and capped black lines imply the particles' central axes that are measured for length. Note that CDW particles selected for tally and/or bulk density sampling may lie outside of the plot boundary. 4



1 DESCRIPTION

1.1 Purpose

This document provides an overview of the data included in this NEON Level 1 data product, the quality controlled product generated from raw Level 0 data, and associated metadata. In the NEON data products framework, the raw data collected in the field, for example the dimensions of logs, are considered the lowest level (Level 0). Raw data that have been quality checked via the steps detailed herein, as well as simple metrics that emerge from the raw data are considered Level 1 data products.

The text herein provides a discussion of measurement theory and implementation, data product provenance, quality assurance and control methods used, and approximations and/or assumptions made during L1 data creation.

1.2 Scope

This document describes the steps needed to generate the L1 data product, Coarse Downed Wood Log Survey - the tally and raw measurement of coarse downed wood particles ≥ 2 cm diameter and ≥ 1 m length - and associated metadata from input data. This document also provides details relevant to the publication of the data products via the NEON data portal, with additional detail available in the file, NEON Data Variables for Coarse Downed Wood Log Survey (DP1.10010.001) (AD[05]), provided in the download package for this data product.

This document describes the process for ingesting and performing automated quality assurance and control procedures on the data collected in the field pertaining to TOS Protocol and Procedure: Coarse Downed Wood (AD[07]). The raw data that are processed in this document are detailed in the file, NEON Raw Data Validation for Coarse Downed Wood Log Survey (DP0.10010.001) (AD[04]), provided in the download package for this data product. Please note that raw data products (denoted by 'DP0') may not always have the same numbers (e.g., '10010') as the corresponding L1 data product.



2 RELATED DOCUMENTS AND ACRONYMS

2.1 Associated Documents

AD[01]	NEON.DOC.000001	NEON Observatory Design (NOD) Requirements
AD[02]	NEON.DOC.000913	TOS Science Design for Spatial Sampling
AD[03]	NEON.DOC.002652	NEON Data Products Catalog
AD[04]	NEON.DP0.10010.001_dataValidation.csv	NEON Raw Data Validation for Coarse Downed Wood Log Survey (DP0.10010.001)
AD[05]	NEON.DP1.10010.001_variables.csv	NEON Data Variables for Coarse Downed Wood Log Survey (DP1.10010.001)
AD[06]	NEON.DOC.000914	TOS Science Design for Plant Biomass, Productivity, and Leaf Area Index
AD[07]	NEON.DOC.001711	TOS Protocol and Procedure: Coarse Downed Wood
AD[08]	NEON.DOC.000913	TOS Science Design for Spatial Sampling
AD[09]	NEON.DOC.000913	TOS Science Design for Plant Diversity
AD[10]	NEON.DOC.000008	NEON Acronym List
AD[11]	NEON.DOC.000243	NEON Glossary of Terms
AD[12]	NEON.DOC.000987	TOS Protocol and Procedure: Measurement of Vegetation Structure
AD[13]	NEON.DOC.001710	TOS Protocol and Procedure: Measurement of Litterfall and Fine Woody Debris
AD[14]	NEON.DOC.004825	NEON Algorithm Theoretical Basis Document: OS Generic Transitions
AD[15]	Available on NEON data portal	NEON's Ingest Conversion Language (NICL) specifications

2.2 Acronyms

Acronym	Definition
CDW	Coarse Downed Wood
DST	Decay class x size category x taxonID combination
LIDS	Line Intercept Distance Sampling



3 DATA PRODUCT DESCRIPTION

The Coarse Downed Wood Log Survey data product provides volume estimates of downed logs ≥ 2 cm diameter across NEON sites. Tally surveys are completed according to the Line Intercept Distance Sampling method (LIDS; Affleck 2008). LIDS sampling tallies logs with probability proportional to volume and restricts the search for logs to a transect or group of radial transects at each sampling point (Affleck 2008, 2010). By using transects, detection errors in brushy or complex terrain are minimized compared to other techniques that require searching for logs over large areas (Jordan et al. 2004). The length of the transect is not fixed with the LIDS method; instead, the length of the transect increases for large-volume logs, ensuring that a representative sample of large logs is tallied across multiple field sites (Affleck 2008).

Qualifying particles of CDW are divided into three different diameter size categories, as defined by Keller et al. (2004): 2-5 cm, 5-10 cm, and ≥ 10 cm diameter. To qualify for inclusion, logs or portions of logs within a given size category must also be ≥ 1 m in length (Harmon and Sexton 1996). Dead trees (i.e. snags) that have not yet fallen to an angle $> 45^\circ$ from vertical are accounted for via the Vegetation Structure protocol (AD[12]), and logs that are suspended in the air > 2 m above the ground at the transect intersection point are ignored. Woody particles with diameter < 2 cm at the transect intersection point are considered fine woody debris and are sampled according to the Litterfall and Fine Woody Debris protocol (AD[13]).

All logs are classified into unique decay class x size category x taxonID (DST) combinations. Size category is based on diameter but is supplemented by additional physical characteristics such length that allow for calculation of volume for each DST. Decay class is supplemented with data on the presence/absence of leaves and branches, % bark cover, etc. The distance and azimuth angle from the transect origin is also recorded for mapping purposes.

3.1 Spatial Sampling Design

Coarse downed wood surveys are completed at all terrestrial NEON sites that contain woody vegetation. At sites where qualifying CDW particles are present, sampling occurs at each Tower Plot (20-30 plots), and a subset of randomly selected Distributed Plots (20 plots maximum), for a maximum of 50 plots at any given site. At each plot, three LIDS transects radiate outward from the plot centroid, with 120° separating each transect (Figure 1). The azimuthal orientation of each group of transects is chosen randomly for each plot, in order to minimize effects of topography, directional blowdown, logging management, etc., on selection of CDW particles across all sampling locations. See TOS Science Design for Plant Biomass, Productivity, and Leaf Area Index (AD[06]), TOS Protocol and Procedure: Coarse Downed Wood (AD[07]), TOS Science Design for Spatial Sampling (AD[08]) and TOS Science Design for Plant Diversity (AD[09]) for further details on sampling design, specific protocol steps, plot allocation, and species identification, respectively.

As much as possible, sampling occurs in the same locations over the lifetime of the Observatory. However, over time some sampling locations may become impossible to sample, due to disturbance or other local changes. When this occurs, the location and its location ID are retired. A location may also shift to slightly different coordinates. Refer to the locations endpoint of the NEON API for details about locations that have been moved or retired: <https://data.neonscience.org/data-api/endpoints/locations/>




Figure 1: Illustration of three randomly oriented LIDS transects (arrow vectors) superimposed over a 20m x 20m plot (left), and a 40m x 40m plot (right). There is a 3m gap between the sampling reference point (i.e. the plot centroid) and the start of each transect, which minimizes the chance that CDW particles close to the plot centroid get double- or triple-counted. Gray shapes represent CDW particles, dashed lines show the location and orientation of cross-sectional area measurement(s) (ovals) for those particles that intersect a LIDS transect, and capped black lines imply the particles' central axes that are measured for length. Note that CDW particles selected for tally and/or bulk density sampling may lie outside of the plot boundary.

3.2 Temporal Sampling Design

Coarse downed wood tally bouts are completed in all selected plots once every five years, typically temporally distributed such that each site within a domain is sampled in a different year. Sampling at each site should take no longer than 3 months to complete. New coarse downed wood particles are typically produced during periods of high wind or rain events associated with seasonal storm activity. Sampling bouts at any given site are scheduled to occur after the expected storm season. For example, in the Southeast (D03) and Atlantic Neotropical domains (D04), CDW tally sampling occurs during either the winter/spring season, outside of the summer/fall hurricane seasons. In contrast, in the Pacific Northwest (D16), most storm activity occurs primarily in the winter, and CDW tally sampling occurs in the summer/autumn seasons.

3.3 Sampling Design Changes

In 2018 the coarse downed wood tally sampling frequency was reduced from once every three years to once every five years as a cost-savings measure.



3.4 Variables Reported

All variables reported from the field or laboratory (L0 data) are listed in the file, NEON Raw Data Validation for Coarse Downed Wood Log Survey (DP0.10010.001) (AD[04]). All variables reported in the published data (L1 data) are also provided separately in the file, NEON Data Variables for Coarse Downed Wood Log Survey (DP1.10010.001) (AD[05]).

Field names have been standardized with Darwin Core terms (<http://rs.tdwg.org/dwc/>; accessed 16 February 2014), the Global Biodiversity Information Facility vocabularies (<http://rs.gbif.org/vocabulary/gbif/>; accessed 16 February 2014), and the VegCore data dictionary (<https://projects.nceas.ucsb.edu/nceas/projects/bien/wiki/VegCore>; accessed 16 February 2014), where applicable. NEON TOS spatial data employs the World Geodetic System 1984 (WGS84) for its fundamental reference datum and GEOID09 for its reference gravitational ellipsoid. Latitudes and longitudes are denoted in decimal notation to six decimal places, with longitudes indicated as negative west of the Greenwich meridian.

Some variables described in this document may be for NEON internal use only and will not appear in downloaded data.

3.5 Spatial Resolution and Extent

The finest resolution at which spatial data are reported is the point location of the log along a transect.

The spatial hierarchy from finest to coarsest resolution is:

point (see below to calculate) → **plotID** → **siteID** → **domainID**

The basic spatial data included in the data downloaded include the latitude, longitude, and elevation of the centroid of the plot where sampling occurred + associated uncertainty due to GPS error and plot width. Shapefiles of all NEON Terrestrial Observation System sampling locations can be found in the Document Library: <http://data.neonscience.org/documents>.

To derive a more precise estimate of the location of each log, there are two options:

- Use the getLocTOS function from the geoNEON package, available here: <https://github.com/NEONScience/NEON-geolocation>
- Or follow these steps to perform the same calculation:
 1. The namedLocation field in the data is the named location of the plot.
 2. Use the API (<http://data.neonscience.org/api>; e.g. http://data.neonscience.org/api/v0/locations/HARV_052.basePlot.cdw) to query for elevation("locationElevation"), easting("locationUtmEasting"), northing("locationUtmNorthing"), coordinateUncertainty ("Value for Coordinate uncertainty"), elevationUncertainty ("Value for Elevation uncertainty"), and utmZone ("locationUtmZone") as inputs to the next step.
 3. Calculate the geolocation of each log according to the equation:



Title: NEON User Guide to Coarse Downed Wood Log Survey (DP1.10010.001)	Date: 04/11/2022
Author: Katherine Jones	Revision: C

$$Easting = locationUtmEasting + d * \sin \theta \quad (1)$$

and

$$Northing = locationUtmNorthing + d * \cos \theta \quad (2)$$

where,

$$\theta = \frac{logsAzimuth * \pi}{180} \quad (3)$$

$$locationUtmEasting = \quad (4)$$

the easting value associated with the plotID (plot centroid)

$$locationUtmNorthing = \quad (5)$$

the northing value associated with the plotID (plot centroid)

$$d = logDistance + 3 \quad (6)$$

Note: logDistance is the distance along transect, but transects start 3 m from plot centroid.

- Increase coordinateUncertainty by an appropriate amount to account for error introduced by navigating along the transect. Additional error may be introduced due to tape stretching and drift from a compass heading.

3.6 Temporal Resolution and Extent

The finest resolution at which temporal data are reported for a given log tally record is the range between **startDate** and **endDate** associated with when the transect was tallied.

date (date range a transect was tallied) → **eventID** (bout-level identifier)



<i>Title:</i> NEON User Guide to Coarse Downed Wood Log Survey (DP1.10010.001)	<i>Date:</i> 04/11/2022
<i>Author:</i> Katherine Jones	<i>Revision:</i> C

3.7 Associated Data Streams

The Coarse Downed Wood Log Survey data product (DP1.10010.001) is tightly integrated with the Coarse Downed Wood Bulk Density Sampling data product (DP1.10014.001). Together, these two data products allow calculation of CDW dry mass per sizeCategory per decayClass per taxonID. Although these two data products are related, logs that are tagged during Log Survey are specifically avoided for Bulk Density Sampling.

The Woody and Non-Herbaceous Perennial Vegetation Structure data products (DP1.10098.001 and DP1.10045.001, respectively) are also linked to the CDW Log Survey product due to the fact that individuals tagged and measured while living may be sampled in the CDW Log Survey once they have fallen to the ground. In the event that data in CDW Log Survey are collected from an individual that was previously part of one of these two datasets, the **vstTagID** field in the CDW dataset may be matched to the **tagID** field in the Woody and Non-Woody Perennial Vegetation Structure products.

3.8 Product Instances

A maximum of 50 plots will be selected for coarse downed wood survey, once every five years, at a given site. Each plot has 3 transects, with the total number of logs tallied along all 3 transects for a given plot ranging between 0-50. Thus, each site should generate no more than approximately 2,500 data product instances over a five year period.

3.9 Data Relationships

The protocol dictates that each transect is surveyed only once in a sampling year. If no particles ≥ 2 cm diameter and ≥ 1 m length are present, only a single record will be generated for the transect, and these records will have **targetTaxaPresent**= 'No', indicating that the transect was checked and no qualifying particles were discovered. For transects where **targetTaxaPresent**= 'Yes', each qualifying particle along the transect results in a distinct record. Duplicates and/or missing data may exist where protocol and/or data entry aberrations have occurred; users should check data carefully for anomalies before joining field tally table with other data products.

cdw_fieldtally.csv → One record expected per **plotID** + **lidsAzimuth** + **logID** combination per year.

An **individualID** and **individualBarcode** is generated for each tallied particle with logMaxDiameter ≥ 10 cm, and the **individualID** is built from the **logID** for these particles. No **individualID** is generated for logs with logMaxDiameter < 10 cm. For particles < 10 cm diameter, **logID** values are incremented values that reset for each **plotID**, and are denoted with the letter "L" preceding an incremented value (e.g. L005), and the **individualID** field will be null. Note that the CDW **individualID** cannot be matched with the **individualID** created for other NEON plant data products (described below).

Recently downed logs in the CDW Log Survey dataset that were previously measured while standing according to one or more protocols are identified in the CDW Log Survey dataset by the **vstTagID** field. To match **vstTagID** with the linked **individualID** reported in other NEON data products, an end-user must perform the following steps:



1. Flag **vstTagID** with leading zeroes until the total string length is 6 characters
2. Concatenate: "NEON.PLA.D##.SITE" + "." + "**vstTagID**", where:
 - D## = **domainID**
 - SITE = **siteID**
3. Match the resulting string (e.g., NEON.PLA.D16.WREF.000047) with **individualID** in any/all of the following associated data products:
 - Woody Plant Vegetation Structure (DP1.10095.001)
 - Plant Canopy Leaf Mass per Area (DP1.10048.001)
 - Plant Foliar Stable Isotopes (DP1.10053.001)
 - Plant Phenology Observations (DP1.10055.001)

Data downloaded from the NEON Data Portal are provided in separate data files for each site and month requested. The neonUtilities R package contains functions to merge these files across sites and months into a single file for each table described above. The neonUtilities package is available from the Comprehensive R Archive Network (CRAN; <https://cran.r-project.org/web/packages/neonUtilities/index.html>) and can be installed using the install.packages() function in R. For instructions on using neonUtilities to merge NEON data files, see the Download and Explore NEON Data tutorial on the NEON website: <https://www.neonscience.org/download-explore-neon-data>

3.10 Special Considerations

A subset of records for coarse downed wood particles < 10 cm diameter, measured between 2016-04-04 and 2017-06-19, were recorded without **decayClass**. These records have been flagged as a **dataQF**="decayClassNotRecorded". Affected records can be used to calculate volume of coarse downed wood, but do not contain sufficient information to calculate wood density or mass when combined with the Coarse Downed Wood Bulk Density product.

4 TAXONOMY

NEON manages taxonomic entries by maintaining a master taxonomy list based on the community standard, if one exists. Through the master taxonomy list, synonyms submitted in the data are converted to the appropriate name in use by the standard. The master taxonomy for plants is the USDA PLANTS Database (USDA, NRCS. 2014. <https://plants.usda.gov>). Taxon ID codes used to identify taxonomic concepts in the NEON master taxonomy list are alpha-numeric codes, 4-6 characters in length based on the accepted scientific name. Each code is composed of the first two letters of the genus, followed by the first two letters of the species and first letter of the terminal infraspecific name (if applicable) then, if needed, a tiebreaking number to address duplicate codes. Genus and family symbols are the first five (genus) or six (family) letters of the name, plus tiebreaking number (if needed). Symbols were first used in the Soil Conservation Service's National List of Scientific Plant Names (NLSPN) and have been perpetuated in the PLANTS system. The portions of the PLANTS Database included in the NEON plant master taxonomy list



includes native and naturalized plants present in NEON observatory sampling area including the Lower 48 U.S. States, Alaska, Hawaii, and Puerto Rico. NEON plans to keep the taxonomy updated in accordance with USDA PLANTS Database starting in 2020 and annually thereafter.

The master taxonomy list includes geographic range and nativity as described by the USDA PLANTS Database. A list for each NEON domain includes those species with ranges that overlap the domain as well as nativity designations - introduced or native - in that part of the range. Errors are generated if a species is reported at a location outside of its known range. If the record proves to be a reliable report, the master taxonomy table is updated to reflect the distribution change.

Prior to the 2022 data release, publication of species identifications were obfuscated to a higher taxonomic rank when the taxon was found to be listed as threatened, endangered, or sensitive at the state level where the observation was recorded. The state-level obfuscation routine was removed from the data publication process at all locations excluding sites located in D01 and D20, and data have been reprocessed to remove the obfuscation of state-listed taxa for all years. Federally listed threatened and endangered or sensitive species remain obfuscated at all sites and sensitive species remain redacted at National Park sites.

The full master taxonomy lists are available on the NEON Data Portal for browsing and download: <https://data.neonscience.org/taxonomic-lists>.

5 DATA QUALITY

5.1 Data Entry Constraint and Validation

Many quality control measures are implemented at the point of data entry within a mobile data entry application or web user interface (UI). For example, data formats are constrained and data values controlled through the provision of dropdown options, which reduces the number of processing steps necessary to prepare the raw data for publication. An additional set of constraints are implemented during the process of ingest into the NEON database. The product-specific data constraint and validation requirements built into data entry applications and database ingest are described in the document NEON Raw Data Validation for Coarse Downed Wood Log Survey (DPO.10010.001), provided with every download of this data product. Contained within this file is a field named 'entryValidationRulesForm', which describes syntactically the validation rules for each field built into the data entry application. Data entry constraints are described in NiCl syntax in the validation file provided with every data download, and the NiCl language is described in NEON's Ingest Conversion Language (NICL) specifications ([AD[15]]). Data collected prior to 2017 were processed using a paper-based workflow that did not implement the full suite of quality control features associated with the interactive digital workflow.

5.2 Automated Data Processing Steps

Following data entry into a mobile application or web user interface, the steps used to process the data through to publication on the NEON Data Portal are detailed in the NEON Algorithm Theoretical Basis Document: OS Generic Transitions (AD[14]).



5.3 Data Revision

All data are provisional until a numbered version is released. Annually, NEON releases a static version of all or almost all data products, annotated with digital object identifiers (DOIs). The first data Release was made in 2021. During the provisional period, QA/QC is an active process, as opposed to a discrete activity performed once, and records are updated on a rolling basis as a result of scheduled tests or feedback from data users. The Issue Log section of the data product landing page contains a history of major known errors and revisions.

5.4 Quality Flagging

The **dataQF** field in each data record is a quality flag for known errors applying to the record. Please see the *Special Considerations* section of this document for a list of known errors that may be present in the data, and below for an explanation of **dataQF** codes specific to this product.

Table 1: Descriptions of the dataQF codes for quality flagging

fieldName	value	definition
dataQF	legacyData	Data recorded using a paper-based workflow that did not implement the full suite of quality control features associated with the interactive digital workflow
dataQF	decayClass NotRecorded	Records collected with no decayClass; see <i>Special Considerations</i> section above for more information.

Records of land management activities, disturbances, and other incidents of ecological note that may have a potential impact are found in the Site Management and Event Reporting data product (DP1.10111.001)

6 REFERENCES

Affleck, D. L. R. 2008. A line intersect distance sampling strategy for downed wood inventory. *Canadian Journal of Forest Research* 38:2262-2273.

Affleck, D. L. R. 2010. On the efficiency of line intersect distance sampling. *Canadian Journal of Forest Research* 40:1086-1094.

Harmon, M. E., and J. Sexton. 1996. Guidelines for measurements of woody detritus in forest ecosystems. Publication No. 20 Long-term Ecological Research Network Office, University of Washington:1-42

Jordan, G. J., M. J. Ducey, and J. H. Gove. 2004. Comparing line-intersect, fixed-area, and point relascope sampling for dead and downed coarse woody material in a managed northern hardwood forest. *Canadian Journal of Forest Research* 34:1766-1775.



<i>Title:</i> NEON User Guide to Coarse Downed Wood Log Survey (DP1.10010.001)	<i>Date:</i> 04/11/2022
<i>Author:</i> Katherine Jones	<i>Revision:</i> C

Keller, M., M. Palace, G. P. Asner, R. Pereira, and J. N. M. Silva. 2004. Coarse woody debris in undisturbed and logged forests in the eastern Brazilian Amazon. *Global Change Biology* 10:784-795.

USDA, NRCS. 2014. The PLANTS Database (<http://plants.usda.gov>, 25 August 2014). National Plant Data Team, Greensboro, NC 27401-4901 USA.