



<i>Title:</i> NEON User Guide to Soil physical and chemical properties, periodic (DP1.10086.001)	<i>Date:</i> 03/02/2022
<i>Author:</i> Lee Stanish	<i>Revision:</i> C

NEON USER GUIDE TO SOIL PHYSICAL AND CHEMICAL PROPERTIES, PERIODIC (DP1.10086.001)

PREPARED BY	ORGANIZATION
Lee Stanish	TOS
Samantha Weintraub-Leff	TOS



<i>Title:</i> NEON User Guide to Soil physical and chemical properties, periodic (DP1.10086.001)	<i>Date:</i> 03/02/2022
<i>Author:</i> Lee Stanish	<i>Revision:</i> C

CHANGE RECORD

REVISION	DATE	DESCRIPTION OF CHANGE
A	04/20/2017	Initial Release
B	09/01/2020	Included statement about use of neonUtilities R package and possible sampling location changes. Added text on publication of soil chemistry and stable isotope data along with soil field collections and pH/moisture measurements as part of this data product. Sections 3 and 5: Updated reference for pH method; Section 3.2: Revised temporal design description to match current design; Section 3.3: Added Sampling Design Changes section and included changes to sampling frequency for microbial analyses; Section 3.5: Updated instructions on obtaining sample location data using the NEON API; Section 3.6: Introduced eventID; Section 3.8: Added information for missed sampling; Section 4.4: Added new dataQF option for early Alaska samples.
C	03/02/2022	Section 3.3: Added details for sample identifier changes that occurred in 2021 and 2022; minor text updates and corrections throughout; Section 4.3: Updated Data Revision with latest information regarding data release.



<i>Title:</i> NEON User Guide to Soil physical and chemical properties, periodic (DP1.10086.001)	<i>Date:</i> 03/02/2022
<i>Author:</i> Lee Stanish	<i>Revision:</i> C

TABLE OF CONTENTS

1	DESCRIPTION	1
1.1	Purpose	1
1.2	Scope	1
2	RELATED DOCUMENTS AND ACRONYMS	2
2.1	Associated Documents	2
3	DATA PRODUCT DESCRIPTION	3
3.1	Spatial Sampling Design	4
3.2	Temporal Sampling Design	5
3.3	Sampling Design Changes	8
3.4	Variables Reported	8
3.5	Spatial Resolution and Extent	9
3.6	Temporal Resolution and Extent	10
3.7	Associated Data Streams	10
3.8	Product Instances	10
3.8.1	Product Instances for Missed or Incomplete Sampling	11
3.9	Data Relationships	11
3.10	Special Considerations	12
4	DATA QUALITY	13
4.1	Data Entry Constraint and Validation	13
4.2	Automated Data Processing Steps	13
4.3	Data Revision	13
4.4	Quality Flagging	14
5	REFERENCES	14

LIST OF TABLES AND FIGURES

Table 1	Descriptions of the dataQF codes for quality flagging	14
---------	---	----



<i>Title:</i> NEON User Guide to Soil physical and chemical properties, periodic (DP1.10086.001)	<i>Date:</i> 03/02/2022
<i>Author:</i> Lee Stanish	<i>Revision:</i> C

Figure 1 Overview of soil field sampling and analysis workflow. 3

Figure 2 Representation of a NEON site with Tower and Distributed plots shown 4

Figure 3 Overview of soil field sampling and laboratory analyses based on bout type. At the highest level, within a year sampling bouts at a site are either Off-year or Coordinated. Off-year bouts (majority) perform routine measurements and collect various microbial samples for downstream analyses and archive. Coordinated bouts (every 5 years) perform routine measurements, microbial sample collections, and a full suite of biogeochemical analyses. 6

Figure 4 Soil field and laboratory workflows. Arrows indicate which data tables are generated by each step in the soil sampling process and under what conditions. Data ingest table names follow the convention: moduleID_tableName_in, for example sls_soilcorecollection_in 7

1 DESCRIPTION

1.1 Purpose

This document provides an overview of the data included in this NEON Level 1 data product, the quality controlled product generated from raw Level 0 data, and associated metadata. In the NEON data products framework, the raw data collected in the field - for example, soil temperature from a single collection event - are considered the lowest level (Level 0). Raw data that have been quality checked via the steps detailed herein, as well as simple metrics that emerge from the raw data are considered Level 1 data products.

The text herein provides a discussion of measurement theory and implementation, data product provenance, quality assurance and control methods used, and approximations and/or assumptions made during L1 data creation.

1.2 Scope

This document describes the steps needed to generate the L1 data product Soil physical and chemical properties, periodic, which encompasses field and laboratory measurements of soil temperature, moisture, and pH, the generation of subsamples used for microbial analyses, and measurements of soil carbon (C) and nitrogen (N) pools. As of August 2020, the Soil chemical properties (DP1.10078.001), Soil stable isotopes (DP1.10101.001), and Soil inorganic nitrogen pools and transformations (DP1.10080.001) data products are published as part of Soil physical and chemical properties, periodic (DP1.10086.001). However, this User Guide is focused on the subset of tables dealing with field collection, subsampling, and moisture/pH measurements, while two related Data Product User Guides detail the biogeochemical measurements included in this product. Data from the microbial subsamples can be found in the related data products listed below.

This document also provides details relevant to the publication of the data products via the NEON data portal, with additional detail available in the files NEON Data Variables for Soil physical and chemical properties, periodic (DP1.10086.001) (AD[05]) and NEON Categorical Codes for Soil physical and chemical properties, periodic (AD[06]), provided in the download package for this data product.

This document describes the process for ingesting and performing automated quality assurance and control procedures on the data collected in the field pertaining to TOS Protocol and Procedure: Soil Biogeochemical and Microbial Sampling (AD[09]), or TOS Standard Operating Procedure: Wetland Soil Sampling (AD[10]) if the site is a wetland. The raw data that are processed in this document are detailed in the file, NEON Raw Data Validation for Soil physical properties (Distributed periodic) (DPO.10086.001) (AD[04]), provided in the download package for this data product. Please note that raw data products (denoted by 'DPO') may not always have the same numbers (e.g., '10033') as the corresponding L1 data product.

2 RELATED DOCUMENTS AND ACRONYMS

2.1 Associated Documents

AD[01]	NEON.DOC.000001	NEON Observatory Design (NOD) Requirements
AD[02]	NEON.DOC.000913	TOS Science Design for Spatial Sampling
AD[03]	NEON.DOC.002652	NEON Data Products Catalog
AD[04]	Available with data download	Variables csv
AD[05]	Available with data download	Validation csv
AD[06]	Available with data download	Categorical Codes csv
AD[07]	NEON.DOC.000906	TOS Science Design for Terrestrial Biogeochemistry
AD[08]	NEON.DOC.000908	TOS Science Design for Terrestrial Microbial Diversity
AD[09]	NEON.DOC.014048	TOS Protocol and Procedure: Soil Biogeochemical and Microbial Sampling
AD[10]	NEON.DOC.004130	TOS Standard Operating Procedure: Wetland Soil Sampling
AD[11]	NEON.DOC.000008	NEON Acronym List
AD[12]	NEON.DOC.000243	NEON Glossary of Terms
AD[13]	NEON.DOC.004825	NEON Algorithm Theoretical Basis Document: OS Generic Transitions
AD[14]	Available on NEON data portal	NEON Ingest Conversion Language Function Library
AD[15]	Available on NEON data portal	NEON Ingest Conversion Language

3 DATA PRODUCT DESCRIPTION

The Soil physical and chemical properties, periodic data product is derived from TOS Protocol and Procedure: Soil Biogeochemical and Microbial Sampling (AD[09]), or TOS Standard Operating Procedure: Wetland Soil Sampling (AD[10]) if the site is a wetland. The sampling plan implements the guidelines and requirements described in the Science Designs for TOS Terrestrial Biogeochemistry (AD[07]) and Microbial Diversity (AD[08]). All accompanying field and laboratory data are reported at the spatial resolution of a single soil sampling location, e.g., an x,y coordinate (+/- 0.5 meters) within a subplot within a NEON plot, except for microbial metagenomic samples, which represent a pooled, plot-level composite and are thus reported at the scale of a NEON plot. For all samples, the temporal resolution is that of a single collection date.

Soils are sampled by horizon type (organic or mineral, Figure 1) to a maximum depth of 30 cm. Where possible, users can validate the technician designation of horizon type using measured C content in the soil carbon and nitrogen table (the USDA generally considers a horizon 'organic' if % organic C >= 20%). The type of device used to collect soils varies based on local soil types and is recorded for each sample (for details about site-specific sampling devices, refer to the soil sampling protocol, NEON.DOC.014048). At each x,y sampling location, in-situ soil temperature is measured with a hand-held probe and soil samples are removed and homogenized. Subsamples for microbial community analyses are immediately frozen on dry ice.




Figure 1: Overview of soil field sampling and analysis workflow.

Following field sampling, gravimetric soil moisture and pH measurements are conducted at NEON domain support facilities, following the methods outlined in Robertson et al. (1999) and Burt (2014), respectively. Prior to 2020, NEON measured soil pH using the method outlined in Robertson et al. (1999). For select bouts, subsamples are also prepared for total C and N and inorganic N measurements.

Data on soil physical and chemical properties help to elucidate the constraints on microbial activity, nutrient cycling, and carbon storage in soils at the plot, site, and continental scales. They also provide essential data for understanding change in soil microbial and biogeochemical dynamics over time.

3.1 Spatial Sampling Design

Soil biogeochemical and microbial sampling is executed at all terrestrial NEON sites. Soils are sampled from three pre-determined, randomly assigned x,y locations in 3 of 4 randomly selected subplots per 40 x 40 meter plot (Figure 1). If large rocks, tree roots, animal burrows, or other significant impediments are encountered at a pre-determined sampling location, technicians proceed down a list of random, alternate locations until an acceptable one is found. Ten plots per site are sampled, four within the tower airshed (Figure 2) and six others distributed across the landscape, located in dominant vegetation types. Tower plots are selected using a random spatial design, while distributed plots follow a random stratified design based on National Land Cover Database (NLCD) vegetation class. The number of distributed plots within each NLCD class are proportional to the percent coverage of that class. See AD[02] for further details on the NEON spatial design.

As much as possible, sampling occurs in the same locations over the lifetime of the Observatory. However, over time some sampling locations may become impossible to sample, due to disturbance or other local changes. When this occurs, the location and its location ID are retired. A location may also shift to slightly different coordinates. Refer to the locations endpoint of the NEON API for details about locations that have been moved or retired: <https://data.neonscience.org/data-api/endpoints/locations/>




Figure 2: Representation of a NEON site with Tower and Distributed plots shown

<i>Title:</i> NEON User Guide to Soil physical and chemical properties, periodic (DP1.10086.001)	<i>Date:</i> 03/02/2022
<i>Author:</i> Lee Stanish	<i>Revision:</i> C

3.2 Temporal Sampling Design

Soil sampling for a suite of physical, chemical and microbial analyses occurs up to three times per year on either an annual or interannual time frame, the frequency of collection varying by analysis type (Figure 3). A suite of physical measurements are made at all sites during every sampling bout which consists of basic field measurements, soil moisture, pH, and subsampling for microbial genetic archive. At all sites, sampling occurs during the period of peak greenness/productivity as indicated either by remotely sensed vegetation indices (e.g. NDVI) or historic changes in precipitation. At most sites, soil sampling occurs two additional times per year, usually bracketing the peak greenness window and aimed at capturing seasonal transitions in microbial activity. Sampling frequency is reduced in arctic and boreal sites, which are only sampled during peak greenness.

	Off-Year			Coordinated bout			
N-trans Bout Type	No			T initial			T final
Sample Timing	T1	PG	T2	T1	PG	T2	T1, PG, T2
Bout Type	microbes/ microbes- Biomass	microbes/ microbes- Biomass	microbes/ microbes- Biomass	microbes- Biomass	microbes- BiomassBGC	microbes- Biomass	fieldOnly
Field-generated Samples	Bulk -gen* -gaX	Bulk -gaX -gen* -comp*	Bulk -gen* -gaX	Bulk -gen -gaX	Bulk -gen -gaX -comp	Bulk -gen -gaX	Bulk
Lab-generated Samples	-bm*	-bm*	-bm*	-bm -kcl	-bm; -kcl -cn; -ba	-bm -kcl	kcl
Lab measurements	pH moisture	pH moisture	pH moisture	pH moisture	pH moisture	pH moisture	moisture
* Core sites only							
Abbreviations							
Sample Timing:							
T1: Transition 1							
PG: Peak Greenness							
T2: Transition 2							
Sample:							
Bulk: Homogenized soil used for all subsamples and analyses							
Subsamples:							
-gen: microbial genetic analysis subsample							
-gaX: microbial genetic archive subsample, X denotes subsample number of 1-5 (for up to 5 vials)							
-comp: plot-level composited microbial metagenomics subsample							
-bm: microbial biomass subsample							
-kcl: KCl extraction sample for nitrogen transformation rate measurement							
-cn: BGC analysis subsample							
-ba: BGC archive subsample							

Figure 3: Overview of soil field sampling and laboratory analyses based on bout type. At the highest level, within a year sampling bouts at a site are either Off-year or Coordinated. Off-year bouts (majority) perform routine measurements and collect various microbial samples for downstream analyses and archive. Coordinated bouts (every 5 years) perform routine measurements, microbial sample collections, and a full suite of biogeochemical analyses.

Once every five years each NEON site will conduct Coordinated sampling in which soils collected during routine sampling bouts for that year are utilized for additional downstream analyses. These analyses include incubation and processing of soil cores to determine inorganic N pools and net transformation rates, plus total soil organic C and total N measurements and creation of an air-dried subsample for the Biorepository (during the peak greenness bout only, Figure 4). See the associated Data Product



User Guides for more details on these biogeochemistry measurements. The sites rotate on a pre-defined schedule such that every site will complete Coordinated sampling over a 5-year period.




Figure 4: Soil field and laboratory workflows. Arrows indicate which data tables are generated by each step in the soil sampling process and under what conditions. Data ingest table names follow the convention: moduleID_tableName_in, for example sls_soilcorecollection_in

A plot-level, pooled metagenomics sample is also collected during peak greenness. At core terrestrial sites, this sample is generated every year, while gradient sites generate a metagenomics sample once every five years during Coordinated bouts.

Subsamples for additional microbial analyses are also collected during routine bouts according to Figure 3. Subsamples are collected for microbial marker gene sequencing (16S and ITS sequencing) and microbial biomass analyses during all bouts at core terrestrial sites, while gradient sites generate subsamples for these analyses once every five years during Coordinated sampling.

For each of the five data tables discussed in this User Guide,

- the field collection, pH, and moisture tables will contain data for 1-3 sampling events per site per year,
- the metagenomics pooling table will contain data for 1 sampling event per core site per year as well as for gradient sites conducting Coordinated sampling,
- the biogeochemistry subsampling table will contain data for 1 sampling event per site, every 5 years.

3.3 Sampling Design Changes

Over the course of early operations, the design for soil periodic sampling evolved. Below is a list of previous sampling strategies that differ from the current design, with applicable years indicated.

- 2013 - 2018: All terrestrial sites generated a metagenomics sample annually.
- 2013 - 2018: Subsamples were collected for microbial marker gene sequencing analyses (16S and ITS sequencing) during every bout and for all sites.
- 2013 - 2018: Subsamples were collected for microbial group abundances analysis during every bout and for all sites.
- 2015 - 2017: The field methods for distinguishing litter from organic soil were deemed not optimal for arctic tundra and boreal sites, and the method was refined in 2018. See Special Considerations for details.
- 2017 - 2019: Subsamples for microbial biomass analysis were collected once every five years only, during Coordinated sampling bouts.

Additionally, there have been changes to the way samples are uniquely identified.

- Pre-2021: Genetic archive samples had identifiers linked to the descriptive sampleID (ex: CPER_001-M-1-2.5-20200415-ga1). From mid-2021 onward they are identified by scannable barcodes only (ex: A00000234432).
- Pre-2022: The **sampleID** field contained the sampled x,y coordinate (ex: WREF_001-O-35-4.5-20200715). From 2022 onward, it contains the subplot instead (ex: WREF_001-O-23-20200715).

3.4 Variables Reported

All variables reported from the field or laboratory technician (L0 data) are listed in the file, NEON Raw Data Validation for Soil physical properties (Distributed periodic) (DP0.10086.001) (AD[04]). All variables reported in the published data (L1 data) are also provided separately in the file, NEON Data Variables for Soil physical and chemical properties, periodic (DP1.10086.001) (AD[05]).

Field names have been standardized with Darwin Core terms (<http://rs.tdwg.org/dwc/>; accessed 16 February 2014), the Global Biodiversity Information Facility vocabularies (<http://rs.gbif.org/vocabulary/gbif/>; accessed 16 February 2014), and the VegCore data dictionary (<https://projects.nceas.ucsb.edu/nceas/projects/bien/wiki/VegCore>; accessed 16 February 2014), where applicable. NEON TOS spatial data

employs the World Geodetic System 1984 (WGS84) for its fundamental reference datum and GEOID09 for its reference gravitational ellipsoid. Latitudes and longitudes are denoted in decimal notation to six decimal places, with longitudes indicated as negative west of the Greenwich meridian.

Some variables described in this document may be for NEON internal use only and will not appear in downloaded data. These are indicated with downloadPkg = “none” in NEON Data Variables for Soil physical and chemical properties, periodic (DP1.10086.001) (AD[05]).

3.5 Spatial Resolution and Extent

The finest resolution at which spatial data are reported is a single x,y sampling location.

sampleID (unique ID given to the individual soil sampling location and horizon) → **subplotID** (ID of subplot within plot) → **plotID** (ID of plot within site) → **siteID** (ID of NEON site) → **domainID** (ID of NEON domain).

The basic spatial data included in the data downloaded include spatial location (northing and easting) and elevation of the *centroid* of the plot where sampling occurred, plus associated uncertainty due to GPS error and plot width. Shapefiles of all NEON Terrestrial Observation System sampling locations can be found in the Document Library: <http://data.neonscience.org/documents>.

To derive a more precise estimate of the location of each soil sampling location, there are two options:

- Use the getLocTOS function from the geoNEON package, available here: <https://github.com/NEONScience/NEON-geolocation>
- Or follow these steps to perform the same calculation:
 1. Find the namedLocation field in the data; this is the named location of the plot. For example, ‘HARV_052.basePlot.bgc’ is a complete named location for plotID ‘HARV_052’.
 2. Use the API (<http://data.neonscience.org/api>; e.g. http://data.neonscience.org/api/v0/locations/HARV_052.basePlot.bgc?history=true) to query for easting(“locationUtmEasting”), northing(“locationUtmNorthing”), and coordinateUncertainty (“Value for Coordinate uncertainty”) that match the sampling date. These are the location data from the plot centroid, and will be used in the next steps.
 3. Calculate the actual northing of the x,y location. This can be accomplished by subtracting 20 meters from locationUtmNorthing, then adding the y-coordinate value. Use the following formula:

$$yCoordNorthing = locationUtmNorthing - 20m + coreCoordinateY$$

4. Calculate the actual easting of the x,y location. This can be accomplished by subtracting 20 meters from locationUtmEasting, then adding the x-coordinate value. Use the following formula:

$$xCoordEasting = locationUtmEasting - 20m + coreCoordinateX$$

5. Increase coordinateUncertainty by an appropriate amount to account for variance in sampling location area (+/- 0.5 m), plus the error introduced by technicians navigating within plots, stretching meter tapes to navigate to x,y locations, etc. This uncertainty, on average, will be +/- 1 m, but will vary by site based on terrain heterogeneity and density of vegetation.

3.6 Temporal Resolution and Extent

The finest resolution at which temporal data are reported is collectDate. All samples associated with a sampling event have collectDates within a ~14-day window and are categorized by the target season or seasonal transition (e.g. wet-dry transition, winter-spring transition, peak greenness). The total number of sampling events per year will vary among sites, based on the length of the growing season. It is expected that 3 sampling events will occur annually for microbial sampling in most sites, except those in arctic and boreal regions.

In 2020 the **eventID** field was added in order to facilitate grouping of samples collected during the same sampling event or bout. The eventID is created by merging the following fields:

siteID + . + sampleTiming + . + collectDate (year portion only)

Example. WOOD.peakGreenness.2020

For the handful of sites where sampling seasons straddle 2 calendar years (for example, Transition 1 in Oct, Peak Green in Jan, Transition 2 in March), the eventID contains both years separated by a dash.

Example. LAJA.peakGreenness.2021-2022

3.7 Associated Data Streams

sampleID is the linking variable that can be used to join field metadata, moisture/pH, and soil chemistry data across tables. In addition, field and laboratory data from the Soil physical and chemical properties, periodic data product will be necessary to interpret and utilize several related soil microbial data products. For Soil microbe community composition (DP1.10081.001), Soil microbe group abundances (DP1.10109.001), and Soil microbe marker gene sequences (DP1.10108.001) **geneticSampleID** is the variable name that links samples and their associated metadata from field to external laboratory data tables. For Soil microbe biomass (DP1.10104.001), this variable is **biomassID**, while for Soil microbe metagenome sequences (DP1.10107.001), the linking variable name is **compositeSampleID**.

3.8 Product Instances

Soil samples are collected at all terrestrial NEON sites. A maximum of 10 plots are sampled at every site at a frequency of 3 times per year (except for arctic and boreal sites, which are sampled once). For each

soil horizon (maximum of 2, organic or mineral), 3 samples per plot are collected, 1 in each of 3 randomly selected subplots. When organic and mineral horizons are present at a sampling location, both horizons are sampled only during Coordinated bouts, else only the top horizon is sampled.

Off-years: At most sites, expect soil sampling will result in 90 unique soil samples per site per year analyzed for moisture and pH. At boreal/arctic sites, expect 30 unique soil samples per site per year. Each unique sample may result in 0-8 sub-samples used for additional analyses and archiving.

Coordinated years: At most sites, expect soil sampling will result in 90-180 unique soil samples per site per year, analyzed for pH, moisture, and chemistry, and another 90-180 analyzed for net N transformation rates (incubated cores). At boreal/arctic sites, expect 30-60 unique soil samples analyzed for pH, moisture, and chemistry and another 30-60 analyzed for net N transformation rates (incubated cores). Each unique sample may result in 0-11 sub-samples used for additional analyses and archiving.

3.8.1 Product Instances for Missed or Incomplete Sampling

Beginning in 2020, missed or incomplete sampling bouts are recorded using the **samplingImpractical** field located in the *sls_soilCoreCollection* data table. Any value for **samplingImpractical** that is not 'OK' indicates that sampling did not occur. A record will be created for every expected sampling location that was unable to be sampled. A **sampleID** will be generated for these records that consists of the following:

- plotID + "-" + subplotID + "-" + collectDate (intended) + "-" + samplingImpractical value
- Example. "NIWO_013-23-20200501-locationFrozen"

The same number of records as outlined above are expected in *sls_soilCoreCollection* for an impractical sampling event. However, records in any downstream tables (e.g. *sls_Moisture*, *sls_pH*, *sls_bgcSubsampling*, *sls_metagenomicsPooling*, all chemistry data tables) are not generated.

3.9 Data Relationships

The protocol dictates that each x,y location within a randomly selected subplot yields a unique sampleID per horizon per **collectDate** (day of year, local time) in *sls_soilCoreCollection*. A record from *sls_soilCoreCollection* may have zero or one child records in *sls_soilpH* and *sls_soilMoisture*; a given *sls_soilCoreCollection.sampleID* is expected to be sampled only once. Depending on the type of bout and time of year (Figure 3), a record from *sls_soilCoreCollection* may have zero or one child records in *sls_metagenomicsPooling* and in *sls_bgcSubsampling*. Records for soil subsamples used in chemistry measurements will appear in associated downstream tables, described in the two related Data Product User Guides. Duplicates and/or missing data may exist where protocol and/or data entry aberrations have occurred; *users should check data carefully for anomalies before joining tables.*

sls_soilCoreCollection.csv -> One record expected per **sampleID**. Depending upon **boutType**, each **sampleID** generates up to a single **geneticSampleID**, up to 5 interchangeable **geneticArchiveSamples**, and up to a single **biomassID**. Generates samples used in Soil microbe community composition (DP1.10081.001), Soil microbe group abundances (DP1.10109.001), Soil microbe marker gene sequences (DP1.10108.001),

and Soil microbe biomass (DP1.10104.001). Additionally, subsamples generated from soil **sampleIDs** are used to measure soil inorganic N pools and transformations

sls_soilpH.csv -> One record expected per **sampleID**, generates a single **pHSampleID**.

sls_soilMoisture.csv -> One record expected per **sampleID**, generates a single **moistureSampleID**. Missing records in this table indicate failure to carry out the measurement.

sls_metagenomicsPooling.csv -> One record expected per **plotID** per **horizon** per **collectDate** (day of year, local time). Record represents a mixture of the samples collected in a plot (listed in **toCompositeSampleIDList**). Each record generates a single **compositeSampleID**, used in Soil microbe metagenome sequences (DP1.10107.001).

sls_bgcSubsampling.csv -> One record expected per **sampleID**, generates a single **cnSampleID** and **bgcArchiveID** used for chemistry measurements and air-dried archive respectively. Chemistry values are reported in a separate carbon-nitrogen table.

sampleIDs will be generated for each unique physical soil sample created during a collection event, and **sampleBarcodes** may also be generated. Each **sampleID** will yield between one to five frozen archive subsamples, stored in 5 mL cryovials at -80C or liquid nitrogen. When a collection event includes soil chemistry and isotope analyses, an air-dried archive subsample (**bgcArchiveID**) will also be created with an associated **bgcArchiveMass**. If not, following pH and moisture measurements, any remaining soil material will be discarded at the end of the calendar year.

Data downloaded from the NEON Data Portal are provided in separate data files for each site and month requested. The neonUtilities R package contains functions to merge these files across sites and months into a single file for each table described above. The neonUtilities package is available from the Comprehensive R Archive Network (CRAN; <https://cran.r-project.org/web/packages/neonUtilities/index.html>) and can be installed using the install.packages() function in R. For instructions on using neonUtilities to merge NEON data files, see the Download and Explore NEON Data tutorial on the NEON website: <https://www.neonscience.org/download-explore-neon-data>

3.10 Special Considerations

1. Every five years, incubation and processing of soil cores occurs in order to measure net nitrogen transformation rates. Incubated cores, designated with **nTransBoutType = tfinal**, will appear intermingled with non-incubated samples throughout the sls_soilCoreCollection and sls_soilMoisture data tables. However, they will not be measured for soil pH and will not be subsampled for microbial diversity or bulk soil chemistry and isotopic analyses. Refer to the related inorganic N Data Product User Guide for more details.
2. *Wetland sites*: At sites in which the Wetland SOP is carried out, standing water may be encountered during sampling. Soil temperature is not measured for locations where **standingWaterDepth** > 0.
3. *Alaska-specific consideration*: Beginning in 2018 the process for differentiating between organic soil and litter in arctic tundra and boreal sites was revised to more accurately reflect the slow rates of organic matter decomposition and the extensive fibric organic material at the TOOL, BARR, DEJU, BONA, and HEAL sites. Prior to 2018, instructions to technicians may have resulted in a portion of

the top-most fibric material being included in measurements of litter depth, and excluded from O-horizon soil samples. From 2018 onward, the Protocol (AD[09]) dictates that all non-green, friable fibric material that has roots growing in it should be classified as organic soil and sampled as such, rather than being measured as litter. Users should use caution when comparing soil data from these five sites collected before and after 2018.

4 DATA QUALITY

4.1 Data Entry Constraint and Validation

Many quality control measures are implemented at the point of data entry within a mobile data entry application or web user interface (UI). For example, data formats are constrained and data values controlled through the provision of dropdown options, which reduces the number of processing steps necessary to prepare the raw data for publication. An additional set of constraints are implemented during the process of ingest into the NEON database. The product-specific data constraint and validation requirements built into data entry applications and database ingest are described in the document NEON Raw Data Validation for Soil physical properties (Distributed periodic) (DP0.10086.001), provided with every download of this data product. Contained within this file is a field named 'entryValidationRulesForm', which describes syntactically the validation rules for each field built into the data entry application. Data entry constraints are described using a standardized data validation language (NiCl) internal to NEON. Please see AD[14] and AD[15] for more information about the NiCl language.

Data collected prior to 2017 were processed using a paper-based workflow that did not implement the full suite of quality control features associated with the interactive digital workflow.

4.2 Automated Data Processing Steps

Following data entry into a mobile application or web user interface, the steps used to process the data through to publication on the NEON Data Portal are detailed in the NEON Algorithm Theoretical Basis Document: OS Generic Transitions (AD[13]). Additionally, the methods used to create calculated fields from the raw data (soil moisture, water-to-solution ratios for pH measurements) are detailed in NEON Raw Data Validation for Soil physical properties (Distributed periodic) (DP0.10086.001).

4.3 Data Revision

All data are provisional until a numbered version is released. Annually, NEON releases a static version of all or almost all data products, annotated with digital object identifiers (DOIs). The first data Release was made in 2021. During the provisional period, QA/QC is an active process, as opposed to a discrete activity performed once, and records are updated on a rolling basis as a result of scheduled tests or feedback from data users. The Issue Log section of the data product landing page contains a history of major known errors and revisions.

4.4 Quality Flagging

The **dataQF** field in each data record is a catch-all quality flag for known errors applying to the record. The dataQF codes specific to the five soil tables discussed in this User Guide are detailed below. The issue with Alaska soils is detailed further in the *Special Considerations* section above.

Table 1: Descriptions of the dataQF codes for quality flagging

fieldName	value	definition
dataQF	legacyData	Data recorded using a paper-based workflow that did not implement the full suite of quality control features associated with the interactive digital workflow
dataQF	alaskaDeprecatedMethod	Different methods used for measuring litter depth and the boundaries between soil horizons prior to 2018, use caution when comparing measurements to data collected in 2018 and later

Additionally, several other condition and quality fields have been added to the field collection and lab processing tables over time in order to communicate anomalous sampling conditions or method deviations. These include variables such as **horizonDetails**, **biophysicalCriteria**, **sampleCondition**, and others. Definitions for the categorical codes used for these fields are included in the file NEON Categorical Codes for Soil physical and chemical properties, periodic (AD[06]), provided in the download package for this data product. Fields have been added over time and entries may be missing in older data.

Records of land management activities, disturbances, and other incidents of ecological note that may have a potential impact are found in the Site Management and Event Reporting data product (DP1.10111.001).

5 REFERENCES

Burt, R. (2014). Kellogg Soil Survey Laboratory Methods Manual. Report No. 42 Version 5.0. Soil Survey Laboratory Investigations. United States Department of Agriculture, Natural Resources Conservation Service. 1031 pp.

Robertson, G. P. (1999). Standard soil methods for long-term ecological research (Vol. 2). Oxford University Press on Demand.