

NEON USER GUIDE FOR AQUATIC MICROBE CELL COUNT SAMPLING (NEON.DP1.20138)

PREPARED BY	ORGANIZATION	DATE
Stephanie Parker	AQU	06/12/2017
Caren Scott	AQU	06/12/2017

<i>Title:</i> NEON User Guide for Aquatic Microbe Cell Count Sampling (NEON.DP1.20138)	<i>Date:</i> 06/12/2017
<i>Author:</i> Stephanie Parker	<i>Revision:</i> A

CHANGE RECORD

REVISION	DATE	DESCRIPTION OF CHANGE
A	06/21/2017	Initial Release

TABLE OF CONTENTS

1	DESCRIPTION	1
1.1	Purpose	1
1.2	Scope	1
2	RELATED DOCUMENTS AND ACRONYMS	2
2.1	Associated Documents	2
2.2	Acronyms	2
3	DATA PRODUCT DESCRIPTION	3
3.1	Spatial Sampling Design	3
3.2	Temporal Sampling Design	5
3.3	Variables Reported	5
3.4	Spatial Resolution and Extent	5
3.5	Temporal Resolution and Extent	5
3.6	Associated Data Streams	6
3.7	Product Instances	6
3.8	Data Relationships	6
3.9	Special Considerations	6
4	DATA ENTRY CONSTRAINT AND VALIDATION	7
5	DATA PROCESSING STEPS	7
6	REFERENCES	7

LIST OF TABLES AND FIGURES

Figure 1	Generic aquatic site layouts with surface water cell count sampling locations highlighted in red boxes for wadeable streams, lakes, and rivers.	4
Figure 2	Lake and river sampling depths in non-stratified and stratified water columns.	4
Figure 3	Schematic of the applications used by field technicians to enter water chemistry and aquatic cell count field data	8

1 DESCRIPTION

1.1 Purpose

This document provides an overview of the data included in this NEON Level 1 data product, the quality controlled product generated from raw Level 0 data, and associated metadata. In the NEON data products framework, the raw data collected in the field, for example, the dry weights of litter functional groups from a single collection event are considered the lowest level (Level 0). Raw data that have been quality checked via the steps detailed herein, as well as simple metrics that emerge from the raw data are considered Level 1 data products.

The text herein provides a discussion of measurement theory and implementation, data product provenance, quality assurance and control methods used, and approximations and/or assumptions made during L1 data creation.

1.2 Scope

This document describes the steps needed to generate the L1 data product Surface water cell count - the count of bacterial cells per liter of surface water in streams, rivers, and lakes - and associated field and external lab metadata. This document also provides details relevant to the publication of the data products via the NEON data portal, with additional detail available in the file, NEON Data Variables for Aquatic Microbe Cell Counts (NEON.DP1.20138) (AD[05]), provided in the download package for this data product.

This document describes the process for ingesting and performing automated quality assurance and control procedures on the data collected in the field pertaining to AOS Protocol and Procedure: Aquatic Microbial Sampling (AD[06]). The raw data that are processed in this document are detailed in the file, NEON Raw Data Validation for Aquatic Microbe Cell Counts (NEON.DP0.20138) (AD[04]), provided in the download package for this data product. Please note that raw data products (denoted by 'DP0') may not always have the same numbers (e.g., '20138') as the corresponding L1 data product.

2 RELATED DOCUMENTS AND ACRONYMS

2.1 Associated Documents

AD[01]	NEON.DOC.000001	NEON Observatory Design (NOD) Requirements
AD[02]	NEON.DOC.001152	NEON Aquatic Sampling Strategy
AD[03]	NEON.DOC.002652	NEON Level 1, Level 2 and Level 3 Data Products Catalog
AD[04]	NEON.DP0.20138.001_dataValidation.csv	NEON Raw Data Validation for Aquatic Microbe Cell Counts (NEON.DP0.20138)
AD[05]	NEON.DP1.20138.001_variables.csv	NEON Data Variables for Aquatic Microbe Cell Counts (NEON.DP1.20138)
AD[06]	NEON.DOC.003041	AOS Protocol and Procedure: Aquatic Microbial Sampling
AD[07]	NEON.DOC.000008	NEON Acronym List
AD[08]	NEON.DOC.000243	NEON Glossary of Terms
AD[09]	NEON.DOC.002905	AOS Protocol and Procedure: Water Chemistry Sampling in Surface Waters and Groundwater
AD[10]	OS_Generic_Transitions.pdf	NEON Algorithm Theoretical Basis Document: OS Generic Transitions
AD[11]		NEON's Ingest Conversion Language (NICL) specifications

2.2 Acronyms

Acronym	Definition
PI	Propidium iodide

<i>Title:</i> NEON User Guide for Aquatic Microbe Cell Count Sampling (NEON.DP1.20138)	<i>Date:</i> 06/12/2017
<i>Author:</i> Stephanie Parker	<i>Revision:</i> A

3 DATA PRODUCT DESCRIPTION

This data product contains the quality-controlled, field sampling metadata and associated total bacterial cell count data provided by a contracted lab. Field samples are collected in the water column of wadeable streams, rivers, and lakes in conjunction with surface water chemistry samples. Cell count samples are preserved with formaldehyde in the field, kept in the dark at 4°C, and shipped to an external lab for analysis.

Surface water microbes are collected 12 times per year in wadeable streams and 6 times per year in lakes and non-wadeable streams, at the same time and location as standard recurrent (monthly) water chemistry samples (AD[09]). Details on sampling locations and timing are provided in NEON Raw Data Validation for Aquatic Microbe Cell Counts (NEON.DP0.20138) (AD[06]) and the Surface Water Chemistry Sampling in Aquatic Habitats protocol (AD[09]). Cell count samples are collected as grab samples from the water column at sampling locations near the S2 sensor set in streams, and near the buoy in rivers, and near buoy, inlet, and outlet sensor sets in lakes. In lakes and rivers with a stratified water column, samples are collected at multiple depths.

An 18 mL aliquot of field sample is preserved with 2 mL of 10% formaldehyde (final concentration of ~1% formaldehyde) in the field. Samples are sent to an external analytical lab within 60 days of collection and analyzed using a propidium iodide (PI) staining method and epifluorescence microscopy (Boulos et al. 1999). Cell counts are enumerated using an image analysis program calibrated for NEON samples. Additional quality assurance data related to the automated counts, including counts of a standard reference photo, are available in the expanded package.

3.1 Spatial Sampling Design

Aquatic microbial cell count samples are collected at all NEON aquatic sites at the same time and location as surface water chemistry samples (AD[09]). At stream sites, 1 sample is collected from the thalweg <1 m downstream of the downstream sensor set (S2) on each sampling date. Samples represent the water column, so care is taken to avoid stirring up sediments that may contaminate the sample.

At river (non-wadeable stream) sites, cell count samples are collected just downstream of the sensor set or profiling buoy (station = 'c0') (Figure 1). If the river is non-stratified, samples are collected at 0.5 m depth. If the river is stratified, an epilimnion sample is collected at 0.5 m (station = 'c1') and an integrated sample is collected from the hypolimnion (station = 'c2'). Care is taken to avoid contamination from sediments suspended by the boat motor or anchor.

At lake sites, samples are collected near the profiling buoy, the inlet sensor, and outlet sensor (Figure 2). Near the buoy, sampling depth is dependent on the presence or absence of lake stratification. In an unstratified lake, the sample is collected near the surface at 0.5 m depth. In a stratified lake, additional samples are collected from the hypolimnion, in addition to the surface water sample. In lakes with a shallow hypolimnion (<4 m), the sample is collected from the midpoint of the hypolimnion. In lakes with a deeper hypolimnion (>4 m), an integrated sample is collected throughout the hypolimnion. Samples collected near the inlet and outlet sensor sets are collected near the surface at 0.5 m depth. See AOS Protocol and Procedure: Aquatic Microbial Sampling (AD[06]) and AOS Protocol and Procedure: Water Chemistry Sampling in Surface Waters and Groundwater (AD[09]) for additional details.




Figure 1: Generic aquatic site layouts with surface water cell count sampling locations highlighted in red boxes for wadeable streams, lakes, and rivers.




Figure 2: Lake and river sampling depths in non-stratified and stratified water columns.

Title: NEON User Guide for Aquatic Microbe Cell Count Sampling (NEON.DP1.20138)	Date: 06/12/2017
Author: Stephanie Parker	Revision: A

3.2 Temporal Sampling Design

Surface water microbe cell count samples are collected at the same time and location as standard recurrent water chemistry sampling: once per month in wadeable streams (12 times per year). At streams sites, samples are collected year-round including when the stream is frozen over if the ice can be broken by hand. When the ice becomes too thick, sampling is suspended and noted as **samplingImpractical** in the field data. At lake and river sites, microbe cell count samples are collected every other month with standard recurrent water chemistry samples. At northern sites, samples are collected year round and collected under the ice during winter. See NEON Aquatic Sampling Strategy (AD[02]), AOS Protocol and Procedure: Aquatic Microbial Sampling (AD[06]) and AOS Protocol and Procedure: Water Chemistry Sampling in Surface Waters and Groundwater (AD[09]) for additional details.

3.3 Variables Reported

All variables reported from the field or laboratory technician (L0 data) are listed in the file, NEON Raw Data Validation for Aquatic Microbe Cell Counts (NEON.DP0.20138) (AD[04]). All variables reported in the published data (L1 data) are also provided separately in the file, NEON Data Variables for Aquatic Microbe Cell Counts (NEON.DP1.20138) (AD[05]).

Field names have been standardized with Darwin Core terms (<http://rs.tdwg.org/dwc/>; accessed 16 February 2014), the Global Biodiversity Information Facility vocabularies (<http://rs.gbif.org/vocabulary/gbif/>; accessed 16 February 2014), the VegCore data dictionary (<https://projects.nceas.ucsb.edu/nceas/projects/bien/wiki/VegCore>; accessed 16 February 2014), where applicable. NEON OS spatial data employs the World Geodetic System 1984 (WGS84) for its fundamental reference datum and Earth Gravitational Model 96 (EGM96) for its reference gravitational ellipsoid. Latitudes and longitudes are denoted in decimal notation to six decimal places, with longitudes indicated as negative west of the Greenwich meridian.

Some variables described in this document may be for NEON internal use only and will not appear in downloaded data.

3.4 Spatial Resolution and Extent

The finest resolution at which spatial data are reported is at a single station within a site. For example, data may be collected at a specific depth in the water column of a lake. The basic spatial data included in the data downloaded include the latitude, longitude, and elevation of the named location at the aquatic site (e.g., the aquatic location) or the latitude and longitude of an alternate location if the named location is not suitable for sampling.

namedLocation (unique ID given to the location within the site) → **siteID** (ID of NEON site) → **domainID** (ID of a NEON domain)

3.5 Temporal Resolution and Extent

The finest resolution at which temporal data are reported is at **collectDate**, the date and time of day when the sample was collected in the field.

<i>Title:</i> NEON User Guide for Aquatic Microbe Cell Count Sampling (NEON.DP1.20138)	<i>Date:</i> 06/12/2017
<i>Author:</i> Stephanie Parker	<i>Revision:</i> A

3.6 Associated Data Streams

Cell count samples are related to water chemistry samples collected at the same time and location. Water chemistry data are available in the 'Chemical properties of surface water' data product (NEON.DP1.20093).

Cell count samples are also related to aquatic microbe sequencing data generated from subsamples of the same parent sample (linked with **parentSampleID**), including Surface water microbe community composition (NEON.DP1.20141), Surface water microbe group abundances (NEON.DP1.20278), Surface water microbe marker gene sequences (NEON.DP1.20282), and Surface water microbe metagenome sequences (NEON.DP1.20281).

3.7 Product Instances

At each stream site, there will be 12 samples collected per year. At a lake or river site, there will be a minimum of 6 samples and a maximum of 9 samples collected per year (maximum if water the column is stratified). Each sample generates one cell count record at the external lab.

3.8 Data Relationships

A record in `amc_fieldCellCounts` must have a corresponding record in `amc_fieldSuperParent` describing measurement depth and abiotic variables during sample collection. Each record in `amc_fieldCellCounts` may be linked to a record in `amc_cellCounts`, which contains data from the external laboratory. Duplicates and/or missing data may exist where protocol and/or data entry aberrations have occurred; users should check data carefully for anomalies before joining tables.

`amc_fieldSuperParent.csv` -> One record is created for each cell count that comes from a surface water sample, and contains metadata that applies to cell counts.

`amc_fieldCellCounts.csv` -> One record is created by field personnel for each cell count sample. Each field record has a corresponding `amc_fieldSuperParent` **parentSampleID**. The field **cellCountSampleID** is also created here and will be used to track the sample through to the external lab.

`amc_cellCounts.csv` -> One record is created by the external lab for each cell count sample, linked to the field data by the field **cellCountSampleID**. If samples need to be re-analyzed for QA reasons, there may be more than 1 record per **cellCountSampleID**.

`amc_cellCountLabSummary.csv` -> QA data are recorded using a **labSpecificStartDate** and **labSpecificEndDate**. The QA data pertain to the lab analysis if the **testedDate** in `amc_cellCounts` falls between this date range.

3.9 Special Considerations

The cell count data results comes from an external lab, in the field **rawMicrobialAbundance**. **rawMicrobialAbundance** is NOT corrected for preservative volume, so data users will need to apply this correction using data from **cellCountPreservantVolume** from the `amc_fieldCellCounts` for an accurate cell count value.

Title: NEON User Guide for Aquatic Microbe Cell Count Sampling (NEON.DP1.20138)	Date: 06/12/2017
Author: Stephanie Parker	Revision: A

$$\frac{\text{microbialAbundancePerMl}_i = \text{amc_cellCountsrawMicrobialAbundance}_i}{\text{amc_fieldCellCounts_cellCountSampleVolume}_i + \text{amc_fieldCellCounts_cellCountPreservantVolume}_i} \text{amc_fieldCellCounts_cellCountSampleVolume}_i \quad (1)$$

Where 'i' is a unique **cellCountSampleID**

See the external lab SOP (referenced in amc_cellCounts.csv) for calculations applied to the data by the external laboratory.

4 DATA ENTRY CONSTRAINT AND VALIDATION

Many quality control measures are implemented at the point of data entry within a mobile data entry application or web user interface (UI). For example, data formats are constrained and data values controlled through the provision of dropdown options, which reduces the number of processing steps necessary to prepare the raw data for publication. The data entry workflow for collecting aquatic cell count data as part of the water sampling is diagrammed in Figure 3. An additional set of constraints are implemented during the process of ingest into the NEON database. The product-specific data constraint and validation requirements built into data entry applications and database ingest are described in the document NEON Raw Data Validation for Aquatic Microbe Cell Counts (NEON.DP0.20138), provided with every download of this data product. Contained within this file is a field named 'entryValidationRulesForm', which describes syntactically the validation rules for each field built into the data entry application. Data entry constraints are described in NiCl syntax in the validation file provided with every data download, and the NiCl language is described in NEON's Ingest Conversion Language (NICL) specifications ([AD[11]).

5 DATA PROCESSING STEPS

Following data entry into a mobile application or web user interface, the steps used to process the data through to publication on the NEON Data Portal are detailed in the NEON Algorithm Theoretical Basis Document: OS Generic Transitions (AD[10]).

6 REFERENCES

Boulos, L., M. Prevost, B. Barbeau, J. Coallier, and R. Desjardins. 1999. LIVE/DEAD®*BacLight*TM: application of a new rapid staining method for direct enumeration of viable and total bacteria in drinking water. *Journal of Microbiological Methods* 37: 77-86.




Figure 3: Schematic of the applications used by field technicians to enter water chemistry and aquatic cell count field data